

Vectors

Last modified: 2026-10-10 11:17:20 (PDT)

Vectors

Exercise 0.1 (Store a data point as one object). A house has a floor area of 120 square meters, 3 bedrooms, and an age of 15 years.

1. Write these three numbers as a single vertical list $\tilde{x}$, in the order given, and call its entries x_1 , x_2 and x_3 . What is x_2 ?
2. Does the list $(3, 120, 15)$ describe the same house?

Solution

Solution 0.1.

1. Stacking the three numbers in order gives

$$\tilde{x} = \begin{bmatrix} 120 \\ 3 \\ 15 \end{bmatrix},$$

so $x_2 = 3$, the number of bedrooms.

2. No. The list $(3, 120, 15)$ would describe a house with 3 square meters of floor area and 120 bedrooms. An entry's position says what it measures, so the order of the entries matters.

Definition 0.1 (Column vector). A **column vector** of length p is an ordered list of p numbers, written vertically:

$$\tilde{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix}$$

Remark 0.1 (Vectors are columns by default). In these notes, as in most statistics textbooks, a vector is a column vector unless stated otherwise. A vector written inline as a list in parentheses is still a column vector: for example, $\tilde{x} = (2, -1, 5)$ means

$$\tilde{x} = \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix}.$$

Definition 0.2 (The set $\mathbb{R}^p$). For a positive integer p , $\mathbb{R}^p$ is the set of all column vectors of length p whose entries are real numbers.

Remark 0.2 (Reading $\tilde{x} \in \mathbb{R}^p$). Writing $\tilde{x} \in \mathbb{R}^p$ says, in one symbol, that $\tilde{x}$ is a column vector with p real entries. For example, $(1, -2, 0.5) \in \mathbb{R}^3$. The vector $(1, 2)$ is not in $\mathbb{R}^3$, because it has only 2 entries; it is in $\mathbb{R}^2$. In Exercise 0.1, $\tilde{x} \in \mathbb{R}^3$.

Definition 0.3 (Scalar). In linear algebra, a **scalar** is a single real number, as opposed to a vector or a matrix, which holds several numbers.

Example 0.1 (Scalars and a vector). The numbers 5 and -0.2 are scalars. The column vector $(5, -0.2)$ is not a scalar: it has 2 entries, so it is in $\mathbb{R}^2$ (Definition 0.2), and each of its 2 entries is a scalar.

Definition 0.4 (Real-valued (scalar-valued) and vector-valued functions). A function $f : A \rightarrow \mathbb{R}$, whose values are numbers, is **real-valued** (also called **scalar-valued**, since its values are scalars). A function $f : A \rightarrow \mathbb{R}^m$, whose values are vectors (Definition 0.2), is **vector-valued**.

Example 0.2 (A real-valued and a vector-valued function). On $\mathbb{R}^2$, $f(\tilde{x}) = x_1 + x_2$ is real-valued and $g(\tilde{x}) = \begin{bmatrix} x_1 + x_2 \\ x_1 - x_2 \end{bmatrix}$ is vector-valued:

$$f\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = 3 + 4 = 7, \quad g\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = \begin{bmatrix} 3 + 4 \\ 3 - 4 \end{bmatrix} = \begin{bmatrix} 7 \\ -1 \end{bmatrix}$$

Definition 0.5 (Transpose). The **transpose** of a column vector $\tilde{x}$ is the row vector with the same [sequence](#) of entries, written horizontally:

$$\tilde{x}^\top \equiv \tilde{x}' \equiv [x_1, x_2, \dots, x_p]$$

Example 0.3 (Transposing a column vector). The transpose of the column vector with entries 2, −1, 5 is the row vector with the same entries:

$$\begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix}^\top = [2, -1, 5].$$

Definition 0.6 (Vector addition). The **sum** of two column vectors $\tilde{x}$ and $\tilde{y}$ of the same length p is the column vector $\tilde{x} + \tilde{y}$ of length p obtained by adding entry by entry:

$$(\tilde{x} + \tilde{y})_i \stackrel{\text{def}}{=} x_i + y_i, \quad i = 1, \dots, p$$

Example 0.4 (Adding two vectors).

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 1 + 4 \\ 2 + 5 \\ 3 + 6 \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \\ 9 \end{bmatrix}$$

Definition 0.7 (Dot product). For any two real-valued vectors $\tilde{x} = (x_1, \dots, x_p)$ and $\tilde{y} = (y_1, \dots, y_p)$ of the same length p , the **dot product** of $\tilde{x}$ and $\tilde{y}$ is:

$$\tilde{x} \cdot \tilde{y} = \tilde{x}^\top \tilde{y} \stackrel{\text{def}}{=} \sum_{i=1}^p x_i y_i \tag{1}$$

See also the definitions in:

- Dobson and Barnett (2018), Section 1.3 (equation 1.1, page 7)

- Kaplan (2022), chapter on vectors
- [wikipedia](#)

The dot product has a different generalization for two matrices; see [wikipedia](#) for more.

Example 0.5 (A dot product). For $\tilde{x} = (1, 2, 3)$ and $\tilde{y} = (4, 5, 6)$:

$$\begin{aligned}\tilde{x} \cdot \tilde{y} &= 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6 && \text{(definition of the dot product)} \\ &= 4 + 10 + 18 && \text{(multiply)} \\ &= 32 && \text{(add)}\end{aligned}$$

Definition 0.8 (Linear combination). A **linear combination** of the numbers $a_1, \dots, a_k$ with **coefficients** $c_1, \dots, c_k$ is the weighted sum

$$c_1 a_1 + \dots + c_k a_k = \sum_{i=1}^k c_i a_i$$

A linear combination of vectors $\tilde{v}_1, \dots, \tilde{v}_k$ of the same length is defined entry by entry: entry j of $\sum_{i=1}^k c_i \tilde{v}_i$ is the linear combination of the j th entries of $\tilde{v}_1, \dots, \tilde{v}_k$ with the same coefficients. For example, the linear combination of $(1, 0)^\top$ and $(0, 1)^\top$ with coefficients 2 and 3 is $(2 \cdot 1 + 3 \cdot 0, 2 \cdot 0 + 3 \cdot 1)^\top = (2, 3)^\top$.

Remark 0.3 (The dot product is a linear combination). The dot product $\tilde{x} \cdot \tilde{y}$ is a linear combination (Definition 0.8) of the entries of $\tilde{y}$, with coefficients $x_1, \dots, x_p$. In Example 0.5, $\tilde{x} \cdot \tilde{y} = 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6$ is the linear combination of 4, 5, and 6 with coefficients 1, 2, and 3.

The dot product is also the standard *inner product* on $\mathbb{R}^p$; “inner product” is the general notion (Definition 1 in Inner Products and Orthogonality, later in these notes), of which the dot product is one example.

Theorem 0.1 (Dot product is symmetric). *The dot product is symmetric:*

$$\tilde{x} \cdot \tilde{y} = \tilde{y} \cdot \tilde{x}$$

i Proof

Proof.

$$\begin{aligned}\tilde{x} \cdot \tilde{y} &= \sum_{i=1}^p x_i y_i && \text{(definition of the dot product)} \\ &= \sum_{i=1}^p y_i x_i && \text{(products of numbers are symmetric)} \\ &= \tilde{y} \cdot \tilde{x} && \text{(definition of the dot product)}\end{aligned}$$

□

Special vectors

Definition 0.9 (Zero vector). The **zero vector** $\tilde{0}$ of length p has all entries equal to zero:

$$\tilde{0} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Remark 0.4 (The zero vector is the additive identity). Adding the zero vector to a vector leaves it unchanged (Definition 0.6): $\tilde{x} + \tilde{0} = \tilde{x}$ for any vector $\tilde{x}$ of the same length, so $\tilde{0}$ is the **identity element** for vector addition. For example,

$$\begin{aligned}(2, -1) + (0, 0) &= (2 + 0, -1 + 0) \\ &= (2, -1).\end{aligned}$$

Definition 0.10 (Mean (average, sample mean)). The **mean** (or **average**) of n numbers $y_1, \dots, y_n$ is their **sum** divided by how many there are:

$$\bar{y} \stackrel{\text{def}}{=} \frac{1}{n} \sum_{i=1}^n y_i$$

When the numbers are observed data values, $\bar{y}$ is also called the **sample mean**.

Example 0.6 (The mean of three numbers). For $y_1 = 2$, $y_2 = -1$ and $y_3 = 5$:

$$\begin{aligned}
\bar{y} &= \frac{1}{3} (2 + (-1) + 5) && \text{(definition of the mean, with } n = 3\text{)} \\
&= \frac{1}{3} \cdot 6 && \text{(add)} \\
&= 2 && \text{(multiply)}
\end{aligned}$$

Definition 0.11 (Ones vector). The **ones vector** $\tilde{\mathbf{1}}$ of length p has all entries equal to one:

$$\tilde{\mathbf{1}} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$$

Remark 0.5 (The ones vector sums the entries). The dot product (Definition 0.7) of $\tilde{\mathbf{1}}$ with $\tilde{x}$ is the sum of the entries of $\tilde{x}$:

$$\begin{aligned}
\tilde{\mathbf{1}} \cdot \tilde{x} &= \sum_{i=1}^p 1 \cdot x_i \\
&= \sum_{i=1}^p x_i.
\end{aligned}$$

For example, for $\tilde{x} = (2, -1, 5)$,

$$\begin{aligned}
\tilde{\mathbf{1}} \cdot \tilde{x} &= 2 + (-1) + 5 \\
&= 6,
\end{aligned}$$

and dividing by the length $p = 3$ gives the [mean](#) of the entries, $6/3 = 2$.

Definition 0.12 (Indicator vector (standard basis vector)). The j -th **indicator vector** (or **standard basis vector**) $\tilde{e}_j$ of length p has a 1 in position j and 0s elsewhere:

$$(\tilde{e}_j)_i = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \quad \tilde{e}_j = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \leftarrow \text{position } j$$

Remark 0.6 (Indicator vectors build every vector). For $p = 3$, the indicator vectors are $\tilde{e}_1 = (1, 0, 0)$, $\tilde{e}_2 = (0, 1, 0)$, and $\tilde{e}_3 = (0, 0, 1)$. Every vector of length 3 is a weighted sum $c_1\tilde{e}_1 + c_2\tilde{e}_2 + c_3\tilde{e}_3$ of them, with weights equal to its entries; for example, $(2, -1, 5) = 2\tilde{e}_1 - 1\tilde{e}_2 + 5\tilde{e}_3$. No other weights work: entry j of $c_1\tilde{e}_1 + c_2\tilde{e}_2 + c_3\tilde{e}_3$ is c_j , so each weight c_j must equal entry j of the vector.

Theorem 0.2 (Indicator vectors select entries). *For any vector $\tilde{x}$ of length p and any $j \in \{1, \dots, p\}$:*

$$\tilde{e}_j \cdot \tilde{x} = x_j$$

i Proof

Proof. Writing the dot product componentwise:

$$\begin{aligned} \tilde{e}_j \cdot \tilde{x} &= \sum_{i=1}^p (\tilde{e}_j)_i x_i && \text{(definition of the dot product)} \\ &= \sum_{i=1}^p \begin{cases} 1 \cdot x_i & \text{if } i = j \\ 0 \cdot x_i & \text{if } i \neq j \end{cases} && \text{(definition of } \tilde{e}_j) \\ &= x_j && \text{(only the } i = j \text{ term is nonzero)} \end{aligned}$$

□

Theorem 0.3 (A vector is a linear combination of indicator vectors). *Every vector $\tilde{x} \in \mathbb{R}^p$ is the linear combination (Definition 0.8) of the indicator vectors $\tilde{e}_1, \dots, \tilde{e}_p$ (Definition 0.12) with coefficients $x_1, \dots, x_p$:*

$$\tilde{x} = \sum_{j=1}^p x_j \tilde{e}_j$$

i Proof

Proof. Compare the two sides entry by entry. For each $i \in \{1, \dots, p\}$:

$$\begin{aligned}
\left(\sum_{j=1}^p x_j \tilde{e}_j\right)_i &= \sum_{j=1}^p x_j (\tilde{e}_j)_i && \text{(Definition 0.8, entry by entry)} \\
&= \sum_{j=1}^p \begin{cases} x_j \cdot 1 & \text{if } j = i \\ x_j \cdot 0 & \text{if } j \neq i \end{cases} && \text{(Definition 0.12)} \\
&= x_i \cdot 1 + \sum_{j \neq i} x_j \cdot 0 && \text{(split off the } j = i \text{ term)} \\
&= x_i + 0 && \text{(times 1 changes nothing; times 0 gives 0)} \\
&= x_i && \text{(adding 0 changes nothing)}
\end{aligned}$$

□

Example 0.7 (Expanding a vector in indicator vectors).

$$\begin{aligned}
\begin{bmatrix} 4 \\ -1 \\ 2 \end{bmatrix} &= 4 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + (-1) \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
&= 4\tilde{e}_1 - \tilde{e}_2 + 2\tilde{e}_3
\end{aligned}$$

Orthogonality

Definition 0.13 (Orthogonal vectors). Two vectors $\tilde{x}$ and $\tilde{y}$ of the same length are **orthogonal** (written $\tilde{x} \perp \tilde{y}$) if their dot product (Definition 0.7) is zero:

$$\tilde{x} \perp \tilde{y} \iff \tilde{x} \cdot \tilde{y} = 0$$

Remark 0.7 (Orthogonal means perpendicular). Orthogonality extends the geometric idea of perpendicular lines to vectors with any number of entries. In the plane, $(1, 2)$ and $(-2, 1)$ are perpendicular, and their dot product is $1 \cdot (-2) + 2 \cdot 1 = 0$. The same test works in $\mathbb{R}^3$, where pictures are harder to draw:

$$\begin{aligned}
(1, 1, 0) \cdot (1, -1, 5) &= 1 - 1 + 0 \\
&= 0,
\end{aligned}$$

so $(1, 1, 0) \perp (1, -1, 5)$.

Example 0.8 (Vectors that are not orthogonal).

$$\begin{aligned}(1, 2) \cdot (1, 1) &= 1 \cdot 1 + 2 \cdot 1 \\ &= 3 \\ &\neq 0,\end{aligned}$$

so $(1, 2)$ and $(1, 1)$ are not orthogonal.

Definition 0.14 (Euclidean norm (length, L_2 norm)). The **Euclidean norm** (or **length**, also called the L_2 **norm** and written $\|\tilde{x}\|_2$) of a vector $\tilde{x}$ of length p is

$$\|\tilde{x}\| \stackrel{\text{def}}{=} \sqrt{\tilde{x} \cdot \tilde{x}} = \sqrt{\sum_{i=1}^p x_i^2} \quad (2)$$

where $\sqrt{\cdot}$ is the [square root](#).

Example 0.9 (The length of a vector). For $\tilde{x} = (3, 4)$:

$$\begin{aligned}\|\tilde{x}\| &= \sqrt{3^2 + 4^2} \quad (\text{definition of the norm}) \\ &= \sqrt{25} \quad (\text{square and add}) \\ &= 5\end{aligned}$$

The vector $(0.6, 0.8)$ has norm $\sqrt{0.36 + 0.64} = 1$.

Definition 0.15 (Unit vector). A **unit vector** is a vector whose Euclidean norm (Definition [0.14](#)) is 1. A unit vector is said to have *unit length*.

Example 0.10 (A unit vector and a vector that is not one). For $\tilde{x} = (0.6, 0.8)$:

$$\begin{aligned}\|\tilde{x}\| &= \sqrt{0.6^2 + 0.8^2} \quad (\text{definition of the norm}) \\ &= \sqrt{0.36 + 0.64} \quad (\text{square}) \\ &= \sqrt{1} \quad (\text{add}) \\ &= 1 \quad (\text{take the square root})\end{aligned}$$

so $(0.6, 0.8)$ is a unit vector. The vector $(1, 1)$ is not:

$$\begin{aligned}\|(1, 1)\| &= \sqrt{1^2 + 1^2} \\ &= \sqrt{2} \\ &\neq 1.\end{aligned}$$

Exercise 0.2 (How far apart are two points?). Let $\tilde{x} = (1, 2)$ and $\tilde{y} = (4, 6)$.

1. Compute the vector $\tilde{x} - \tilde{y}$.
2. Compute the length of $\tilde{x} - \tilde{y}$, using the [Euclidean norm](#).
3. Is the result the same if you compute the length of $\tilde{y} - \tilde{x}$ instead?

Solution

Solution 0.2.

1.

$$\begin{aligned}\tilde{x} - \tilde{y} &= (1 - 4, 2 - 6) \\ &= (-3, -4).\end{aligned}$$

2.

$$\begin{aligned}\|\tilde{x} - \tilde{y}\| &= \sqrt{(-3)^2 + (-4)^2} \\ &= \sqrt{25} \\ &= 5.\end{aligned}$$

3. $\tilde{y} - \tilde{x} = (3, 4)$, and

$$\begin{aligned}\|\tilde{y} - \tilde{x}\| &= \sqrt{3^2 + 4^2} \\ &= 5.\end{aligned}$$

Yes, the result is the same.

Definition 0.16 (Euclidean distance). The **Euclidean distance** between two vectors $\tilde{x}$ and $\tilde{y}$ of length p is the [Euclidean norm](#) of their difference:

$$d(\tilde{x}, \tilde{y}) \stackrel{\text{def}}{=} \|\tilde{x} - \tilde{y}\| = \sqrt{\sum_{i=1}^p (x_i - y_i)^2} \quad (3)$$

Remark 0.8 (Distance from the origin). The distance from the origin $\tilde{0}$ to a vector $\tilde{x}$ is the norm of $\tilde{x}$:

$$\begin{aligned} d(\tilde{x}, \tilde{0}) &= \|\tilde{x} - \tilde{0}\| \\ &= \|\tilde{x}\| \end{aligned}$$

(Goodfellow et al. 2016, 39).

For example, the distance from $\tilde{0}$ to $(3, 4)$ is $\|(3, 4)\| = 5$, as in Example 0.9.

Definition 0.17 (Orthonormal vectors). A set of vectors $\{\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_k\}$ is **orthonormal** if the vectors are mutually orthogonal (Definition 0.13) and each has norm 1 (Definition 0.14), that is, **unit length**:

$$\tilde{x}_i \cdot \tilde{x}_j = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

Example 0.11 (Indicator vectors are orthonormal). The indicator vectors $\tilde{e}_1, \tilde{e}_2, \dots, \tilde{e}_p$ (Definition 0.12) form an orthonormal set. By Theorem 0.2, $\tilde{e}_i \cdot \tilde{e}_j$ is entry i of $\tilde{e}_j$, which is 1 if $i = j$ and 0 if $i \neq j$. For $p = 2$:

$$\begin{aligned} \tilde{e}_1 \cdot \tilde{e}_1 &= 1 \cdot 1 + 0 \cdot 0 \\ &= 1, \end{aligned}$$

$$\begin{aligned} \tilde{e}_2 \cdot \tilde{e}_2 &= 0 \cdot 0 + 1 \cdot 1 \\ &= 1, \end{aligned}$$

and

$$\begin{aligned} \tilde{e}_1 \cdot \tilde{e}_2 &= 1 \cdot 0 + 0 \cdot 1 \\ &= 0. \end{aligned}$$

Example 0.12 (Two ways to fail to be orthonormal).

- $(1, 1)$ and $(1, -1)$ are orthogonal, since $1 \cdot 1 + 1 \cdot (-1) = 0$, but not orthonormal: $(1, 1) \cdot (1, 1) = 2 \neq 1$.
- $(1, 0)$ and $(0.6, 0.8)$ both have unit length, since $(1, 0) \cdot (1, 0) = 1$ and

$$\begin{aligned} (0.6, 0.8) \cdot (0.6, 0.8) &= 0.36 + 0.64 \\ &= 1, \end{aligned}$$

but they are not orthonormal: $(1, 0) \cdot (0.6, 0.8) = 0.6 \neq 0$.

Exercise 0.3 (Compute an inner product and a norm). Let

$$\tilde{a} = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}, \quad \tilde{b} = \begin{bmatrix} 0 \\ 4 \\ -2 \end{bmatrix}$$

Compute $\tilde{a}^\top \tilde{b}$ and $\|\tilde{a}\|_2$.

Solution

Solution 0.3. Multiply entry by entry and add:

$$\begin{aligned} \tilde{a} \cdot \tilde{b} &= (3)(0) + (-1)(4) + (2)(-2) \\ &= 0 - 4 - 4 \\ &= -8 \end{aligned}$$

For the norm, take the inner product of $\tilde{a}$ with itself first:

$$\begin{aligned} \tilde{a} \cdot \tilde{a} &= 3^2 + (-1)^2 + 2^2 \\ &= 9 + 1 + 4 \\ &= 14 \end{aligned}$$

so

$$\|\tilde{a}\|_2 = \sqrt{14} \approx 3.742$$

The inner product came out negative while the norm cannot: a norm is a square root of a sum of squares, so it is never below zero.

Python has both operations built in, and R builds them from `sum`.

```
import numpy as np

a = np.array([3.0, -1.0, 2.0])
b = np.array([0.0, 4.0, -2.0])

print(a @ b, round(float(np.linalg.norm(a)), 3))
```

```
a <- c(3, -1, 2)
b <- c(0, 4, -2)

print(c(sum(a * b), round(sqrt(sum(a^2)), 3)))
#> [1] -8.000 3.742
```

Both print -8 and 3.742 .

Video lecture

Hutchinson's [Linear Function Basics](#) (24 min) covers dot products and the Euclidean norm, and goes on to hyperplanes ([Definition 19 in Matrices](#)) and [level sets](#) (Hutchinson, n.d.). The login for the video site is posted [on Canvas](#).

Dobson, Annette J, and Adrian G Barnett. 2018. *An Introduction to Generalized Linear Models*. 4th ed. CRC press. <https://doi.org/10.1201/9781315182780>.

Goodfellow, Ian, Yoshua Bengio, and Aaron Courville. 2016. *Deep Learning*. MIT Press. <https://www.deeplearningbook.org/>.

Hutchinson, Brian. n.d. *DATA 471/571 (Machine Learning) and CSCI 481/581 (Deep Learning) Video Lectures*. Western Washington University. Accessed September 28, 2026. https://facultyweb.cs.wvu.edu/~hutchib2/video_lectures/data371/.

Kaplan, Daniel. 2022. *MOSAIC Calculus*. Wwww.mosaic-web.org. www.mosaic-web.org.