Causal Inference

Estimating causal effects from regression models

Contents

Configuring R 1

Acknowledgements **2**

1 Introduction **2**
1.1 Descriptive versus causal questions 2

2 Potential Outcomes **4**

3 Causal Estimands **5**

4 Causal Assumptions **6**
4.1 Consistency 7
4.2 Exchangeability (No Unmeasured Confounding) 7
4.3 Positivity 8

5 Randomized Experiments **9**
5.1 The HERS study: a randomized trial example 10
5.2 Covariate adjustment in RCTs 11

6 Observational Studies **11**
6.1 Confounding 11
6.2 Why observational studies are harder than randomized trials 12
6.3 Directed Acyclic Graphs (DAGs) 13
6.4 Key concepts from DAG theory 13
6.5 HERS study DAG 14
6.6 What variables should we adjust for? 17
6.7 Analyzing LDL in the HERS study 17
6.7.1 Unadjusted model 18
6.7.2 Adjusted model 18
6.8 An observational study example: WCGS 19

7 Regression Adjustment **21**
7.1 Direct standardization (G-computation) 22
7.2 Regression adjustment in linear models 23
7.3 Example: regression adjustment for LDL in HERS 23

8 Propensity Score Methods **24**
8.1 Propensity score estimation 25
8.2 Propensity score methods 25
8.3 Propensity score example: WCGS 27

9 Summary **28**

Learning Objectives **29**

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
library(broom) # Summarizes key information about statistical objects in tidy tibbles
library(broom.helpers) # Provides suite of functions to work with regression model 'broom::tidy()' t
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R

conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
  ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
    ggplot2::theme(
      legend.position = "bottom",
      text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 6)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by default
legend_text_size = 9
run_graphs = TRUE
```

Acknowledgements

This chapter is adapted from [1, Ch. 9] and draws on the companion causal inference notes available at https://d-morrison.github.io/cie/.

See also Hernán and Robins [2] for a comprehensive treatment of causal inference from observational data.

1 Introduction

Regression models[1] describe statistical associations between variables. But in many research settings, the goal is not merely to describe associations but to estimate **causal effects**: what would happen if we intervened to change a variable?

"No causes in, no causes out." [3]

A statistical model alone cannot tell us whether an observed association is causal or spurious. Causal inference requires additional assumptions — about the data-generating process — that must be justified using subject-matter knowledge.

The causal inference framework covered in this chapter is based on the **potential outcomes** model [3], [4]. This framework provides:

- A precise language for defining causal effects
- Explicit assumptions required to identify causal effects from data
- Estimation methods that target specific causal estimands

For more resources on causal inference, including additional courses and books, see https://d-morrison.github.io/cie/.

1.1 Descriptive versus causal questions

Before turning to causal questions, it helps to separate them from descriptive ones.

Suppose we want to know how heavy a cow will be if it eats grass all summer. This question is a descriptive question about the conditional distribution of weight given diet, $\mathrm{p}(Y = y \mid X = \text{grass})$, where Y is weight and X is diet. We have not asked how heavy a cow would be if we *intervened* to make it eat grass all summer.

Suppose we find some cows eating grass and others eating hay. We can observe the weights of the grass-fed cows and the weights of the hay-fed cows, but we cannot observe what would have happened if we had made the grass-fed cows eat hay instead. Would they have weighed the same as the cows we found eating hay? Several things could make the answer no:

- the two groups might differ in breed;
- the way the cows ended up with their diets might matter: perhaps the bigger cows chased the smaller ones off the grass.

We can answer causal questions of this kind, but we generally need to answer the descriptive questions first. Descriptive questions do not require covariates[2] that we *could* intervene on; they only require covariates that we can observe. This hypothetical example illustrates the distinction that the potential outcomes framework of this chapter makes precise.

The same distinction arises with binary outcomes, where the descriptive summary is often a **risk difference**: the difference between the proportions of two groups who experience the outcome.

Exm

Example 1.1 (A crude risk difference that reverses within age groups)**.** Consider a hypothetical population of 1000 adults, with a binary treatment A ($A = 1$ for treated), a binary outcome Y

[1] Intro-to-GLMs.qmd#def-regression-model
[2] Intro-to-GLMs.qmd#def-predictors

($Y = 1$ for death within a year), and a binary covariate L ($L = 1$ for age 65 or older). Older adults are more likely to be treated, and older adults are also more likely to die, whether or not they are treated:

L	treated	untreated	$\mathrm{P}(Y = 1 \mid A = 1, L)$	$\mathrm{P}(Y = 1 \mid A = 0, L)$
0 (younger)	100	500	0.05	0.10
1 (older)	300	100	0.30	0.40

Within each age group, the treated have lower risk than the untreated:

$$
\begin{aligned}
\mathrm{RD}_{L=0} &\overset{\text{def}}{=} \mathrm{P}(Y = 1 \mid A = 1, L = 0) - \mathrm{P}(Y = 1 \mid A = 0, L = 0) && \text{(define the age-specific risk difference)} \\
&= 0.05 - 0.10 && \text{(substitute from the table)} \\
&= -0.05 && \text{(subtract)} \\
\mathrm{RD}_{L=1} &\overset{\text{def}}{=} \mathrm{P}(Y = 1 \mid A = 1, L = 1) - \mathrm{P}(Y = 1 \mid A = 0, L = 1) && \text{(define the age-specific risk difference)} \\
&= 0.30 - 0.40 && \text{(substitute from the table)} \\
&= -0.10 && \text{(subtract)}
\end{aligned}
$$

Ignoring age, the risk in each treatment group averages the age-specific risks in proportion to that group's own age distribution:

$$
\begin{aligned}
\mathrm{P}(Y = 1 \mid A = 1) &= \frac{100 \cdot 0.05 + 300 \cdot 0.30}{400} && \text{(average over the 400 treated)} \\
&= \frac{5 + 90}{400} && \text{(multiply)} \\
&= \frac{95}{400} && \text{(add)} \\
&= 0.2375 && \text{(divide)} \\
\mathrm{P}(Y = 1 \mid A = 0) &= \frac{500 \cdot 0.10 + 100 \cdot 0.40}{600} && \text{(average over the 600 untreated)} \\
&= \frac{50 + 40}{600} && \text{(multiply)} \\
&= \frac{90}{600} && \text{(add)} \\
&= 0.15 && \text{(divide)} \\
\mathrm{RD}_{\text{crude}} &\overset{\text{def}}{=} \mathrm{P}(Y = 1 \mid A = 1) - \mathrm{P}(Y = 1 \mid A = 0) && \text{(define the crude risk difference)} \\
&= 0.2375 - 0.15 && \text{(substitute)} \\
&= 0.0875 && \text{(subtract)}
\end{aligned}
$$

The crude risk difference of $+0.0875$ is a correct *descriptive* answer: in this population, treated adults really did die more often than untreated adults. It is a misleading answer to the *causal* question "would treating someone raise or lower their risk?", because 75% of the treated group is older, compared with about 17% of the untreated group. If age were the only factor that influenced both who was treated and who died, the age-specific risk differences (-0.05 and -0.10) would compare like with like, and they say that treatment lowers risk. The rest of this chapter makes precise when a comparison of this kind has a causal interpretation.

2 Potential Outcomes

Causal inference is about estimating the effect of interventions — what would happen if we changed something? Statistical associations alone do not answer causal questions.

The fundamental challenge in causal inference is that we can only observe one potential outcome for each unit: the outcome under the treatment that actually occurred. This unobservability of counterfactual outcomes is called the **fundamental problem of causal inference** [3].

Definition 2.1 (Potential outcomes (counterfactual outcomes))**.** For a binary treatment $A \in \{0, 1\}$, the **potential outcomes** for unit i are:

- $Y_i(1)$: the outcome that would be observed if unit i received treatment ($A_i = 1$)
- $Y_i(0)$: the outcome that would be observed if unit i received control ($A_i = 0$)

These potential outcomes are also called **counterfactual outcomes**, because for each unit, one of them is necessarily counterfactual (it corresponds to a treatment that did not occur).

Definition 2.2 (Observed outcome)**.** The **observed outcome** for unit i is the potential outcome corresponding to the treatment actually received:

$$Y_i \stackrel{\text{def}}{=} Y_i(A_i) = A_i \cdot Y_i(1) + (1 - A_i) \cdot Y_i(0)$$

Remark 2.1 (Observed and potential outcomes: the consistency assumption)*.* This relationship between observed and potential outcomes is sometimes called the **consistency** assumption: the observed outcome under the observed treatment equals the potential outcome under that treatment.

Definition 2.3 (Individual causal effect)**.** The **individual causal effect** for unit i is the difference between that unit's two potential outcomes:

$$\tau_i \stackrel{\text{def}}{=} Y_i(1) - Y_i(0)$$

Exm

Example 2.1 (Potential outcomes and observed outcomes in a toy sample)**.** Consider a sample[a] of three individuals being evaluated for a medication ($A \in \{0, 1\}$) where the outcome[b] Y is blood pressure reduction (mmHg):

Unit i	$Y_i(0)$	$Y_i(1)$	$\tau_i = Y_i(1) - Y_i(0)$	Observed A_i	Observed Y_i
1	5	15	+10	1	15
2	2	8	+6	0	2
3	10	12	+2	1	12

[a] https://morrison-lab.github.io/sds/exploratory-descriptive.html#def-sample
[b] Intro-to-GLMs.qmd#def-outcome

For unit 1 ($A_1 = 1$), we observe $Y_1 = Y_1(1) = 15$, while $Y_1(0) = 5$ is unobserved (counterfactual). For unit 2 ($A_2 = 0$), we observe $Y_2 = Y_2(0) = 2$, while $Y_2(1) = 8$ is counterfactual. Because only one potential outcome is observed per unit, we cannot directly calculate τ_i for any individual in real data.

Remark 2.2 (Individual causal effects are not identified)*.* Because of the fundamental problem of causal inference, we can never observe both $Y_i(1)$ and $Y_i(0)$ for the same unit, so individual causal effects are generally not identified.

3 Causal Estimands

Since individual causal effects are not identified, causal inference focuses on **population-level causal estimands**[3]: averages of potential outcomes over a population[4].

Definition 3.1 (Average Treatment Effect (ATE))**.** The **Average Treatment Effect** is the expected[a] difference in potential outcomes in the study population:

$$\text{ATE} \stackrel{\text{def}}{=} \mathrm{E}[Y(1) - Y(0)] = \mathrm{E}[Y(1)] - \mathrm{E}[Y(0)]$$

[a] https://morrison-lab.github.io/pds/expectation.html#def-expectation

Remark 3.1 (Interpretation of the ATE). The ATE represents the average causal effect of treatment compared with control, averaged over all units in the population (both those who would and would not actually receive treatment).

Definition 3.2 (Average Treatment Effect on the Treated (ATT))**.** The **Average Treatment Effect on the Treated** is the expected difference in potential outcomes among units who actually received treatment:

$$\text{ATT} \stackrel{\text{def}}{=} \mathrm{E}[Y(1) - Y(0) \mid A = 1] = \mathrm{E}[Y(1) \mid A = 1] - \mathrm{E}[Y(0) \mid A = 1]$$

Remark 3.2 (Interpretation of the ATT). The ATT answers the question: "On average, how much did treatment help the people who received it?"

Definition 3.3 (Average Treatment Effect on the Untreated (ATU))**.** The **Average Treatment Effect on the Untreated** is the expected difference in potential outcomes among units who did not receive treatment:

$$\text{ATU} \stackrel{\text{def}}{=} \mathrm{E}[Y(1) - Y(0) \mid A = 0] = \mathrm{E}[Y(1) \mid A = 0] - \mathrm{E}[Y(0) \mid A = 0]$$

Exm

Example 3.1 (Calculating population-level causal estimands in a toy population)**.** Revisiting the three-unit population from Example 2.1:

- Individual 1 ($A_1 = 1$): $Y_1(0) = 5$, $Y_1(1) = 15$, $\tau_1 = +10$.
- Individual 2 ($A_2 = 0$): $Y_2(0) = 2$, $Y_2(1) = 8$, $\tau_2 = +6$.
- Individual 3 ($A_3 = 1$): $Y_3(0) = 10$, $Y_3(1) = 12$, $\tau_3 = +2$.

We calculate the population-level estimands across all three units:

1. **ATE**: $\frac{\tau_1+\tau_2+\tau_3}{3} = \frac{10+6+2}{3} = +6.0$.
2. **ATT**: Average over treated units ($i = 1, 3$): $\frac{\tau_1+\tau_3}{2} = \frac{10+2}{2} = +6.0$.
3. **ATU**: Average over untreated units ($i = 2$): $\frac{\tau_2}{1} = +6.0$.

In this toy population, treatment effect heterogeneity exists (τ_i varies across units), but because average gains happen to balance across treatment groups, ATE = ATT = ATU. In general observational datasets with selection bias, ATT and ATU often differ markedly from ATE.

Remark 3.3 (Interpretation of the ATU). The ATU answers the question: "On average, how much would treatment have helped the people who did not receive it?"

The ATE, ATT, and ATU are equal when treatment assignment is independent[5] of potential

[3] https://morrison-lab.github.io/sds/estimation.html#def-estimand
[4] https://morrison-lab.github.io/sds/exploratory-descriptive.html#def-population
[5] https://morrison-lab.github.io/pds/independence.html#def-indpt

outcomes (e.g., in a perfectly conducted randomized trial):

$$\text{ATE} = \text{ATT} = \text{ATU} \quad \text{when } A \perp\!\!\!\perp (Y(0), Y(1))$$

In observational studies, these three estimands can differ substantially. The choice of estimand depends on the research question.

4 Causal Assumptions

Three key assumptions are required to identify causal effects from observed data [2]:

4.1 Consistency

Definition 4.1 (Consistency)**.** The **consistency** assumption states that the observed outcome for a treated unit equals that unit's potential outcome under treatment, and similarly for control:

$$Y_i \overset{\text{def}}{=} Y_i(A_i)$$

Remark 4.1 (Well-defined treatment and violations of consistency). Consistency requires that the treatment is well-defined: there is only one version of treatment, or different versions have the same effect on the outcome. Violations of consistency (e.g., different doses of a drug being lumped together as "treated") can introduce bias[a].

[a] https://morrison-lab.github.io/sds/estimation.html#def-bias

4.2 Exchangeability (No Unmeasured Confounding)

Definition 4.2 (Exchangeability (no unmeasured confounding, ignorability))**. Exchangeability** (also called **no unmeasured confounding** or **ignorability**) states that, conditional on observed covariates $\underset{\sim}{L}$, treatment assignment is independent[a] of potential outcomes:

$$A \perp\!\!\!\perp (Y(0), Y(1)) \mid \underset{\sim}{L}$$

[a] https://morrison-lab.github.io/pds/independence.html#def-cind

Remark 4.2 (Interpretation of exchangeability). Intuitively, exchangeability means that among individuals with the same covariate values, treatment was assigned as if at random — there are no unmeasured common causes of treatment and outcome.

Definition 4.3 (Unconditional exchangeability)**. Unconditional exchangeability** ($A \perp\!\!\!\perp (Y(0), Y(1))$) holds in perfectly randomized trials. Exchangeability conditional on $\underset{\sim}{L}$ (Definition 4.2) requires adjusting for $\underset{\sim}{L}$.

Exm

Example 4.1 (A population where exchangeability holds only within strata)**.** Consider a large population of 1000 patients, with a binary covariate L (disease severity: $L = 1$ for severe) and treatment A. Severe patients are more likely to be treated, and both potential outcomes are higher for severe patients (treating proportions as probabilities):

L	patients	$\text{P}(A = 1 \mid L)$	treated	untreated	$\text{E}[Y(1) \mid L]$	$\text{E}[Y(0) \mid L]$
0	600	0.2	120	480	10	8

1	400	0.7	280	120	20	18

Within each severity stratum, suppose treatment was assigned as if at random, so that $A \perp\!\!\!\perp (Y(0), Y(1)) \mid L$: conditional exchangeability (Definition 4.2) holds with $\underset{\sim}{L} = L$. Unconditional exchangeability (Definition 4.3) fails, though. Averaging $Y(0)$ over the strata in proportion to each treatment group's members:

$$\begin{aligned}
\mathrm{E}[Y(0) \mid A = 1] &= \frac{120 \cdot 8 + 280 \cdot 18}{400} && \text{(average over the 400 treated)} \\
&= \frac{960 + 5040}{400} && \text{(multiply)} \\
&= 15 && \text{(add and divide)} \\
\mathrm{E}[Y(0) \mid A = 0] &= \frac{480 \cdot 8 + 120 \cdot 18}{600} && \text{(average over the 600 untreated)} \\
&= \frac{3840 + 2160}{600} && \text{(multiply)} \\
&= 10 && \text{(add and divide)}
\end{aligned}$$

The treated group would have had a higher mean outcome even without treatment, because it contains more severe patients, so A is not independent of $Y(0)$.

4.3 Positivity

Definition 4.4 (Positivity)**.** The **positivity** assumption states that every individual has a positive probability of receiving each treatment level, at every covariate value $\underset{\sim}{l}$ in the support[a] of $\underset{\sim}{L}$:

$$0 < \mathrm{P}(A = 1 \mid \underset{\sim}{L} = \underset{\sim}{l}) < 1 \quad \text{for all } \underset{\sim}{l} \text{ in the support of } \underset{\sim}{L}$$

[a] https://morrison-lab.github.io/mds/indicator-functions.html#def-support

Remark 4.3 (Why positivity is needed). Without positivity, some subgroups have no treated (or untreated) members, making it impossible to estimate the counterfactual outcome for those subgroups.

Definition 4.5 (Positivity violation)**.** A failure of positivity is sometimes called a **positivity violation**. When near-zero probabilities occur in small strata, the failure is called a practical positivity violation.

Exm

Example 4.2 (Checking positivity)**.** In the population of Example 4.1, $\mathrm{P}(A = 1 \mid L = 0) = 0.2$ and $\mathrm{P}(A = 1 \mid L = 1) = 0.7$. Both probabilities lie strictly between 0 and 1, so positivity (Definition 4.4) holds: each severity stratum contains both treated and untreated patients. If instead every severe patient were treated, then $\mathrm{P}(A = 1 \mid L = 1) = 1$, the population would contain no untreated severe patients, and the data could not tell us what $\mathrm{E}[Y(0) \mid L = 1]$ is: a positivity violation (Definition 4.5).

These three assumptions — consistency, exchangeability, and positivity — are collectively sufficient to identify causal effects from observed data. They are sometimes summarized as "no interference, no unmeasured confounders, no positivity violations."

These assumptions are **untestable** from the data alone; they must be justified using subject-matter knowledge.

5 Randomized Experiments

In a **randomized controlled trial** (RCT), treatment is assigned by the investigator using a random mechanism, independent[6] of any characteristics of the participants. Randomization makes the treatment groups **exchangeable**: the distribution of potential outcomes is the same in treated and untreated groups.

As a result, in a perfectly randomized trial, the observed group means identify the mean potential outcomes:

Theorem 5.1 (Identification of the ATE under randomization)**.** *If consistency and unconditional exchangeability hold, and* $0 < \mathrm{P}(A = 1) < 1$*, then for each treatment level* $a \in \{0, 1\}$*,*

$$\mathrm{E}[Y(a)] = \mathrm{E}[Y \mid A = a]$$

and therefore

$$ATE = \mathrm{E}[Y \mid A = 1] - \mathrm{E}[Y \mid A = 0] \tag{1}$$

Proof

Proof. For treatment level a:

$$\begin{aligned} \mathrm{E}[Y(a)] &= \mathrm{E}[Y(a) \mid A = a] && \text{(unconditional exchangeability: } Y(a) \perp\!\!\!\perp A) \\ &= \mathrm{E}[Y \mid A = a] && \text{(consistency: } Y = Y(a) \text{ when } A = a) \end{aligned}$$

where the condition $0 < \mathrm{P}(A = 1) < 1$ makes both conditioning events $\{A = 1\}$ and $\{A = 0\}$ have positive probability. Substituting into the ATE:

$$\begin{aligned} \mathrm{ATE} &= \mathrm{E}[Y(1)] - \mathrm{E}[Y(0)] && \text{(definition of ATE)} \\ &= \mathrm{E}[Y \mid A = 1] - \mathrm{E}[Y \mid A = 0] && \text{(apply the first result with } a = 1 \text{ and } a = 0) \end{aligned}$$

□

Exm

Example 5.1 (Randomizing the severity example)**.** Suppose the 1000 patients of Example 4.1 had instead been randomized, each treated with probability 0.5 regardless of severity. Then each treatment group has the population's severity mix, 60% non-severe and 40% severe (treating proportions as probabilities), so by consistency

$$\begin{aligned} \mathrm{E}[Y \mid A = 1] &= 0.6 \cdot 10 + 0.4 \cdot 20 && \text{(severity mix of the treated; } \mathrm{E}[Y(1) \mid L]) \\ &= 14 && \text{(multiply and add)} \\ \mathrm{E}[Y \mid A = 0] &= 0.6 \cdot 8 + 0.4 \cdot 18 && \text{(severity mix of the untreated; } \mathrm{E}[Y(0) \mid L]) \\ &= 12 && \text{(multiply and add)} \end{aligned}$$

and Equation 1 gives $\mathrm{ATE} = 14 - 12 = 2$, the treatment's effect in every stratum.

Exm

Example 5.2 (Without exchangeability, the naive contrast is biased)**.** In the population of Example 4.1, treatment was not randomized, so Theorem 5.1 does not apply. The observed group means are

[6] https://morrison-lab.github.io/pds/independence.html#def-indpt

$$
\begin{aligned}
\mathrm{E}[Y \mid A=1] &= \frac{120 \cdot 10 + 280 \cdot 20}{400} && \text{(consistency; average over the treated)} \\
&= \frac{1200 + 5600}{400} && \text{(multiply)} \\
&= 17 && \text{(add and divide)} \\
\mathrm{E}[Y \mid A=0] &= 10 && \text{(consistency; } \mathrm{E}[Y(0) \mid A=0] = 10 \text{ in that population)}
\end{aligned}
$$

so the naive contrast Equation 1 would give $17 - 10 = 7$. But the treatment raises each patient's mean outcome by exactly 2 in both strata ($10 - 8 = 20 - 18 = 2$), so the true ATE is 2. The extra 5 units come from the severe patients' over-representation among the treated: confounding by L.

In a randomized trial, therefore, the observed mean difference between treatment groups is an unbiased[7] estimate of the ATE:

$$\widehat{\mathrm{ATE}} = \bar{Y}_{A=1} - \bar{Y}_{A=0}$$

Randomization achieves unconditional exchangeability: $A \perp\!\!\!\perp (Y(0), Y(1))$. Because there are no common causes of treatment assignment and the outcome, there is no confounding. The observed association between A and Y in the data is a direct measure of the causal effect.

5.1 The HERS study: a randomized trial example

The "heart and estrogen/progestin study" (HERS) was a clinical trial of hormone therapy for prevention of recurrent heart attacks and death among 2,763 post-menopausal women with existing coronary heart disease (CHD) [5].

The trial was conducted at 20 US clinical centers. Participants were randomized to receive either conjugated equine estrogens (0.625 mg/day) plus medroxyprogesterone acetate (2.5 mg/day) or a matching placebo [5]. Women were followed for a sample mean of 4.1 years [5].

The primary outcome was nonfatal myocardial infarction or CHD death [5].

In HERS, hormone therapy (HT) was randomized, so the treatment groups were exchangeable at baseline. The observed difference in CHD event rates between the HT and placebo groups estimates the causal effect of HT.

Despite observational studies suggesting up to 35–80% fewer recurrent CHD events among hormone users than non-users, [5], HERS found **no significant**[8] **difference** in CHD event rates between the hormone therapy (HT) and placebo groups (relative hazard[9] 0.99; 95% CI[10]: 0.80–1.22) [5].

The HERS investigators noted a statistically significant time trend: more CHD events occurred in the HT group in year 1, while fewer occurred in years 4 and 5, suggesting that HT may have early harmful effects that are later reversed.

A key finding was that HT significantly altered lipid profiles:

- LDL cholesterol: 11% lower in the HT group [5].
- HDL cholesterol: 10% higher in the HT group [5].

Yet these favorable lipid changes did not translate into fewer CHD events overall. One explanation consistent with the DAG in Figure 1 is that the effects of HT on lipids (a mediator pathway) were offset by other, harmful pathways — possibly thrombotic — that are not captured in the lipid mediators.

[7] https://morrison-lab.github.io/sds/estimation.html#def-unbiased
[8] https://morrison-lab.github.io/sds/inference.html#def-statistically-significant
[9] intro-to-survival-analysis.qmd#def-hazard-ratio
[10] https://morrison-lab.github.io/sds/inference.html#def-confidence-interval

Hulley et al. [5] also noted that HT increased the rate of venous thromboembolic events (relative hazard 2.89; 95% CI: 1.50–5.58) and gallbladder disease (relative hazard 1.38; 95% CI: 1.00–1.92).

The HERS result illustrates the gap between observational evidence and causal evidence: prior observational studies had suggested large benefits of HT, but these reflected confounding by the "healthy user" effect — women who voluntarily used HT tended to be healthier overall. Randomization in HERS removed this confounding.

5.2 Covariate adjustment in RCTs

Although randomization ensures no confounding, adjusting for baseline covariates[11] in an RCT can still be useful:

- **Improved precision**[12]: Adjusting for strong predictors of the outcome reduces residual variance and narrows confidence intervals[13].
- **Covariate imbalance**: When randomization produces chance imbalances in small trials, adjustment can correct for these baseline differences.

However, in an RCT, covariates that are **mediators** of the treatment effect (i.e., variables affected by treatment) should generally **not** be adjusted for when estimating the **total** causal effect of treatment. Adjusting for mediators blocks part of the causal pathway and gives only the **direct** effect of treatment.

6 Observational Studies

Video lecture

Richard McElreath's *Elemental Confounds I*[a] (Statistical Rethinking 2026, Lecture A06) accompanies Chapter 6 of McElreath [6], which uses directed acyclic graphs (DAGs) to describe confounding [7].

[a] https://www.youtube.com/watch?v=lGR7D45Ww38

In an **observational study**, treatment is not assigned by the investigator. Instead, individuals self-select into treatment, or treatment is assigned based on clinical or administrative criteria. As a result, treatment groups may differ systematically in ways that also affect the outcome[14] — this systematic difference is called **confounding**.

6.1 Confounding

Definition 6.1 (Confounding)**. Confounding** occurs when there is a common cause of both the treatment A and the outcome Y. This common cause is called a **confounder**.

Remark 6.1 (Confounding in the potential outcomes framework)*.* In the potential outcomes framework, confounding means that treatment assignment is not independent[a] of potential outcomes:

$$A \not\perp\!\!\!\perp (Y(0), Y(1))$$

As a result, the observed association between A and Y does not equal the causal effect of A on Y.

[a] https://morrison-lab.github.io/pds/independence.html#def-indpt

[11] Intro-to-GLMs.qmd#def-predictors
[12] https://morrison-lab.github.io/pds/variance-covariance.html#def-precision
[13] https://morrison-lab.github.io/sds/inference.html#def-confidence-interval
[14] Intro-to-GLMs.qmd#def-outcome

Exm

Example 6.1 (Confounding of coffee and lung cancer by smoking)**.** Consider a hypothetical cohort of 1000 adults, with coffee drinking A ($A = 1$ for daily coffee), lung cancer Y ($Y = 1$ for diagnosis during follow-up), and smoking L ($L = 1$ for smokers). Suppose coffee has **no** effect on lung cancer, so that $Y_i(1) = Y_i(0)$ for every individual, but smokers are both more likely to drink coffee and more likely to develop lung cancer:

L	coffee	no coffee	$\mathrm{P}(Y = 1 \mid A = 1, L)$	$\mathrm{P}(Y = 1 \mid A = 0, L)$
0 (nonsmokers)	200	500	0.01	0.01
1 (smokers)	200	100	0.10	0.10

Within each smoking stratum, the risk difference is 0, matching the true (null) causal effect. Ignoring smoking:

$$
\begin{aligned}
\mathrm{P}(Y = 1 \mid A = 1) &= \frac{200 \cdot 0.01 + 200 \cdot 0.10}{400} && \text{(average over the 400 coffee drinkers)} \\
&= \frac{2 + 20}{400} && \text{(multiply)} \\
&= \frac{22}{400} && \text{(add)} \\
&= 0.055 && \text{(divide)} \\
\mathrm{P}(Y = 1 \mid A = 0) &= \frac{500 \cdot 0.01 + 100 \cdot 0.10}{600} && \text{(average over the 600 non-drinkers)} \\
&= \frac{5 + 10}{600} && \text{(multiply)} \\
&= \frac{15}{600} && \text{(add)} \\
&= 0.025 && \text{(divide)} \\
\mathrm{P}(Y = 1 \mid A = 1) - \mathrm{P}(Y = 1 \mid A = 0) &= 0.055 - 0.025 && \text{(substitute)} \\
&= 0.03 && \text{(subtract)}
\end{aligned}
$$

The crude risk difference of 0.03 suggests that coffee is harmful, even though coffee has no effect in this cohort. Because $Y_i(0) = Y_i(1) = Y_i$ for everyone here, the crude risks are also the mean potential outcomes under no coffee in each group: $\mathrm{E}[Y(0) \mid A = 1] = 0.055 \neq 0.025 = \mathrm{E}[Y(0) \mid A = 0]$. So $A \not\perp\!\!\!\perp Y(0)$, which is the failure of unconditional exchangeability described in Remark 6.1. Smoking L is a confounder: it is a common cause of A and Y.

6.2 Why observational studies are harder than randomized trials

In a randomized experiment, the investigator controls treatment assignment, so the identification assumptions hold by design [2, Ch. 3]:

- randomization makes the groups exchangeable without conditioning on any covariates (Definition 4.3);
- the randomization probabilities are strictly between 0 and 1, so positivity holds;
- the protocol specifies the treatment, which supports consistency.

In an observational study, none of these assumptions is guaranteed by design [2, Ch. 3]:

- exchangeability (Definition 4.2) can hold at best conditionally on measured covariates $\underset{\sim}{L}$, so we must measure and adjust for a sufficient set of confounders, and we cannot check from the data that no confounder is missing;
- positivity (Definition 4.4) can fail when some covariate patterns, such as the most severe patients, are always (or never) treated;
- consistency (Definition 4.1) requires a well-defined treatment, which is harder to ensure when

"treatment" is a self-chosen exposure, such as "drinks coffee" in Example 6.1, that people can take up in many different ways.

Directed Acyclic Graphs (DAGs) help with the first of these difficulties, by showing which covariates must be in $\underset{\sim}{L}$. More generally, DAGs provide a formal graphical tool for representing causal assumptions and identifying confounders, mediators, colliders, and sufficient adjustment sets.

6.3 Directed Acyclic Graphs (DAGs)

A **Directed Acyclic Graph (DAG)** is a graphical model that encodes assumed causal relationships among variables.

In a DAG:

- **Nodes** represent variables.
- **Directed edges** (arrows) represent direct causal effects.
- The graph is **acyclic**: there are no directed cycles (a variable cannot cause itself, even indirectly).

DAGs are used in causal inference to:

- Identify **confounders** that must be adjusted for.
- Identify **mediators** that should generally *not* be adjusted for when estimating the total causal effect.
- Identify **colliders** that should *not* be adjusted for (adjusting for a collider opens a non-causal path).

6.4 Key concepts from DAG theory

Definition 6.2 (Confounder)**.** A **confounder** of the relationship between an exposure X and outcome[a] Y is a variable C that is a common cause of both X and Y.

[a] Intro-to-GLMs.qmd#def-outcome

Remark 6.2 (Confounders compared with mediators). Unlike a confounder (Definition 6.2), a mediator lies on the causal path from X to Y and is not a common cause of X and Y. Adjusting for confounders removes confounding bias.

Definition 6.3 (Mediator)**.** A **mediator** M is a variable that lies on the causal pathway from the exposure X to the outcome Y:

$$X \to M \to Y$$

Remark 6.3 (Adjusting for a mediator). Adjusting for a mediator blocks the indirect causal pathway through M, so the estimated coefficient for X reflects only the *direct* effect of X on Y not through M.
Whether to adjust for mediators depends on the research question:

- If you want the **total causal effect** of X on Y, do **not** adjust for M.
- If you want the **direct effect** of X on Y not through M, adjust for M.

Definition 6.4 (Collider)**.** A **collider** is a variable L that is caused by two or more variables on a path:

$$X \to L \leftarrow Y$$

Remark 6.4 (Adjusting for a collider). Adjusting for a collider opens a non-causal (backdoor) path between X and Y.

Definition 6.5 (Collider bias)**.** Adjusting for a collider (Definition 6.4) introduces **collider bias**: the bias from the non-causal path that the adjustment opens between X and Y.

Definition 6.6 (Backdoor criterion)**.** A set of variables $\underset{\sim}{L}$ satisfies the **backdoor criterion** with respect to the ordered pair (A, Y) in a DAG if:

1. No variable in $\underset{\sim}{L}$ is a descendant of A.
2. $\underset{\sim}{L}$ blocks every path between A and Y that has an arrow into A (a "backdoor" path).

Remark 6.5 (Identification by backdoor adjustment). If $\underset{\sim}{L}$ satisfies the backdoor criterion, then adjusting for $\underset{\sim}{L}$ identifies the causal effect of A on Y:

$$\mathrm{E}[Y(a)] = \sum_{\underset{\sim}{l}} \mathrm{E}[Y \mid A = a, \underset{\sim}{L} = \underset{\sim}{l}] \, \mathrm{P}(\underset{\sim}{L} = \underset{\sim}{l})$$

6.5 HERS study DAG

The "heart and estrogen/progestin study" (HERS) was a clinical trial of hormone therapy for prevention of recurrent heart attacks and death among 2,763 post-menopausal women with existing coronary heart disease (CHD) [5].

The trial was conducted at 20 US clinical centers. Participants were randomized to receive either conjugated equine estrogens (0.625 mg/day) plus medroxyprogesterone acetate (2.5 mg/day) or a matching placebo [5]. Women were followed for a sample mean of 4.1 years [5].

The primary outcome was nonfatal myocardial infarction or CHD death [5].

We now examine the assumed causal structure among key baseline variables, treatment, and recurrent CHD events.

Figure 1 shows a DAG for the HERS study, representing the assumed causal structure among key variables.

The main goal of the HERS study was to estimate the causal effect of hormone therapy (HT) on recurrent CHD event risk[15].

Data dictionary for key HERS variables used in the DAG and examples.

Table 1

Variable	Meaning	Type
`HT`	Hormone-therapy assignment (placebo vs estrogen+progestin)	Binary[16] categorical[17]
`CHD`	Recurrent coronary heart disease event outcome	Binary categorical
`age`	Baseline age (years)	Continuous[18]
`smoking`	Baseline smoking status	Binary categorical
`diabetes`	Baseline diabetes status	Binary categorical
`BMI`	Body mass index	Continuous
`LDL`	Low-density lipoprotein cholesterol	Continuous
`HDL`	High-density lipoprotein cholesterol	Continuous

[15] binary-outcome-associations.qmd#def-risk-epi
[16] https://morrison-lab.github.io/sds/data.html#def-binary-var
[17] https://morrison-lab.github.io/sds/data.html#def-categorical-var
[18] https://morrison-lab.github.io/sds/data.html#def-continuous-var

Table 1

Variable	Meaning	Type
`statins`	Baseline statin use	Binary categorical
`SBP`	Systolic blood pressure	Continuous

```
library(dagitty)

hers_dag <- dagitty(
  'dag {
    HT [exposure, pos="0,0"]
    CHD [outcome, pos="4,0"]
    age [pos="2,2"]
    smoking [pos="-1,-2"]
    diabetes [pos="1,-1.5"]
    BMI [pos="-1,1"]
    LDL [pos="2,1"]
    HDL [pos="2,-1"]
    statins [pos="3,2"]
    SBP [pos="3,-2"]

    HT -> CHD
    HT -> HDL
    HT -> LDL
    age -> CHD
    smoking -> CHD
    smoking -> HDL
    BMI -> CHD
    BMI -> diabetes
    BMI -> SBP
    BMI -> LDL
    diabetes -> CHD
    diabetes -> LDL
    LDL -> CHD
    HDL -> CHD
    statins -> LDL
    SBP -> CHD
  }'
)

plot(hers_dag)
```




Figure 1: Directed Acyclic Graph (DAG) for the HERS study. `HT` = hormone therapy (exposure);

```
adjustmentSets(
  hers_dag,
  exposure = "HT",
  outcome = "CHD",
  type = "minimal"
)
#>  {}
```

In Figure 1:

- **age** is prognostic for CHD: older women are at higher risk of CHD events. Since HT was randomized in HERS, age is not a confounder of the HT → CHD relationship, but adjusting for age can improve precision.
- **LDL** and **HDL** are *mediators*: hormone therapy affects lipid levels, which in turn affect CHD risk.
- **BMI** is a common cause of diabetes, SBP, and LDL.
- **statins** (statin use) affects LDL independently of HT (statins are medications prescribed to lower LDL).
- This DAG is simplified; a more complete model might include additional variables such as anti-hypertensive medication use.

6.6 What variables should we adjust for?

Using DAG theory, we can identify the minimum adjustment set needed to estimate the total causal effect of HT on CHD.

Since HT was randomized in HERS, there are no backdoor paths from HT to CHD. The `adjustmentSets()` function confirms that no adjustment is needed to estimate the total causal effect of HT on CHD.

The variables `LDL` and `HDL` are mediators on the path HT → LDL/HDL → CHD, so they should **not** be included when estimating the *total* causal effect of HT on CHD.

Including `LDL` and `HDL` in the model would block a part of the causal effect of HT, giving us only the *direct* effect of HT on CHD (not through lipids).

6.7 Analyzing LDL in the HERS study

The HERS baseline dataset includes LDL and HDL cholesterol, allowing us to examine the effect of hormone therapy on lipid levels.

Here we build models to predict LDL cholesterol, illustrating how DAG thinking guides predictor selection[19].

```
library(dplyr)

hers <- rmb::hers |> haven::zap_labels()

hers_ldl <-
  hers |>
  filter(!is.na(LDL)) |>
  mutate(
    HT = factor(HT, levels = c(0, 1), labels = c("Placebo", "HT")),
    statins = factor(statins, levels = c(0, 1), labels = c("No", "Yes")),
    diabetes = factor(diabetes, levels = c(0, 1), labels = c("No", "Yes")),
    smoking = factor(smoking, levels = c(0, 1), labels = c("No", "Yes"))
  )
```

[19] predictor-selection.qmd#sec-pred-sel-strategies

Table 3: Summary table for LDL analysis variables in HERS

```
library(gtsummary)

hers_ldl |>
  dplyr::select(LDL, HT, age, BMI, diabetes, smoking, statins) |>
  tbl_summary(by = HT) |>
  add_overall() |>
  bold_labels()
```

Characteristic	**Overall** N = 2,752[1]	**Placebo** N = 1,379[1]	**HT** N = 1,373[1]
LDL cholesterol (mg/dl)	141 (120, 166)	141 (119, 165)	141 (120, 167)
age in years	67 (62, 72)	68 (62, 72)	67 (62, 72)
BMI (kg/m^2)	27.7 (24.6, 31.7)	27.6 (24.5, 31.7)	27.9 (24.7, 31.8)
Unknown	5	4	1
diabetes	727 (26%)	350 (25%)	377 (27%)
smoking	355 (13%)	180 (13%)	175 (13%)
statins	998 (36%)	516 (37%)	482 (35%)

[1]Median (Q1, Q3); n (%)

Table 4: Unadjusted linear regression for LDL in HERS

```
model_ldl_unadj <-
  lm(LDL ~ HT, data = hers_ldl)

model_ldl_unadj |>
  tbl_regression() |>
  bold_labels()
```

Characteristic	**Beta**	**95% CI**	**p-value**
HT			
Placebo	—	—	
HT	0.36	-2.5, 3.2	0.8

Abbreviation: CI = Confidence Interval

6.7.1 Unadjusted model

6.7.2 Adjusted model

Based on the DAG in Figure 1, BMI is a common cause of LDL and other outcomes (but age does not confound the HT → LDL relationship, since HT was randomized).

In a randomized trial, treatment assignment is independent[20] of pre-treatment covariates[21], so confounding adjustment is not required to estimate the causal effect. However, adjusting for strong predictors[22] of the outcome[23] can improve precision[24].

Note that `statins` is a strong predictor of LDL (statin medications are prescribed specifically to lower LDL). Including `statins` substantially reduces residual variability, improving precision of the HT coefficient.

[20]https://morrison-lab.github.io/pds/independence.html#def-indpt
[21]Intro-to-GLMs.qmd#def-predictors
[22]Intro-to-GLMs.qmd#def-predictors
[23]Intro-to-GLMs.qmd#def-outcome
[24]https://morrison-lab.github.io/pds/variance-covariance.html#def-precision

Table 5: Adjusted linear regression for LDL in HERS

```
model_ldl_adj <-
  lm(LDL ~ HT + age + BMI + diabetes + statins, data = hers_ldl)

model_ldl_adj |>
  tbl_regression() |>
  bold_labels()
```

Characteristic	Beta	95% CI	p-value
HT			
Placebo	—	—	
HT	-0.14	-2.9, 2.6	>0.9
age in years	-0.23	-0.44, -0.02	0.035
BMI (kg/m^2)	0.46	0.19, 0.72	<0.001
diabetes			
No	—	—	
Yes	-4.6	-7.9, -1.4	0.005
statins			
No	—	—	
Yes	-17	-20, -14	<0.001

Abbreviation: CI = Confidence Interval

6.8 An observational study example: WCGS

The HERS study is a randomized trial, so confounding of the treatment-outcome relationship is not a concern. To illustrate DAG-guided confounder selection in an **observational study**, we use the Western Collaborative Group Study (WCGS).

The WCGS was a prospective cohort study of 3,154 men employed by California companies, followed from 1960 to 1969 [8]. The exposure of primary interest is **Type A behavior pattern** (a personality trait characterized by competitive drive, time urgency, and hostility), hypothesized to increase CHD risk[25] independently of classic cardiovascular risk factors[26].

Data dictionary for key WCGS variables used in the DAG and examples.

Table 6

Variable	Meaning	Type
`personality_type`	Personality type (Type A vs Type B behavior pattern)	Binary categorical
`CHD`	Coronary heart disease event outcome	Binary categorical
`age`	Baseline age (years)	Continuous
`smoking`	Baseline smoking status	Binary categorical
`chol`	Serum cholesterol	Continuous
`sbp`	Systolic blood pressure	Continuous
`bmi`	Body mass index	Continuous

[25] binary-outcome-associations.qmd#def-risk-epi
[26] Intro-to-GLMs.qmd#def-predictors

```
wcgs_dag <- dagitty(
  'dag {
    personality_type [exposure, pos="0,0"]
    CHD [outcome, pos="4,0"]
    age [pos="1.4,2"]
    bmi [pos="-0.3,1.1"]
    smoking [pos="1.1,-1.8"]
    chol [pos="2.7,1"]
    sbp [pos="2.8,-1.2"]

    personality_type -> CHD
    age -> CHD
    age -> chol
    age -> sbp
    age -> bmi
    age -> personality_type
    personality_type -> smoking
    smoking -> CHD
    chol -> CHD
    sbp -> CHD
    bmi -> CHD
    bmi -> chol
    bmi -> sbp
  }'
)

plot(wcgs_dag)
```




Figure 2: Directed Acyclic Graph (DAG) for the WCGS study. `personality_type` = personality type (exposure; Type A vs Type B); `CHD` = coronary heart disease events (outcome); `chol` = serum cholesterol; `sbp` = systolic blood pressure.

In Figure 2, `age` is a **confounder**: it has direct causal arrows to both the exposure (`personality_type`) and the outcome (`CHD`). We assume `personality_type -> smoking` rather than `smoking -> personality_type` as an explicit modeling assumption for this teaching example: we treat Type A behavior pattern as a relatively stable trait that can influence smoking behavior. Under that assumption, `smoking` is a mediator on the path `personality_type -> smoking -> CHD`, not a confounder. BMI is causally affected by age. Cholesterol and SBP are causally affected by both age and BMI. All three have direct effects on CHD. Unlike in HERS, where HT was randomized to estimate its causal effect on CHD event risk, we cannot assume the exposure is unconfounded here.

```
wcgs_adj <- adjustmentSets(
  wcgs_dag,
  exposure = "personality_type",
  outcome = "CHD",
  type = "minimal"
)
wcgs_adj
#> { age }
```

The `adjustmentSets()` function returns the minimal set(s) of variables that block all backdoor paths from `personality_type` to `CHD`. Adjusting for any of these minimal sets is sufficient to estimate the total causal effect of Type A behavior on CHD (under the assumptions encoded in the DAG).

7 Regression Adjustment

Video lecture

Richard McElreath's *Good and Bad Controls*[a] (Statistical Rethinking 2026, Lecture A07) also accompanies Chapter 6 of McElreath [6], and discusses which covariates[b] a regression model[c] should and should not adjust for [7].

[a] https://www.youtube.com/watch?v=sTieMzOcreQ
[b] Intro-to-GLMs.qmd#def-predictors
[c] Intro-to-GLMs.qmd#def-regression-model

Under the causal assumptions (consistency, conditional exchangeability, and positivity), the causal effect can be estimated by adjusting for the confounders $\underset{\sim}{L}$ in a regression model[27]. The identification result behind that adjustment is standardization:

Theorem 7.1 (Identification by standardization)**.** *Suppose $\underset{\sim}{L}$ is discrete, and that consistency, conditional exchangeability given $\underset{\sim}{L}$, and positivity hold. Then for each treatment level $a \in \{0, 1\}$,*

$$\mathrm{E}[Y(a)] = \sum_{\underset{\sim}{l}} \mathrm{E}[Y \mid A = a, \underset{\sim}{L} = \underset{\sim}{l}] \, \mathrm{P}(\underset{\sim}{L} = \underset{\sim}{l}) \tag{2}$$

where the sum runs over the support of $\underset{\sim}{L}$.

Proof

Proof. For treatment level a:

[27] Intro-to-GLMs.qmd#def-regression-model

$$
\begin{aligned}
\mathrm{E}[Y(a)] &= \sum_{\underset{\sim}{l}} \mathrm{E}[Y(a) \mid \underset{\sim}{L} = \underset{\sim}{l}]\,\mathrm{P}(\underset{\sim}{L} = \underset{\sim}{l}) && \text{(law of total expectation over } \underset{\sim}{L}) \\
&= \sum_{\underset{\sim}{l}} \mathrm{E}[Y(a) \mid A = a, \underset{\sim}{L} = \underset{\sim}{l}]\,\mathrm{P}(\underset{\sim}{L} = \underset{\sim}{l}) && \text{(conditional exchangeability: } Y(a) \perp\!\!\!\perp A \mid \underset{\sim}{L}) \\
&= \sum_{\underset{\sim}{l}} \mathrm{E}[Y \mid A = a, \underset{\sim}{L} = \underset{\sim}{l}]\,\mathrm{P}(\underset{\sim}{L} = \underset{\sim}{l}) && \text{(consistency: } Y = Y(a) \text{ when } A = a)
\end{aligned}
$$

Positivity guarantees $\mathrm{P}(A = a \mid \underset{\sim}{L} = \underset{\sim}{l}) > 0$ for every $\underset{\sim}{l}$ in the support, so each conditioning event $\{A = a, \underset{\sim}{L} = \underset{\sim}{l}\}$ has positive probability and every term in the second and third lines is defined. □

Exm

Example 7.1 (Standardizing over disease severity)**.** In the population of Example 4.1, $\mathrm{P}(L = 0) = 600/1000 = 0.6$ and $\mathrm{P}(L = 1) = 0.4$, and by consistency the observed stratum means are $\mathrm{E}[Y \mid A = 1, L = l] = \mathrm{E}[Y(1) \mid L = l]$ and $\mathrm{E}[Y \mid A = 0, L = l] = \mathrm{E}[Y(0) \mid L = l]$. By Theorem 7.1:

$$
\begin{aligned}
\mathrm{E}[Y(1)] &= 10 \cdot 0.6 + 20 \cdot 0.4 && \text{(standardization formula with } a = 1) \\
&= 6 + 8 && \text{(multiply)} \\
&= 14 && \text{(add)} \\
\mathrm{E}[Y(0)] &= 8 \cdot 0.6 + 18 \cdot 0.4 && \text{(standardization formula with } a = 0) \\
&= 4.8 + 7.2 && \text{(multiply)} \\
&= 12 && \text{(add)} \\
\mathrm{ATE} &= 14 - 12 && \text{(definition of ATE)} \\
&= 2 && \text{(subtract)}
\end{aligned}
$$

Standardization recovers the true ATE of 2, whereas the naive contrast in Example 5.2 gave 7. Standardization weights both treatment groups' stratum means by the same distribution of L, the whole population's, instead of each group's own distribution of L.

7.1 Direct standardization (G-computation)

The **G-computation** estimator (also called **direct standardization** or **standardization**) estimates the ATE by:

1. Fitting a regression model[28]: $\hat{\mu}(a, \underset{\sim}{l}) \overset{\text{def}}{=} \mathrm{E}[Y \mid \widehat{A = a}, \underset{\sim}{L} = \underset{\sim}{l}]$
2. Predicting the potential outcome mean for each individual under both treatment levels $a = 1$ and $a = 0$:
$$\hat{Y}_i(a) \overset{\text{def}}{=} \hat{\mu}(a, \underset{\sim}{L}_i)$$
3. Averaging over the study population[29]:
$$\widehat{\mathrm{E}[Y(a)]} \overset{\text{def}}{=} \frac{1}{n} \sum_{i=1}^{n} \hat{Y}_i(a)$$
4. Estimating the ATE as the contrast:
$$\widehat{\mathrm{ATE}} \overset{\text{def}}{=} \widehat{\mathrm{E}[Y(1)]} - \widehat{\mathrm{E}[Y(0)]}$$

G-computation is model-based: it relies on the regression model being correctly specified. A misspecified model will produce biased[30] causal estimates even if confounders are included.

[28] Intro-to-GLMs.qmd#def-regression-model
[29] https://morrison-lab.github.io/sds/exploratory-descriptive.html#def-population
[30] https://morrison-lab.github.io/sds/estimation.html#def-bias

G-computation is easily generalized to:

- Binary outcomes (using logistic regression[31])
- Survival outcomes (using Cox[32] or Weibull[33] models)
- Multiple treatments or treatment levels

7.2 Regression adjustment in linear models

When the outcome[34] is continuous[35] and the regression model is linear, G-computation simplifies to the **adjusted treatment coefficient** from a linear regression[36]:

$$Y = \beta_0 + \beta_A A + \underset{\sim}{\beta_L}^\top \underset{\sim}{L} + \varepsilon$$

Under consistency, conditional exchangeability, and positivity, the coefficient β_A equals the average causal effect of A on Y:

$$\begin{aligned}
\text{ATE} &\overset{\text{def}}{=} \text{E}[Y(1) - Y(0)] && \text{(definition of ATE)} \\
&= \text{E}[Y(1)] - \text{E}[Y(0)] && \text{(linearity of expectation)} \\
&= \text{E}[\text{E}[Y(1) \mid \underset{\sim}{L}]] - \text{E}[\text{E}[Y(0) \mid \underset{\sim}{L}]] && \text{(law of iterated expectations)} \\
&= \text{E}[\text{E}[Y \mid A = 1, \underset{\sim}{L}]] - \text{E}[\text{E}[Y \mid A = 0, \underset{\sim}{L}]] && \text{(consistency and conditional exchangeability)} \\
&= \text{E}\left[\beta_0 + \beta_A(1) + \underset{\sim}{\beta_L}^\top \underset{\sim}{L}\right] - \text{E}\left[\beta_0 + \beta_A(0) + \underset{\sim}{\beta_L}^\top \underset{\sim}{L}\right] && \text{(linear outcome model specification)} \\
&= \left(\beta_0 + \beta_A + \underset{\sim}{\beta_L}^\top \text{E}[\underset{\sim}{L}]\right) - \left(\beta_0 + \underset{\sim}{\beta_L}^\top \text{E}[\underset{\sim}{L}]\right) && \text{(linearity of expectation)} \\
&= \beta_A && \text{(algebraic simplification)}
\end{aligned}$$

Thus, when the outcome model is correctly specified and contains no treatment-covariate interactions[37], β_A directly identifies the ATE.

The regression adjustment estimator for the ATE is equivalent to G-computation when the outcome model is linear and has no treatment-covariate interactions. When there are interactions, G-computation and regression adjustment can give different estimates. G-computation is preferred in that case.

7.3 Example: regression adjustment for LDL in HERS

The HERS study DAG (see Section 6.5) shows that `LDL` and `HDL` are mediators of the HT → CHD pathway. For estimating the total causal effect of HT on CHD, we should not adjust for `LDL` or `HDL`.

However, to estimate the causal effect of **HT on LDL** itself, we use a separate regression model with `LDL` as the outcome. In this model, `LDL` is no longer a mediator but the outcome of interest.

Since HT was randomized in HERS, no adjustment is needed for causal identification of HT → LDL. However, adjusting for strong predictors[38] of LDL improves precision[39]:

```
library(dplyr)

hers <- rmb::hers |> haven::zap_labels()

hers_ldl <-
```

[31] logistic-regression.qmd#def-logistic-regression
[32] proportional-hazards-models.qmd#def-ph-model
[33] https://morrison-lab.github.io/pds/distributions.html#def-weibull
[34] Intro-to-GLMs.qmd#def-outcome
[35] https://morrison-lab.github.io/sds/data.html#def-continuous-var
[36] Linear-models-overview.qmd#def-lm-general-form
[37] Intro-to-GLMs.qmd#def-interaction
[38] Intro-to-GLMs.qmd#def-predictors
[39] https://morrison-lab.github.io/pds/variance-covariance.html#def-precision

Table 7: Adjusted regression for the causal effect of HT on LDL in HERS. Since HT was randomized, adjustment is for precision only.

```
library(gtsummary)

model_ldl_adj <-
  lm(LDL ~ HT + age + BMI + diabetes + statins, data = hers_ldl)

model_ldl_adj |>
  tbl_regression() |>
  bold_labels()
```

Characteristic	Beta	95% CI	p-value
HT			
Placebo	—	—	
HT	-0.14	-2.9, 2.6	>0.9
age in years	-0.23	-0.44, -0.02	0.035
BMI (kg/m^2)	0.46	0.19, 0.72	<0.001
diabetes			
No	—	—	
Yes	-4.6	-7.9, -1.4	0.005
statins			
No	—	—	
Yes	-17	-20, -14	<0.001

Abbreviation: CI = Confidence Interval

```
  hers |>
  filter(!is.na(LDL)) |>
  mutate(
    HT = factor(HT, levels = c(0, 1), labels = c("Placebo", "HT")),
    statins = factor(statins, levels = c(0, 1), labels = c("No", "Yes")),
    diabetes = factor(diabetes, levels = c(0, 1), labels = c("No", "Yes"))
  )
```

The estimated causal effect of HT on LDL is approximately -11 mg/dL (negative = lower LDL on HT), with narrower confidence intervals[40] than the unadjusted estimate.

8 Propensity Score Methods

Propensity score methods are an alternative approach to confounding adjustment in observational studies. They replace adjustment for a high-dimensional confounder set $\underset{\sim}{L}$ with adjustment for a single scalar summary: the propensity score.

Definition 8.1 (Propensity score)**.** The **propensity score** is the conditional probability[a] of treatment given the observed covariates[b]:

$$e(\underset{\sim}{L}) \stackrel{\text{def}}{=} \mathrm{P}(A = 1 \mid \underset{\sim}{L})$$

[a] https://morrison-lab.github.io/pds/probability-basics.html#def-conditional-prob
[b] Intro-to-GLMs.qmd#def-predictors

[40] https://morrison-lab.github.io/sds/inference.html#def-confidence-interval

Exm

Example 8.1 (Propensity scores in the severity example)**.** In the population of Example 4.1, the propensity score (Definition 8.1) takes two values: $e(0) = \mathrm{P}(A = 1 \mid L = 0) = 120/600 = 0.2$ and $e(1) = \mathrm{P}(A = 1 \mid L = 1) = 280/400 = 0.7$. A severe patient is $0.7/0.2 = 3.5$ times as likely to be treated as a non-severe patient.

Remark 8.1 (Propensity scores and exchangeability). Rosenbaum and Rubin [9] showed that if $A \perp\!\!\!\perp (Y(0), Y(1)) \mid \underset{\sim}{L}$ (conditional exchangeability given $\underset{\sim}{L}$), then also:

$$A \perp\!\!\!\perp (Y(0), Y(1)) \mid e(\underset{\sim}{L})$$

Definition 8.2 (Balancing score)**.** In other words, the propensity score (Definition 8.1) is a **balancing score**: conditioning on the propensity score is sufficient to remove confounding.

8.1 Propensity score estimation

The propensity score is estimated by fitting a model for the probability of treatment given covariates[41], typically using logistic regression[42]:

$$\mathrm{logit}(\hat{e}(\underset{\sim}{L}_i)) = \hat{\beta}_0 + \hat{\beta}_1 L_{i1} + \cdots + \hat{\beta}_p L_{ip}$$

The estimated propensity score $\hat{e}(\underset{\sim}{L}_i)$ is the predicted probability of treatment for each individual.

8.2 Propensity score methods

There are four main ways to use the propensity score:

1. **Matching**: Match each treated individual to one or more untreated individuals with similar propensity scores. The matched dataset is then analyzed as a simple comparison.
2. **Stratification (subclassification)**: Divide individuals into strata based on propensity score quantiles[43]. Estimate the causal effect within each stratum, then combine across strata.
3. **Inverse probability weighting (IPW)**: Weight each individual by the inverse of their probability of receiving the treatment they actually received. See Definition 8.3.
4. **Covariate adjustment using the propensity score**: Include the estimated propensity score as a covariate[44] in a regression model[45] for the outcome[46].

Definition 8.3 (Inverse probability weighting (IPW))**.** The **inverse probability weighted (IPW)** estimator of the ATE uses weights:

$$w_i \overset{\text{def}}{=} \frac{A_i}{\hat{e}(\underset{\sim}{L}_i)} + \frac{1 - A_i}{1 - \hat{e}(\underset{\sim}{L}_i)}$$

The IPW estimator of $\mathrm{E}[Y(a)]$ is:

$$\widehat{\mathrm{E}[Y(a)]}_{\mathrm{IPW}} \overset{\text{def}}{=} \frac{\sum_{i:A_i=a} w_i Y_i}{\sum_{i:A_i=a} w_i}$$

[41] Intro-to-GLMs.qmd#def-predictors
[42] logistic-regression.qmd#def-logistic-regression
[43] https://morrison-lab.github.io/sds/nonparametric-models.html#def-sample-quantile
[44] Intro-to-GLMs.qmd#def-predictors
[45] Intro-to-GLMs.qmd#def-regression-model
[46] Intro-to-GLMs.qmd#def-outcome

Definition 8.4 (Pseudo-population)**.** Reweighting by the weights w_i of Definition 8.3 creates a **pseudo-population** in which treatment is independent[a] of covariates, effectively removing confounding.

[a] https://morrison-lab.github.io/pds/independence.html#def-indpt

Theorem 8.1 (Identification by inverse probability weighting)**.** *For $a \in \{0,1\}$, write $\pi_a(\underset{\sim}{l}) \overset{def}{=} \mathrm{P}(A = a \mid \underset{\sim}{L} = \underset{\sim}{l})$, so that $\pi_1 = e$ and $\pi_0 = 1 - e$. If consistency, conditional exchangeability given $\underset{\sim}{L}$, and positivity hold, then*

$$\mathrm{E}[Y(a)] = \mathrm{E}\left[\frac{\mathbb{1}(A = a)\, Y}{\pi_a(\underset{\sim}{L})}\right] \tag{3}$$

Proof

Proof.

$$\begin{aligned}
\mathrm{E}\left[\frac{\mathbb{1}(A = a)\, Y}{\pi_a(\underset{\sim}{L})}\right] &= \mathrm{E}\left[\mathrm{E}\left[\frac{\mathbb{1}(A = a)\, Y}{\pi_a(\underset{\sim}{L})} \mid \underset{\sim}{L}\right]\right] && \text{(law of total expectation)} \\
&= \mathrm{E}\left[\frac{1}{\pi_a(\underset{\sim}{L})}\, \mathrm{E}[\mathbb{1}(A = a)\, Y \mid \underset{\sim}{L}]\right] && \text{(pull out } 1/\pi_a(\underset{\sim}{L}), \text{ which positivity keeps finite)} \\
&= \mathrm{E}\left[\frac{1}{\pi_a(\underset{\sim}{L})}\, \mathrm{E}[\mathbb{1}(A = a)\, Y(a) \mid \underset{\sim}{L}]\right] && \text{(consistency: } Y = Y(a) \text{ when } A = a) \\
&= \mathrm{E}\left[\frac{1}{\pi_a(\underset{\sim}{L})}\, \mathrm{E}[\mathbb{1}(A = a) \mid \underset{\sim}{L}]\, \mathrm{E}[Y(a) \mid \underset{\sim}{L}]\right] && \text{(conditional exchangeability: } A \perp\!\!\!\perp Y(a) \mid \underset{\sim}{L}) \\
&= \mathrm{E}\left[\frac{1}{\pi_a(\underset{\sim}{L})}\, \pi_a(\underset{\sim}{L})\, \mathrm{E}[Y(a) \mid \underset{\sim}{L}]\right] && (\,\mathrm{E}[\mathbb{1}(A = a) \mid \underset{\sim}{L}] = \mathrm{P}(A = a \mid \underset{\sim}{L})) \\
&= \mathrm{E}[\mathrm{E}[Y(a) \mid \underset{\sim}{L}]] && \text{(cancel } \pi_a(\underset{\sim}{L})) \\
&= \mathrm{E}[Y(a)] && \text{(law of total expectation)}
\end{aligned}$$

□

Exm

Example 8.2 (The pseudo-population in the severity example)**.** In the population of Example 4.1, with the propensity scores of Example 8.1, the weights of Definition 8.3 are:

Group	patients	weight w_i	weighted count
treated, $L = 0$	120	$1/0.2 = 5$	600
treated, $L = 1$	280	$1/0.7 \approx 1.43$	400
untreated, $L = 0$	480	$1/0.8 = 1.25$	600
untreated, $L = 1$	120	$1/0.3 \approx 3.33$	400

In the pseudo-population, each treatment group has 1000 members, 600 non-severe and 400 severe: the same severity mix as the whole population, so L no longer predicts treatment. The weighted means are:

$$\begin{aligned}
\widehat{\mathrm{E}[Y(1)]}_{\mathrm{IPW}} &= \frac{600 \cdot 10 + 400 \cdot 20}{600 + 400} && \text{(IPW estimator, treated)} \\
&= \frac{6000 + 8000}{1000} && \text{(multiply)} \\
&= 14 && \text{(add and divide)} \\
\widehat{\mathrm{E}[Y(0)]}_{\mathrm{IPW}} &= \frac{600 \cdot 8 + 400 \cdot 18}{600 + 400} && \text{(IPW estimator, untreated)} \\
&= \frac{4800 + 7200}{1000} && \text{(multiply)} \\
&= 12 && \text{(add and divide)}
\end{aligned}$$

These weighted means match Example 7.1, as Theorem 8.1 says they should, and their difference recovers the ATE of 2.

8.3 Propensity score example: WCGS

In the WCGS study, we can use propensity score methods to estimate the causal effect of Type A behavior on CHD. We first fit a logistic regression[47] model predicting Type A behavior from the confounders identified by the DAG (Section 6.8):

```
library(dplyr)

here::here() |>
  fs::path("Data/wcgs.rda") |>
  load()

wcgs_ps <-
  wcgs |>
  mutate(
    dibpat_A = as.integer(dibpat == "Type A"),
    chd69_yn = as.integer(chd69 == "Yes")
  ) |>
  filter(
    !is.na(age), !is.na(dibpat), !is.na(chd69),
    !is.na(bmi), !is.na(sbp), !is.na(chol), !is.na(smoke)
  )
```

```
ps_model <- glm(
  dibpat_A ~ age + bmi + sbp + chol + smoke,
  data = wcgs_ps,
  family = binomial()
)

wcgs_ps <- wcgs_ps |>
  mutate(
    ps = predict(ps_model, type = "response"),
    ipw = dibpat_A / ps + (1 - dibpat_A) / (1 - ps)
  )
```

[47] logistic-regression.qmd#def-logistic-regression

```r
library(ggplot2)

ggplot(wcgs_ps, aes(x = ps, fill = factor(dibpat_A))) +
  geom_histogram(alpha = 0.6, bins = 30, position = "identity") +
  scale_fill_manual(
    values = c("steelblue", "tomato"),
    labels = c("Type B", "Type A"),
    name = "Personality type"
  ) +
  labs(
    x = "Estimated propensity score",
    y = "Count",
    title = "Propensity score distributions by personality type (WCGS)"
  ) +
  theme_bw()
```




Figure 3: Distribution of estimated propensity scores by personality type in the WCGS study. Good overlap between the two groups (the **overlap** or **common support** assumption) supports positivity.

The IPW-weighted odds ratio for Type A vs. Type B behavior estimates the causal odds ratio for CHD, adjusting for confounding by age, BMI, SBP, cholesterol, and smoking.

Compare this to the unadjusted estimate: the difference reflects the degree of confounding in the WCGS data.

9 Summary

Causal inference requires moving beyond statistical association to make statements about what would happen under intervention. The key ideas from this chapter are:

Potential outcomes define what we mean by a causal effect. For each unit, there is a potential outcome under treatment and a potential outcome under control; the individual causal effect is their difference.

Causal estimands — ATE, ATT, ATU — summarize the causal effect at the population level.

Three key assumptions are needed to identify causal effects from observed data:

Table 8: Inverse probability weighted (IPW) logistic regression model for the effect of Type A personality on CHD in WCGS.

```
library(gtsummary)

outcome_ipw <- glm(
  chd69_yn ~ dibpat_A,
  data = wcgs_ps,
  weights = ipw,
  family = quasibinomial()
)

outcome_ipw |>
  tbl_regression(exponentiate = TRUE) |>
  bold_labels()
```

Characteristic	OR	95% CI	p-value
dibpat_A	1.93	1.48, 2.53	<0.001

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

1. **Consistency**: The observed outcome equals the potential outcome under the received treatment.
2. **Exchangeability**: Treatment assignment is independent[48] of potential outcomes, given measured confounders.
3. **Positivity**: Every individual has a positive probability of receiving each treatment level.

Randomized experiments achieve unconditional exchangeability and allow unbiased causal inference from simple treatment comparisons.

In observational studies, confounding violates exchangeability. DAGs help identify which variables are confounders, mediators, and colliders, and the backdoor criterion identifies sufficient adjustment sets.

Estimation methods — regression adjustment (G-computation) and propensity score methods (matching, stratification, IPW) — can recover causal estimates under these three core identification assumptions.

Learning Objectives

After completing this chapter, students should be able to:

1. **Define potential outcomes** and the fundamental problem of causal inference.
2. **Distinguish** the ATE, ATT, and ATU, and explain when they agree and when they differ.
3. **State and interpret** the three core causal assumptions: consistency, exchangeability, and positivity.
4. **Explain why randomization** achieves unconditional exchangeability and allows unbiased causal effect estimation.
5. **Use a DAG** to identify confounders, mediators, and colliders, and apply the backdoor criterion to select sufficient adjustment sets.
6. **Describe the G-computation estimator** and explain how it relates to regression adjustment.
7. **Describe propensity score methods** (matching, stratification, IPW) and explain how they use the propensity score to remove confounding.

[48] https://morrison-lab.github.io/pds/independence.html#def-cind

8. **Contrast causal inference from randomized trials with causal inference from observational studies**, and articulate the additional assumptions required in each setting.

References

[1] E. Vittinghoff, D. V. Glidden, S. C. Shiboski, and C. E. McCulloch, *Regression methods in biostatistics: Linear, logistic, survival, and repeated measures models*, 2nd ed. Springer, 2012. Available: https://doi.org/10.1007/978-1-4614-1353-0

[2] M. A. Hernán and J. M. Robins, *Causal inference: What if*. Chapman & Hall/CRC, 2020. Available: https://www.hsph.harvard.edu/miguel-hernan/causal-inference-book/

[3] P. W. Holland, "Statistics and causal inference," *Journal of the American Statistical Association*, vol. 81, no. 396, pp. 945–960, 1986, doi: 10.2307/2289064[49].

[4] D. B. Rubin, "Estimating causal effects of treatments in randomized and nonrandomized studies," *Journal of Educational Psychology*, vol. 66, no. 5, pp. 688–701, 1974, doi: 10.1037/h0037350[50].

[5] S. Hulley *et al.*, "Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women," *JAMA : the journal of the American Medical Association*, vol. 280, no. 7, pp. 605–613, 1998.

[6] R. McElreath, *Statistical rethinking : A Bayesian course with examples in R and Stan*, Second edition. in Chapman & hall/CRC texts in statistical science series. Boca Raton, FL: CRC Press, 2020.

[7] R. McElreath, "Statistical rethinking 2026." Lecture videos, YouTube, 2026. Available: https://github.com/rmcelreath/stat_rethinking_2026

[8] R. H. Rosenman, R. J. Brand, C. D. Jenkins, M. Friedman, R. Straus, and M. Wurm, "Coronary heart disease in the Western Collaborative Group Study: Final follow-up experience of 8 1/2 years," *JAMA*, vol. 233, no. 8, pp. 872–877, 1975, doi: 10.1001/jama.1975.03260080034016[51].

[9] P. R. Rosenbaum and D. B. Rubin, "The central role of the propensity score in observational studies for causal effects," *Biometrika*, vol. 70, no. 1, pp. 41–55, 1983, doi: 10.1093/biomet/70.1.41[52].

[49] https://doi.org/10.2307/2289064
[50] https://doi.org/10.1037/h0037350
[51] https://doi.org/10.1001/jama.1975.03260080034016
[52] https://doi.org/10.1093/biomet/70.1.41