Inference for Logistic Regression

Last modified: 2026-10-16 10:06:09 (PDT)

Introduction

Logistic regression models a binary outcome, such as whether a patient recovers, from one or more predictors. Its fitted coefficients are maximum likelihood estimates, so the likelihood theory of Introduction to maximum likelihood gives them standard errors, confidence intervals and tests. This page specializes that theory to the logistic model and introduces the deviance, the measure of fit that the likelihood ratio test of a logistic model is built from.

Definition 0.1 (Logistic regression)**.** A **logistic regression** model for binary outcomes $y_1, \dots, y_n \in \{0, 1\}$ with predictor vectors $\underset{\sim}{x}_1, \dots, \underset{\sim}{x}_n$ says that the outcomes are independent and that

$$\Pr\left(Y_i = 1 \mid \underset{\sim}{x}_i\right) = \pi_i = \sigma\left(\underset{\sim}{\beta} \cdot \underset{\sim}{x}_i\right), \qquad \sigma(u) = \frac{1}{1 + e^{-u}}, \tag{1}$$

where $\underset{\sim}{x}_i = \left(1, x_{i1}, \dots, x_{ip}\right)^\top$ and $\underset{\sim}{\beta} = \left(\beta_0, \beta_1, \dots, \beta_p\right)^\top$. The leading 1 in $\underset{\sim}{x}_i$ makes β_0 the intercept, and β_j for $j \geq 1$ is the slope for the jth predictor.

Example 0.1 (A six-patient trial)**.** Six patients are split into a control group, coded $x = -1$, and a treatment group, coded $x = +1$. The outcome is $y = 1$ if the patient recovered. Fitting the model of Equation 1 by maximum likelihood gives the coefficients below.

```r
# group coded -1 (control) and +1 (treatment); y = 1 means recovered
wf_x <- rep(c(-1, 1), each = 3)
wf_y <- c(0, 0, 1, 0, 1, 1)
# maximum likelihood fit of the logistic model
wf_fit <- glm(wf_y ~ wf_x, family = binomial)
# the fitted intercept and slope
round(coef(wf_fit), 3)
#> (Intercept)        wf_x
#>       0.000       0.693
```

The fitted slope is $\hat{\beta}_1 = 0.693$. The control group recovered at rate $1/3$ and the treatment group at $2/3$, so the model fits both group proportions exactly.

Deviance

Definition 0.2 (Deviance)**.** The **deviance** of a fitted logistic regression with labels $y_i \in \{0, 1\}$ is minus twice its maximized log-likelihood:

$$D \stackrel{\text{def}}{=} -2\,\ell\left(\hat{\underset{\sim}{\beta}}\right) = -2\sum_{i=1}^{n}\{y_i \ln \hat{\pi}_i + (1 - y_i)\ln(1 - \hat{\pi}_i)\}, \tag{2}$$

where $\hat{\pi}_i = \sigma\left(\hat{\underset{\sim}{\beta}} \cdot \underset{\sim i}{x}\right)$ is the fitted probability.

Source

Nelder and Wedderburn (1972) introduced the deviance for the whole family of generalized linear models; logistic regression is the case of Bernoulli outcomes with the logit link. Hastie et al. (2009, sec. 4.4.2) defines the residual deviance of a fitted logistic regression as minus twice its log-likelihood.

Example 0.2 (The deviance of the six-patient trial)**.** For the fit of Example 0.1, R's `glm` reports the deviance. We check it against Equation 2 computed by hand.

```r
# fitted probabilities of recovery
wf_p <- fitted(wf_fit)
# minus twice the log-likelihood, from the formula
wf_dev_hand <- -2 * sum(wf_y * log(wf_p) + (1 - wf_y) * log(1 - wf_p))
# the deviance that glm reports
wf_dev_glm <- deviance(wf_fit)
c(by_hand = wf_dev_hand, glm = wf_dev_glm)
#> by_hand     glm
#> 7.63817 7.63817
```

Both give $D = 7.638$.

Remark (Why the deviance measures a shortfall)

Remark 0.1 (Why the deviance measures a shortfall). For labels $y_i \in \{0, 1\}$ the largest log-likelihood any model can reach is 0, by giving each label probability 1. So D is twice the amount by which the maximized log-likelihood falls short of that largest value.

Definition 0.3 (Null deviance)**.** The **null deviance** is the deviance of the model with an intercept and no other predictors:

$$D_0 = -2n\{\bar{y} \ln \bar{y} + (1 - \bar{y}) \ln(1 - \bar{y})\}, \tag{3}$$

where $\bar{y}$ is the fraction of labels equal to 1.

Example 0.3 (The null deviance of the six-patient trial)**.** Three of the six patients recovered, so $\bar{y} = 1/2$ and Equation 3 gives $D_0 = 2 \times 6 \ln 2$.

```r
# null deviance from the formula with ybar = 1/2
wf_null_hand <- 2 * 6 * log(2)
# the null deviance that glm reports
wf_null_glm <- wf_fit$null.deviance
c(by_hand = wf_null_hand, glm = wf_null_glm)
#> by_hand     glm
#> 8.31777 8.31777
```

Both give $D_0 = 8.318$.

Remark (Where the null deviance formula comes from)

Remark 0.2 (Where the null deviance formula comes from). The intercept-only model gives every observation the same fitted probability, and the maximizing value is the overall fraction $\bar{y}$ of labels equal to 1. Substituting $\bar{y}$ for each $\hat{\pi}_i$ in Equation 2 gives Equation 3.

Theorem 0.1 (The drop in deviance is the likelihood ratio statistic)**.** *Let D_0 be the null deviance and D the deviance of a logistic regression with q predictors besides the intercept. Then the likelihood ratio statistic of H_0: all q slopes are zero is*

$$\Lambda = D_0 - D.$$

Proof

Proof. In the notation of the likelihood ratio statistic, the unrestricted MLE $\hat{\theta}_{\mathrm{ML}}$ is the fitted coefficient vector $\hat{\beta}$, and $\hat{\theta}_0$ is the fit of the intercept-only model that H_0 imposes. Then $\ell\big(\hat{\theta}_{\mathrm{ML}}\big) = -D/2$ and $\ell\big(\hat{\theta}_0\big) = -D_0/2$. Substituting both into the definition gives $\Lambda = 2(-D/2 + D_0/2) = D_0 - D$. □

Example 0.4 (The likelihood ratio statistic of the treatment effect)**.** In the six-patient trial, $q = 1$.

```
# drop in deviance, the likelihood ratio statistic
wf_g2 <- wf_fit$null.deviance - deviance(wf_fit)
wf_g2
#> [1] 0.679596
```

The statistic is 0.68.

Corollary 0.1 (Distribution of the drop in deviance)**.** *Suppose the regularity conditions of Wilks' theorem hold for the logistic regression model. Then for a large sample the drop in deviance $D_0 - D$ has approximately a χ^2_q distribution when H_0 holds.*

Proof

Proof. By Theorem 0.1 the drop in deviance is the likelihood ratio statistic of q constraints, and Wilks' theorem gives its approximate distribution. □

Example 0.5 (Testing the treatment effect)**.** For the statistic of Example 0.4, the p-value comes from the chi-square distribution with $q = 1$ degree of freedom.

```
# upper tail of the chi-square distribution with 1 degree of freedom
wf_g2_p <- pchisq(wf_g2, df = 1, lower.tail = FALSE)
wf_g2_p
#> [1] 0.409726
```

The p-value is 0.41, so six patients give no evidence of a treatment effect.

Covariate patterns and the saturated model

Definition 0.4 (Covariate pattern)**.** A **covariate pattern** is a distinct value of the predictor vector among the observations $\underset{\sim}{x}_1, \ldots, \underset{\sim}{x}_n$. The K distinct values are written $\underset{\sim}{x}^{(1)}, \ldots, \underset{\sim}{x}^{(K)}$, so $K \leq n$.

Definition 0.5 (Replicates)**.** The **replicates** of the covariate pattern $\underset{\sim}{x}^{(k)}$ are the observations i with $\underset{\sim}{x}_i = \underset{\sim}{x}^{(k)}$. Their number is written m_k, so $\sum_{k=1}^{K} m_k = n$.

Remark 1: Source: covariate patterns and replicates

Dobson and Barnett (2018, sec. 5.5) introduces both terms: observations that share a combination of factor levels and the same values of any continuous predictors have the same covariate pattern, and are called replicates.

Definition 0.6 (Events at a covariate pattern)**.** The number of **events** at the covariate pattern $\underset{\sim}{x}^{(k)}$ is the sum of the binary outcomes over its replicates:

$$s_k = \sum_{i:\, \underset{\sim}{x}_i = \underset{\sim}{x}^{(k)}} y_i,$$

so $0 \leq s_k \leq m_k$.

Example 0.6 (Covariate patterns in the low-birth-weight data)**.** The `birthwt` data record whether each of 189 babies had low birth weight (`low = 1`, under 2.5 kg), along with the mother's smoking status and race, among other predictors (Hosmer and Lemeshow 1989).

Take `smoke` (0 or 1) and `race` (1, 2 or 3) as the predictors. Counting the distinct pairs, and the babies and low-weight births at each, gives one row per covariate pattern.

```
bw <- read.csv("data/birthwt.csv")
bw$race <- factor(bw$race)
# one row per covariate pattern: replicates and events
bw_grouped <- bw |>
  dplyr::summarize(
    nreps = dplyr::n(),
    nevents = sum(low),
    .by = c(smoke, race)
  ) |>
  dplyr::arrange(smoke, race)
bw_grouped
#>   smoke race nreps nevents
#> 1     0    1    44       4
#> 2     0    2    16       5
#> 3     0    3    55      20
#> 4     1    1    52      19
#> 5     1    2    10       6
#> 6     1    3    12       5
```

There are $K = 6$ covariate patterns among $n = 189$ babies. The pattern with the most replicates has `smoke` $= 0$ and `race` $= 3$, and it has 55 of them.

Definition 0.7 (Saturated model (full model, maximal model))**.** The **saturated model** (also called the **full model** (Dobson and Barnett 2018, sec. 5.5) or **maximal model** (Dobson and Barnett 2018, sec. 5.5)) for binary outcomes gives each pattern its own free event probability:

$$\Pr\left(Y_i = 1 \mid \underset{\sim i}{x} = \underset{\sim}{x}^{(k)}\right) = \pi_k, \qquad k = 1, \ldots, K, \tag{4}$$

with the outcomes independent given the predictors.

Remark 2: Source: saturated model

Dobson and Barnett (2018, sec. 5.5) defines the saturated model as the model with the maximum number of parameters that can be estimated, one per distinct covariate pattern, and says it is also called a maximal or full model. In model selection, "full model" often means something else: the model with every candidate predictor (Dobson and Barnett 2018, sec. 6.3). That model is generally smaller than the saturated model.

Theorem 0.2 (The saturated model fits each pattern's observed proportion)**.** *The maximum likelihood estimate of π_k in the saturated model is the observed proportion of events among the replicates of pattern k:*

$$\hat{\pi}_k = \frac{s_k}{m_k}.$$

Proof

Proof. Independence gives each pattern its own log-likelihood term, and the outcomes at pattern k contribute $s_k \ln \pi_k + (m_k - s_k)\ln(1-\pi_k)$. The terms share no parameters, so each is maximized on its own. First suppose $0 < s_k < m_k$, so that both logarithms are defined for $0 < \pi_k < 1$. Setting the derivative with respect to π_k to zero gives $s_k/\pi_k - (m_k - s_k)/(1-\pi_k) = 0$. Multiplying through by $\pi_k(1-\pi_k)$ gives $s_k(1-\pi_k) = (m_k - s_k)\pi_k$. Expanding gives $s_k - s_k\pi_k = m_k\pi_k - s_k\pi_k$. Adding $s_k\pi_k$ to both sides gives $s_k = m_k\pi_k$. Dividing both sides by m_k gives $\hat{\pi}_k = s_k/m_k$. The log-likelihood term increases to this point and decreases after it, so the point is a maximum. Now suppose $s_k = 0$. A count of 0 times a logarithm is taken as 0, so the term is $m_k \ln(1-\pi_k)$, which decreases in π_k and so peaks at $\pi_k = 0$. Next suppose $s_k = m_k$. By the same convention, the term is $m_k \ln \pi_k$, which peaks at $\pi_k = 1$. Both agree with s_k/m_k. □

Definition 0.8 (Deviance relative to the saturated model)**.** The **deviance relative to the saturated model** of a fitted model is twice the amount by which its maximized log-likelihood $\ell\left(\underset{\sim}{\hat{\beta}}\right)$ falls short of the maximized log-likelihood ℓ_{sat} of the saturated model:

$$D_{\text{sat}} = 2\left(\ell_{\text{sat}} - \ell\left(\underset{\sim}{\hat{\beta}}\right)\right). \tag{5}$$

Theorem 0.3 (The deviance relative to the saturated model, from grouped counts)**.** *For a fitted logistic regression with fitted event probabilities $\hat{\pi}_k$ at the K covariate patterns,*

$$D_{\text{sat}} = 2\sum_{k=1}^{K}\left\{s_k \ln\left(\frac{s_k}{m_k\hat{\pi}_k}\right) + (m_k - s_k)\ln\left(\frac{m_k - s_k}{m_k(1-\hat{\pi}_k)}\right)\right\}, \tag{6}$$

where a term with $s_k = 0$ or $s_k = m_k$ contributes 0 in its zero-count part.

Proof

Proof. Grouping the outcomes by pattern, the fitted model's log-likelihood is $\ell\left(\underset{\sim}{\hat{\beta}}\right) = \sum_{k=1}^{K}\{s_k \ln \hat{\pi}_k + (m_k - s_k)\ln(1-\hat{\pi}_k)\}$. By Theorem 0.2, the saturated log-likelihood is

the same sum with $\hat{\pi}_k$ replaced by s_k/m_k. Subtracting the fitted from the saturated log-likelihood, term by term, replaces each $\ln(s_k/m_k) - \ln\hat{\pi}_k$ by $\ln(s_k/(m_k\hat{\pi}_k))$, and likewise for the non-events. Multiplying by 2, as Equation 5 does, gives Equation 6. A count of 0 times $\ln 0$ is taken as 0, its limit. □

Remark 3: Source: deviance from grouped counts

Dobson and Barnett (2018, sec. 5.6.1) derives the deviance of a binomial model relative to the saturated model, and Section 7.5 writes it in the form of Equation 6. Section 7.5 also notes that when every observation has its own covariate pattern, so each count is 0 or 1, this deviance is not a useful measure of fit.

Corollary 0.2 (The deviance is the saturated-model deviance shifted by a constant)**.** *The deviance of a fitted logistic regression and its deviance relative to the saturated model differ by a constant that depends on the data and not on the fitted model:*

$$D = D_{\text{sat}} - 2\ell_{\text{sat}}.$$

Proof

Proof. Definition 0.2 gives $D = -2\ell\left(\underset{\sim}{\hat{\beta}}\right)$. Distributing the 2 in Equation 5 gives $D_{\text{sat}} = 2\ell_{\text{sat}} - 2\ell\left(\underset{\sim}{\hat{\beta}}\right)$. Substituting $D = -2\ell\left(\underset{\sim}{\hat{\beta}}\right)$ into this expression gives $D_{\text{sat}} = 2\ell_{\text{sat}} + D$. Subtracting $2\ell_{\text{sat}}$ from both sides gives the claim. □

Remark (Why the shift does not change the drop in deviance)

Remark 0.3 (Why the shift does not change the drop in deviance). The constant $2\ell_{\text{sat}}$ in Corollary 0.2 is the same for every model fit to the same data, because the saturated model depends only on the data. So the difference between the deviances of two models, such as the drop in deviance of Theorem 0.1, is the same whether it is computed from D or from D_{sat}. The two deviances differ in their size, though: when every pattern has one replicate, the saturated model fits every label exactly, $\ell_{\text{sat}} = 0$, and $D_{\text{sat}} = D$.

Example 0.7 (The deviance of the low-birth-weight model, ungrouped and grouped)**.** Fit a logistic regression of low birth weight on smoking and race, once to the 189 individual babies and once to the 6 grouped counts of Example 0.6. R's `deviance()` is the deviance of Definition 0.2 for the first fit and the deviance relative to the saturated model of Definition 0.8 for the second. We check the second against Equation 6 computed by hand.

```r
# fit to the individual babies
bw_fit <- glm(low ~ smoke + race, family = binomial, data = bw)
# fit to the grouped counts: events and non-events at each pattern
bw_fit_grouped <- glm(
  cbind(nevents, nreps - nevents) ~ smoke + race,
  family = binomial,
  data = bw_grouped
)
# fitted probability at each pattern
bw_phat <- fitted(bw_fit_grouped)
# one term of the grouped formula; a zero count contributes 0
bw_term <- function(count, expected) {
  ifelse(count == 0, 0, count * log(count / expected))
}
bw_sat_dev_hand <- 2 * sum(
  bw_term(bw_grouped$nevents, bw_grouped$nreps * bw_phat) +
    bw_term(
      bw_grouped$nreps - bw_grouped$nevents,
      bw_grouped$nreps * (1 - bw_phat)
    )
)
c(
  by_hand = bw_sat_dev_hand,
  glm_grouped = deviance(bw_fit_grouped),
  glm_ungrouped = deviance(bw_fit)
)
#>       by_hand   glm_grouped glm_ungrouped
#>       3.15694       3.15694     219.97471
```

The deviance relative to the saturated model is $D_{\text{sat}} = 3.157$ on the grouped counts, and the deviance is $D = 219.975$ on the individual babies.

Example 0.8 (The shift between the two deviances)**.** For the two fits of Example 0.7, Corollary 0.2 says that $D - D_{\text{sat}} = -2\ell_{\text{sat}}$. The left side is 216.818. We compute the right side from the observed proportions of Theorem 0.2.

```r
# twice the saturated log-likelihood, from the observed proportions
bw_phat_sat <- bw_grouped$nevents / bw_grouped$nreps
bw_two_llik_sat <- 2 * sum(
  ifelse(bw_grouped$nevents == 0, 0, bw_grouped$nevents * log(bw_phat_sat)) +
    ifelse(
      bw_grouped$nreps == bw_grouped$nevents,
      0,
      (bw_grouped$nreps - bw_grouped$nevents) * log(1 - bw_phat_sat)
    )
)
c(
  difference = deviance(bw_fit) - deviance(bw_fit_grouped),
  minus_two_llik_sat = -bw_two_llik_sat
)
#>         difference minus_two_llik_sat
#>            216.818            216.818
```

Confidence intervals for the coefficients

Theorem 0.4 (Approximate distribution of the fitted coefficients)**.** *Suppose the outcomes follow the logistic regression model. Let X be the matrix whose rows are $\underset{\sim}{x}_i^\top$, and let W be the diagonal matrix with entries $\pi_i(1-\pi_i)$. Under regularity conditions, for a large sample the fitted coefficients are approximately Gaussian,*

$$\hat{\underset{\sim}{\beta}} \approx \mathrm{N}\left(\underset{\sim}{\beta}, (X^\top W X)^{-1}\right). \tag{7}$$

Proof

Proof. The log-likelihood has score $X^\top\left(\underset{\sim}{y} - \underset{\sim}{\pi}\right)$, and its Hessian is $-X^\top W X$, which does not involve the outcomes. The observed and expected information are therefore both $X^\top W X$. The claim is the Gaussian approximation of the central limit theorem for MLEs, with this information in place of the inverse covariance. The result is stated for logistic regression by Hastie et al. (2009, sec. 4.4.3). For independent but non-identically distributed outcomes it holds under conditions on the predictors that Fahrmeir and Kaufmann (1985) give. □

Corollary 0.3 (Wald interval for a logistic coefficient)**.** *Let $\hat{W}$ be W with each π_i replaced by its fitted value. The estimated standard error of $\hat{\beta}_j$ is the square root of the diagonal entry of $\left(X^\top \hat{W} X\right)^{-1}$ in the row and column of β_j; because β_0 comes first, that is the $(j+1)$th diagonal entry counting from 1, and a Wald confidence interval with level $1-\alpha$ for β_j is*

$$\hat{\beta}_j \pm z_{1-\alpha/2}\, \widehat{\mathrm{se}}\left(\hat{\beta}_j\right).$$

Example 0.9 (A Wald interval from six patients)**.** For the fit of Example 0.1, `vcov` returns $\left(X^\top \hat{W} X\right)^{-1}$.

```
# standard errors: square roots of the diagonal of (X'WX)^-1
wf_se <- sqrt(diag(vcov(wf_fit)))
wf_slope <- coef(wf_fit)[["wf_x"]]
# 95% Wald interval for the slope
wf_ci <- wf_slope + c(-1, 1) * qnorm(0.975) * wf_se[["wf_x"]]
wf_summary <- c(
  slope = wf_slope, se = wf_se[["wf_x"]], lower = wf_ci[1], upper = wf_ci[2]
)
round(wf_summary, 3)
#>  slope     se  lower  upper
#>  0.693  0.866 -1.004  2.391
```

The interval for β_1 is $(-1, 2.39)$. It contains 0, consistent with the test of Example 0.5.

Corollary 0.4 (Interval for the odds factor of a logistic coefficient)**.** *If (L, U) is a confidence interval for β_j from Corollary 0.3, then (e^L, e^U) is a confidence interval with the same level for e^{β_j}, the factor by which a one-unit increase in x_j, holding the other predictors fixed, multiplies the odds of $y = 1$.*

Proof

Proof. The exponential function is increasing, so $L \le \beta_j \le U$ holds exactly when $e^L \le e^{\beta_j} \le e^U$. □

Example 0.10 (How little six patients tell us)**.** The two groups of Example 0.1 sit two units apart on the x scale, so the odds ratio between them is $e^{2\beta_1}$.

```r
# fitted group odds ratio, and its interval: exponentiate twice the slope
wf_group_or_hat <- exp(2 * wf_slope)
wf_group_or <- exp(2 * wf_ci)
c(estimate = wf_group_or_hat, lower = wf_group_or[1], upper = wf_group_or[2])
#>   estimate      lower      upper
#>   4.000000   0.134195 119.229587
```

The fitted odds of recovery in the treatment group are 4 times those in the control group, with interval $(0.13, 119.2)$. The interval runs from a strong harm to a large benefit. Its width shows how little information six observations carry about the slope.

References

Dobson, Annette J, and Adrian G Barnett. 2018. *An Introduction to Generalized Linear Models*. 4th ed. CRC press. https://doi.org/10.1201/9781315182780.

Fahrmeir, Ludwig, and Heinz Kaufmann. 1985. "Consistency and Asymptotic Normality of the Maximum Likelihood Estimator in Generalized Linear Models." *The Annals of Statistics* 13 (1): 342–68. https://doi.org/10.1214/aos/1176346597.

Hastie, Trevor, Robert Tibshirani, and Jerome Friedman. 2009. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. 2nd ed. Springer. https://doi.org/10.1007/978-0-387-84858-7.

Hosmer, David W., and Stanley Lemeshow. 1989. *Applied Logistic Regression*. Wiley.

Nelder, John A., and Robert W. M. Wedderburn. 1972. "Generalized Linear Models." *Journal of the Royal Statistical Society: Series A (General)* 135 (3): 370–84. https://doi.org/10.2307/2344614.