

# Correlation and Simple Linear Regression

Last modified: 2026-10-08 19:43:37 (PDT)

This page reviews two ways to relate two continuous variables: correlation coefficients, with tests of whether they differ from zero, and simple linear regression. It uses the  $t$  reference distribution defined on the [Statistical Inference](#) page. This page is adapted from Vittinghoff et al. (2012), Chapter 3.

## The HERS data

The examples on this page use the HERS data, which the [Comparing Means](#) page describes. The `rmb` R package includes the dataset; `haven::as_factor()` converts its Stata value labels to factors:

```
hers <- rmb::hers |> haven::as_factor()
hers |> head()
#> # A tibble: 6 x 37
#>   HT      age race nonwhite smoking drinkany exercise physact globrat poorfair
#>   <fct> <dbl> <fct>   <fct>   <fct>   <fct>   <fct>   <fct>   <fct>   <fct>
#> 1 plac~    70 Afric~ yes     no     no     no     much m~ good    no
#> 2 plac~    62 Afric~ yes     no     no     no     much l~ good    no
#> 3 horm~    69 White  no     no     no     no     about ~ good    no
#> 4 plac~    64 White  no     yes    yes    no     much l~ good    no
#> 5 plac~    65 White  no     no     no     no     somewh~ good    no
#> 6 horm~    68 Afric~ yes     no     yes    no     about ~ good    no
#> # i 27 more variables: medcond <dbl>, htnmeds <fct>, statins <fct>,
#> #   diabetes <fct>, dmpills <fct>, insulin <fct>, weight <dbl>, BMI <dbl>,
#> #   waist <dbl>, WHR <dbl>, glucose <dbl>, weight1 <dbl>, BMI1 <dbl>,
#> #   waist1 <dbl>, WHR1 <dbl>, glucose1 <dbl>, tchol <dbl>, LDL <dbl>,
#> #   HDL <dbl>, TG <dbl>, tchol1 <dbl>, LDL1 <dbl>, HDL1 <dbl>, TG1 <dbl>,
#> #   SBP <dbl>, DBP <dbl>, age10 <dbl>
```

## Correlation

### Testing the Pearson correlation

**Definition 0.1** (Population correlation). The **population correlation** of two random variables  $X$  and  $Y$  with positive variances is

$$\rho \stackrel{\text{def}}{=} \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}},$$

where  $\text{Cov}(X, Y)$  is their [covariance](#).

*Remark 0.1* (The sample correlation estimates the population correlation). The [Pearson correlation coefficient](#)  $r$  of a sample is the corresponding sample statistic, and estimates  $\rho$ .

**Definition 0.2** (Test of zero correlation). Let  $r$  be the [Pearson correlation coefficient](#) of  $n \geq 3$  pairs, with  $|r| < 1$ . The **t-test of zero correlation** of  $H_0 : \rho = 0$  against  $H_1 : \rho \neq 0$  (Definition 0.1) uses the statistic

$$t \stackrel{\text{def}}{=} \frac{r\sqrt{n-2}}{\sqrt{1-r^2}}.$$

Its p-value is  $\Pr(|T| \geq |t|)$ , where  $T$  has the  $t_{n-2}$  distribution ([t-distribution](#)).

**Theorem 0.1** (Null distribution of the correlation t statistic). *Let the pairs  $(X_1, Y_1), \dots, (X_n, Y_n)$  be independent, and let each  $Y_i$ , given  $X_1, \dots, X_n$ , be Gaussian with a mean and variance that do not depend on the  $X$  values. Then the statistic  $t$  of Definition 0.2 has the  $t_{n-2}$  distribution (Hogg et al. 2019, sec. 9.6, pp. 472-473):*

$$t \sim t_{n-2}.$$

*Remark 0.2* (When the conditions hold). The conditions of Theorem 0.1 hold, for example, when the pairs are independent draws from a bivariate Gaussian distribution with  $\rho = 0$ .

**Example 0.1** (Correlation between BMI and fasting glucose in HERS). Figure 1 plots baseline fasting glucose against BMI.

```

hers |>
  dplyr::filter(!is.na(BMI)) |>
  ggplot2::ggplot() +
  ggplot2::aes(x = BMI, y = glucose) +
  ggplot2::geom_point(alpha = 0.3) +
  ggplot2::geom_smooth(method = "lm", formula = y ~ x) +
  ggplot2::labs(
    x = "BMI (kg/m^2)",
    y = "Fasting glucose (mg/dL)"
  )

```



Figure 1: Baseline fasting glucose against BMI in HERS, with the least-squares line

The statistic of Definition 0.2, using the 2758 participants with a BMI measurement:

```

hers_bmi <- hers |> dplyr::filter(!is.na(BMI))
r <- cor(hers_bmi$BMI, hers_bmi$glucose)
n <- nrow(hers_bmi)
t_cor <- r * sqrt(n - 2) / sqrt(1 - r^2)
c(r = r, t = t_cor, p_value = 2 * pt(-abs(t_cor), df = n - 2))
#>           r           t      p_value
#> 2.72664e-01 1.48779e+01 3.24201e-48

```

`cor.test()` reports the same values, and a confidence interval for  $\rho$ :

```
cor.test(hers$BMI, hers$glucose, method = "pearson")
#>
#> Pearson's product-moment correlation
#>
#> data: hers$BMI and hers$glucose
#> t = 14.88, df = 2756, p-value <2e-16
#> alternative hypothesis: true correlation is not equal to 0
#> 95 percent confidence interval:
#> 0.237760 0.306865
#> sample estimates:
#> cor
#> 0.272664
```

The correlation is positive but modest: glucose tends to be higher at higher BMI, with wide scatter around the line in Figure 1.

## Spearman rank correlation

**Definition 0.3** (Spearman rank correlation). Replace each  $x_i$  by its rank  $u_i$  among  $x_1, \dots, x_n$ , and each  $y_i$  by its rank  $v_i$  among  $y_1, \dots, y_n$ , giving tied values the average of the ranks they span. The **Spearman rank correlation**  $r_S$  is the [Pearson correlation coefficient](#) evaluated at the rank pairs:

$$r_S \stackrel{\text{def}}{=} r((u_1, v_1), \dots, (u_n, v_n)).$$

*Remark 0.3* (What the Spearman correlation measures). Because ranks depend only on the ordering of the values,  $r_S$  measures how close the association is to monotone, whether or not it is linear, and a single extreme value moves  $r_S$  less than it moves  $r$ .

**Example 0.2** (Spearman correlation between BMI and fasting glucose in HERS). The Pearson correlation of the ranks (Definition 0.3), and `cor.test()`'s Spearman test (`exact = FALSE`, because tied values rule out the exact p-value):

```

hers_bmi <- hers |> dplyr::filter(!is.na(BMI))
cor(rank(hers_bmi$BMI), rank(hers_bmi$glucose))
#> [1] 0.333751
cor.test(hers_bmi$BMI, hers_bmi$glucose, method = "spearman", exact = FALSE)
#>
#> Spearman's rank correlation rho
#>
#> data:  hers_bmi$BMI and hers_bmi$glucose
#> S = 2.33e+09, p-value <2e-16
#> alternative hypothesis: true rho is not equal to 0
#> sample estimates:
#>      rho
#> 0.333751

```

$r_S$  is larger than the Pearson  $r$  of Example 0.1: the association is closer to monotone than to linear.

## Simple linear regression

### Model specification

**Definition 0.4** (Conditional Gaussian model). A **conditional Gaussian model** for outcomes  $Y_1, \dots, Y_n$  with covariate values  $x_1, \dots, x_n$  (each a single value or a vector) says that, given the covariates, the outcomes are independent and Gaussian: outcome  $Y_i$  is centered on a mean function  $\mu(x)$  evaluated at  $x_i$ , and has its own variance  $\sigma_i^2$ :

$$Y_i \mid X_i = x_i \sim_{\perp\!\!\!\perp} N(\mu_i, \sigma_i^2),$$

$$\mu_i \stackrel{\text{def}}{=} \mu(x_i).$$

**Definition 0.5** (Deviation from the conditional mean). In a conditional Gaussian model (Definition 0.4), the **deviation**  $\varepsilon_i$  of outcome  $Y_i$  is its **deviation from its mean**, taking the mean conditional on  $X_i = x_i$ :

$$\varepsilon_i \stackrel{\text{def}}{=} Y_i - \mu(x_i).$$

**Exercise 0.1** (Outcome as mean plus deviation). Show that each outcome in a conditional Gaussian model (Definition 0.4) is its conditional mean plus its deviation (Definition 0.5):  $Y_i = \mu_i + \varepsilon_i$ .

## Solution

*Solution 0.1.* Start from the right-hand side:

$$\begin{aligned}\mu_i + \varepsilon_i &= \mu(x_i) + \varepsilon_i && \text{(definition of } \mu_i\text{)} \\ &= \mu(x_i) + (Y_i - \mu(x_i)) && \text{(definition of } \varepsilon_i\text{)} \\ &= \mu(x_i) + Y_i - \mu(x_i) && \text{(remove the parentheses)} \\ &= Y_i + \mu(x_i) - \mu(x_i) && \text{(reorder the terms)} \\ &= Y_i + (\mu(x_i) - \mu(x_i)) && \text{(group the last two terms)} \\ &= Y_i + 0 && (a - a = 0) \\ &= Y_i && (a + 0 = a)\end{aligned}$$

So  $Y_i = \mu_i + \varepsilon_i$ .

Alternatively, solve the definition of  $\varepsilon_i$  for  $Y_i$ :

$$\begin{aligned}\varepsilon_i &= Y_i - \mu(x_i) && \text{(definition of } \varepsilon_i\text{)} \\ \varepsilon_i + \mu(x_i) &= Y_i - \mu(x_i) + \mu(x_i) && \text{(add } \mu(x_i) \text{ to both sides)} \\ \varepsilon_i + \mu(x_i) &= Y_i + (-\mu(x_i)) + \mu(x_i) && \text{(subtracting is adding the negative)} \\ \varepsilon_i + \mu(x_i) &= Y_i + (-\mu(x_i) + \mu(x_i)) && \text{(group the last two terms)} \\ \varepsilon_i + \mu(x_i) &= Y_i + 0 && (-a + a = 0) \\ \varepsilon_i + \mu(x_i) &= Y_i && (a + 0 = a) \\ Y_i &= \varepsilon_i + \mu(x_i) && \text{(swap the two sides)} \\ Y_i &= \mu(x_i) + \varepsilon_i && \text{(reorder the terms)} \\ Y_i &= \mu_i + \varepsilon_i && \text{(definition of } \mu_i\text{)}\end{aligned}$$

**Theorem 0.2** (Additive form of the conditional Gaussian model). *In a conditional Gaussian model (Definition 0.4), each outcome is its conditional mean plus its deviation (Definition 0.5):*

$$Y_i = \mu_i + \varepsilon_i.$$

## i Proof

*Proof.* This is the solution to Exercise 0.1. □

**Exercise 0.2** (Distribution of the deviations). Show that in a conditional Gaussian model (Definition 0.4), given the covariate values, the deviations (Definition 0.5) are independent and each is Gaussian with mean 0 and the same variance  $\sigma_i^2$  as  $Y_i$ :  $\varepsilon_i \mid X_i = x_i \sim \perp$

$N(0, \sigma_i^2)$ .

### Solution

*Solution 0.2.* Every probability, CDF and density in this solution is [conditional](#) on  $X_i = x_i$ ; the condition is left out of the notation to keep the lines short. Write  $F_\varepsilon$  and  $f_\varepsilon$  for the CDF and density of  $\varepsilon_i$ , and  $F_Y$  and  $f_Y$  for those of  $Y_i$ .

First, the CDF of  $\varepsilon_i$ :

$$\begin{aligned} F_\varepsilon(t) &= \Pr(\varepsilon_i \leq t) && \text{(definition of a CDF)} \\ &= \Pr(Y_i - \mu(x_i) \leq t) && \text{(definition of } \varepsilon_i \text{)} \\ &= \Pr(Y_i - \mu_i \leq t) && \text{(definition of } \mu_i \text{)} \\ &= \Pr(Y_i \leq t + \mu_i) && \text{(add } \mu_i \text{ to both sides of the inequality)} \\ &= F_Y(t + \mu_i) && \text{(definition of a CDF)} \end{aligned}$$

Differentiating gives the density:

$$\begin{aligned} f_\varepsilon(t) &= \frac{\partial}{\partial t} F_\varepsilon(t) && \text{(a density is the derivative of its CDF)} \\ &= \frac{\partial}{\partial t} F_Y(t + \mu_i) && \text{(CDF of } \varepsilon_i \text{)} \\ &= f_Y(t + \mu_i) \frac{\partial}{\partial t} (t + \mu_i) && \text{(chain rule)} \end{aligned}$$

The inner derivative:

$$\begin{aligned} \frac{\partial}{\partial t} (t + \mu_i) &= \frac{\partial}{\partial t} t + \frac{\partial}{\partial t} \mu_i && \text{(derivative of a sum)} \\ &= 1 + 0 && (\mu_i \text{ does not depend on } t) \\ &= 1 && \text{(add)} \end{aligned}$$

Plugging the inner derivative back in, and using the [Gaussian density](#) of  $Y_i$ , which has mean  $\mu_i = \mu(x_i)$  and variance  $\sigma_i^2$  (Definition 0.4):

$$\begin{aligned} f_\varepsilon(t) &= f_Y(t + \mu_i) \cdot 1 && \text{(inner derivative is 1)} \\ &= f_Y(t + \mu_i) && \text{(multiply by 1)} \\ &= \frac{1}{\sigma_i \sqrt{2\pi}} e^{-\frac{((t+\mu_i)-\mu_i)^2}{2\sigma_i^2}} && \text{(Gaussian density with mean } \mu_i \text{)} \\ &= \frac{1}{\sigma_i \sqrt{2\pi}} e^{-\frac{t^2}{2\sigma_i^2}} && \text{(subtract)} \end{aligned}$$

The last line is the density of  $N(0, \sigma_i^2)$ , so  $\varepsilon_i \mid X_i = x_i \sim N(0, \sigma_i^2)$ .

For independence, factor the joint CDF of  $\varepsilon_1, \dots, \varepsilon_n$ , using the same CDF step as above for each  $i$  and the independence of the  $Y_i$  given the covariates (Definition 0.4):

$$\begin{aligned}
\Pr(\varepsilon_1 \leq t_1, \dots, \varepsilon_n \leq t_n) &= \Pr(Y_1 \leq t_1 + \mu_1, \dots, Y_n \leq t_n + \mu_n) && \text{(add } \mu_i \text{ to both sides of each inequality)} \\
&= \prod_{i=1}^n \Pr(Y_i \leq t_i + \mu_i) && \text{(the } Y_i \text{ are independent)} \\
&= \prod_{i=1}^n \Pr(\varepsilon_i \leq t_i) && \text{(subtract } \mu_i \text{ from both sides of each inequality)}
\end{aligned}$$

The joint CDF is the product of the marginal CDFs, so  $\varepsilon_1, \dots, \varepsilon_n$  are independent.

**Theorem 0.3** (Distribution of the deviations). *In a conditional Gaussian model (Definition 0.4), given the covariates, the deviations (Definition 0.5) are independent and Gaussian with mean 0 and variance  $\sigma_i^2$ :*

$$\varepsilon_i \mid X_i = x_i \sim_{\perp\!\!\!\perp} N(0, \sigma_i^2).$$

**i** Proof

*Proof.* This is the solution to Exercise 0.2. □

**Definition 0.6** (Homoskedastic model). A conditional Gaussian model (Definition 0.4) is **homoskedastic** if all outcomes share one variance  $\sigma^2$ , which does not depend on  $i$ :

$$\sigma_i^2 = \sigma^2 \text{ for all } i.$$

**Corollary 0.1** (Deviations in a homoskedastic model). *In a homoskedastic model (Definition 0.6), the deviations (Definition 0.5) all have the same variance  $\sigma^2$ :*

$$\varepsilon_i \mid X_i = x_i \sim_{\perp\!\!\!\perp} N(0, \sigma^2).$$

**i** Proof

*Proof.* By Theorem 0.3,  $\varepsilon_i \mid X_i = x_i \sim_{\perp\!\!\!\perp} N(0, \sigma_i^2)$ , and Definition 0.6 sets  $\sigma_i^2 = \sigma^2$  for all  $i$ . □

**Definition 0.7** (Linear regression model). A **linear regression model** is a homoskedastic (Definition 0.6) conditional Gaussian model (Definition 0.4) whose mean function is linear in  $p$  covariates, where observation  $i$  has covariate values  $x_i = (x_{i1}, \dots, x_{ip})$ :

$$\mu(x_1, \dots, x_p) \stackrel{\text{def}}{=} \beta_0 + \sum_{j=1}^p \beta_{x_j} x_j.$$

**Definition 0.8** (Simple linear regression model). A **simple linear regression model** is a linear regression model (Definition 0.7) with  $p = 1$  covariate  $X$ , writing  $x_i$  for  $x_{i1}$ :

$$\mu(x) \stackrel{\text{def}}{=} \beta_0 + \beta_x x.$$

*Remark 0.4* (Interpreting the parameters).

- $\beta_0 = \mu(0)$  is the **intercept**: the mean of  $Y$  among observations with  $X = 0$ .
- $\beta_x = \mu(x + 1) - \mu(x)$  is the **slope**: the difference in the mean of  $Y$  between two groups whose values of  $X$  differ by one unit.
- $\sigma^2$  is the variance of  $Y$  around its mean at each value of  $X$ .

## Ordinary least squares estimation

**Definition 0.9** (Residual sum of squares). For a regression model whose conditional mean  $\mu(x; \tilde{\theta})$  depends on a parameter vector  $\tilde{\theta}$ , the **residual sum of squares** at  $\tilde{\theta}$  is the sum of the squared residuals  $r_i(\tilde{\theta}) \stackrel{\text{def}}{=} y_i - \mu(x_i; \tilde{\theta})$  that the model has when its parameters equal  $\tilde{\theta}$ :

$$\text{RSS}(\tilde{\theta}) \stackrel{\text{def}}{=} \sum_{i=1}^n r_i(\tilde{\theta})^2.$$

At the estimate  $\hat{\tilde{\theta}}$ ,  $r_i(\hat{\tilde{\theta}})$  is the **residual**  $r_i$  of the fitted model.

**Example 0.3** (Residual sum of squares of a simple linear regression). The mean function of a **simple linear regression** model has parameter vector  $\tilde{\theta} = (\beta_0, \beta_x)$ , so write  $\text{RSS}(\beta_0, \beta_x)$  for its residual sum of squares (Definition 0.9):

$$\begin{aligned}
\text{RSS}(\beta_0, \beta_x) &= \sum_{i=1}^n r_i(\beta_0, \beta_x)^2 && \text{(definition of RSS)} \\
&= \sum_{i=1}^n (y_i - \mu(x_i; \beta_0, \beta_x))^2 && \text{(definition of } r_i(\tilde{\theta})\text{)} \\
&= \sum_{i=1}^n (y_i - (\beta_0 + \beta_x x_i))^2 && \text{(definition of } \mu(x)\text{)} \\
&= \sum_{i=1}^n (y_i - \beta_0 - \beta_x x_i)^2 && \text{(distribute the minus sign)}
\end{aligned}$$

**Example 0.4** (Residual sum of squares of a line through three points). For the points  $(0, 1)$ ,  $(1, 2)$ ,  $(2, 2)$  and the line with  $\beta_0 = 1$  and  $\beta_x = 0.5$  (Example 0.3), the residuals are  $1 - 1 = 0$ ,  $2 - 1.5 = 0.5$ , and  $2 - 2 = 0$ , so  $\text{RSS}(1, 0.5) = 0^2 + 0.5^2 + 0^2 = 0.25$  (Figure 2).

```

rss_points <- tibble::tibble(x = c(0, 1, 2), y = c(1, 2, 2)) |>
  dplyr::mutate(fitted = 1 + 0.5 * x)

ggplot2::ggplot(rss_points, ggplot2::aes(x = x, y = y)) +
  ggplot2::geom_abline(intercept = 1, slope = 0.5) +
  ggplot2::geom_segment(
    ggplot2::aes(xend = x, yend = fitted),
    linetype = "dashed",
    color = "firebrick"
  ) +
  ggplot2::geom_point(size = 3) +
  ggplot2::coord_cartesian(xlim = c(-0.25, 2.25), ylim = c(0.75, 2.25))

```



Figure 2: The three points, the line with intercept 1 and slope 0.5, and the one nonzero residual (dashed)

**Exercise 0.3** (Mean squared error and residual sum of squares). For a regression model with conditional mean  $\mu(x; \tilde{\theta})$ , let  $\hat{y}_i \stackrel{\text{def}}{=} \mu(x_i; \tilde{\theta})$  be its predictions of  $n \geq 1$  outcomes  $y_1, \dots, y_n$ , and let  $\text{RSS}(\tilde{\theta})$  (Definition 0.9) be taken over those same outcomes. Write the [mean squared error](#) of these predictions in terms of  $\text{RSS}(\tilde{\theta})$ , and  $\text{RSS}(\tilde{\theta})$  in terms of that mean squared error.

## Solution

*Solution 0.3.* First, each squared **prediction error** equals the squared residual at  $\tilde{\theta}$ :

$$\begin{aligned} e_i^2 &= (\hat{y}_i - y_i)^2 && \text{(definition of prediction error)} \\ &= (\mu(x_i; \tilde{\theta}) - y_i)^2 && \text{(definition of } \hat{y}_i) \\ &= (- (y_i - \mu(x_i; \tilde{\theta})))^2 && \text{(factor out } -1) \\ &= (-r_i(\tilde{\theta}))^2 && \text{(definition of } r_i(\tilde{\theta})) \\ &= r_i(\tilde{\theta})^2 && ((-a)^2 = a^2) \end{aligned}$$

So the mean squared error is the residual sum of squares divided by  $n$ :

$$\begin{aligned} \text{MSE}(\hat{y}) &= \frac{1}{n} \sum_{i=1}^n e_i^2 && \text{(definition of mean squared error)} \\ &= \frac{1}{n} \sum_{i=1}^n r_i(\tilde{\theta})^2 && (e_i^2 = r_i(\tilde{\theta})^2) \\ &= \frac{1}{n} \text{RSS}(\tilde{\theta}) && \text{(definition of RSS)} \end{aligned}$$

Solving for  $\text{RSS}(\tilde{\theta})$ :

$$\begin{aligned} \text{MSE}(\hat{y}) &= \frac{1}{n} \text{RSS}(\tilde{\theta}) && \text{(result above)} \\ n \text{ MSE}(\hat{y}) &= \text{RSS}(\tilde{\theta}) && \text{(multiply both sides by } n) \\ \text{RSS}(\tilde{\theta}) &= n \text{ MSE}(\hat{y}) && \text{(swap the two sides)} \end{aligned}$$

**Theorem 0.4** (Mean squared error is the residual sum of squares divided by  $n$ ). *For a regression model with conditional mean  $\mu(x; \tilde{\theta})$ , the **mean squared error** of its predictions  $\hat{y}_i = \mu(x_i; \tilde{\theta})$  of  $n \geq 1$  outcomes  $y_1, \dots, y_n$  and the residual sum of squares  $\text{RSS}(\tilde{\theta})$  (Definition 0.9) over those same  $n$  outcomes satisfy*

$$\text{MSE}(\hat{y}) = \frac{1}{n} \text{RSS}(\tilde{\theta}),$$

*and equivalently*

$$\text{RSS}(\tilde{\theta}) = n \text{ MSE}(\hat{y}).$$

**i** Proof

*Proof.* Both equations are derived in Exercise 0.3. □

*Remark 0.5* (The same observations on both sides). The identities in Theorem 0.4 need the mean squared error and the residual sum of squares to be taken over the same  $n$  observations. The RSS minimized in fitting is taken over the fitting data, so dividing it by  $n$  gives the mean squared error on the fitting data, not the mean squared error of predictions of new outcomes.

**Definition 0.10** (Ordinary least squares). The **ordinary least squares (OLS) estimate** of a model's parameter vector  $\tilde{\theta}$  is the value of  $\tilde{\theta}$  that minimizes the **residual sum of squares**:

$$\hat{\theta} \stackrel{\text{def}}{=} \arg \min_{\tilde{\theta}} \text{RSS}(\tilde{\theta}).$$

**Example 0.5** (OLS estimates of a simple linear regression). For a simple linear regression, the OLS estimates (Definition 0.10)  $\hat{\beta}_0$  and  $\hat{\beta}_x$  are the values of  $\beta_0$  and  $\beta_x$  that minimize  $\text{RSS}(\beta_0, \beta_x)$  from Example 0.3. Figure 3 shows  $\text{RSS}(\beta_0, \beta_x)$  for the three points of Example 0.4: the OLS estimates sit at the bottom of the bowl, and the line of Example 0.4 sits higher up.

```

rss_grid <- expand.grid(
  b0 = seq(0, 2.5, length.out = 101),
  bx = seq(-0.5, 1.5, length.out = 101)
)
rss_grid$rss <- mapply(
  function(b0, bx) sum((rss_points$y - b0 - bx * rss_points$x)^2),
  rss_grid$b0,
  rss_grid$bx
)
ols_coefs <- coef(lm(y ~ x, data = rss_points))

ggplot2::ggplot(rss_grid, ggplot2::aes(x = b0, y = bx, z = rss)) +
  ggplot2::geom_contour(
    ggplot2::aes(color = ggplot2::after_stat(level)),
    breaks = c(0.25, 0.5, 1, 2, 4, 8)
  ) +
  ggplot2::annotate(
    "point",
    x = ols_coefs[[1]], y = ols_coefs[[2]], size = 3
  ) +
  ggplot2::annotate("point", x = 1, y = 0.5, shape = 4, size = 3) +
  ggplot2::labs(
    x = expression(beta[0]),
    y = expression(beta[x]),
    color = "RSS"
  )

```



```

ols_coefs
#> (Intercept)      x
#>   1.16667    0.50000

```

Figure 3: Contours of  $\text{RSS}(\beta_0, \beta_x)$  for the three points of Example 0.4. The dot marks

**Definition 0.11** (Normal equations). For a model fitted by least squares with parameter vector  $\tilde{\theta}$ , the **normal equations** set the gradient of the **residual sum of squares** to zero:

$$\frac{\partial}{\partial \tilde{\theta}} \text{RSS}(\tilde{\theta}) = \tilde{0}.$$

**Example 0.6** (Normal equations of a simple linear regression). For a simple linear regression, the parameter vector is  $\tilde{\beta} \stackrel{\text{def}}{=} (\beta_0, \beta_x)^\top$ , and  $\text{RSS}(\tilde{\beta})$  is  $\text{RSS}(\beta_0, \beta_x)$  from Example 0.3. Its gradient has one entry per coefficient, so the normal equations (Definition 0.11) are two scalar equations:

$$\frac{\partial \text{RSS}}{\partial \beta_0} = 0, \quad \frac{\partial \text{RSS}}{\partial \beta_x} = 0.$$

**Definition 0.12** (Centered sums of squares and cross-products). For data  $(x_1, y_1), \dots, (x_n, y_n)$ , the **centered sums of squares** of the  $x_i$  and of the  $y_i$ , and their **centered sum of cross-products**, are

$$S_{xx} \stackrel{\text{def}}{=} \sum_{i=1}^n (x_i - \bar{x})^2, \quad S_{yy} \stackrel{\text{def}}{=} \sum_{i=1}^n (y_i - \bar{y})^2, \quad S_{xy} \stackrel{\text{def}}{=} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}).$$

**Exercise 0.4** (Deviations from the mean sum to zero). Show that  $\sum_{i=1}^n (x_i - \bar{x}) = 0$ .

Solution

*Solution 0.4.*

$$\begin{aligned} \sum_{i=1}^n (x_i - \bar{x}) &= \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \quad (\text{split the sum}) \\ &= \sum_{i=1}^n x_i - n\bar{x} \quad (\text{sum of a constant}) \\ &= n\bar{x} - n\bar{x} \quad \left( \sum_{i=1}^n x_i = n\bar{x} \right) \\ &= 0 \quad (\text{subtract}) \end{aligned}$$

The same steps with  $y_i$  in place of  $x_i$  show that  $\sum_{i=1}^n (y_i - \bar{y}) = 0$ .

**Exercise 0.5** (Expanding the centered sum of squares). Show that  $S_{xx} = \sum_{i=1}^n x_i^2 - n\bar{x}^2$  (Definition 0.12).

### Solution

*Solution 0.5.*

$$\begin{aligned}
 S_{xx} &= \sum_{i=1}^n (x_i - \bar{x})^2 && \text{(definition of } S_{xx}) \\
 &= \sum_{i=1}^n (x_i^2 - 2\bar{x}x_i + \bar{x}^2) && \text{(expand the square)} \\
 &= \sum_{i=1}^n x_i^2 - \sum_{i=1}^n 2\bar{x}x_i + \sum_{i=1}^n \bar{x}^2 && \text{(split the sum)} \\
 &= \sum_{i=1}^n x_i^2 - 2\bar{x} \sum_{i=1}^n x_i + \sum_{i=1}^n \bar{x}^2 && \text{(constant factor out of the sum)} \\
 &= \sum_{i=1}^n x_i^2 - 2\bar{x} \sum_{i=1}^n x_i + n\bar{x}^2 && \text{(sum of a constant)} \\
 &= \sum_{i=1}^n x_i^2 - 2\bar{x} \cdot n\bar{x} + n\bar{x}^2 && \left(\sum_{i=1}^n x_i = n\bar{x}\right) \\
 &= \sum_{i=1}^n x_i^2 - 2n\bar{x}^2 + n\bar{x}^2 && \text{(multiply)} \\
 &= \sum_{i=1}^n x_i^2 - n\bar{x}^2 && \text{(collect the } \bar{x}^2 \text{ terms)}
 \end{aligned}$$

**Exercise 0.6** (Derivative of RSS with respect to the intercept). Find the partial derivative of the simple linear regression residual sum of squares  $\text{RSS}(\beta_0, \beta_x)$  (Example 0.3) with respect to the intercept,  $\partial \text{RSS} / \partial \beta_0$ , and write it in terms of  $\bar{x}$  and  $\bar{y}$ .

### Solution

*Solution 0.6.* Differentiate one operation at a time, starting from the definition (Example 0.3):

$$\begin{aligned}
 \frac{\partial \text{RSS}}{\partial \beta_0} &= \frac{\partial}{\partial \beta_0} \sum_{i=1}^n (y_i - \beta_0 - \beta_x x_i)^2 && \text{(definition of RSS)} \\
 &= \sum_{i=1}^n \frac{\partial}{\partial \beta_0} (y_i - \beta_0 - \beta_x x_i)^2 && \text{(derivative of a sum)} \\
 &= \sum_{i=1}^n 2(y_i - \beta_0 - \beta_x x_i) \frac{\partial}{\partial \beta_0} (y_i - \beta_0 - \beta_x x_i) && \text{(chain rule)}
 \end{aligned}$$

The inner derivative:

$$\begin{aligned}
\frac{\partial}{\partial \beta_0}(y_i - \beta_0 - \beta_x x_i) &= \frac{\partial y_i}{\partial \beta_0} - \frac{\partial \beta_0}{\partial \beta_0} - \frac{\partial(\beta_x x_i)}{\partial \beta_0} && \text{(derivative of a sum)} \\
&= 0 - 1 - 0 && (y_i \text{ and } \beta_x x_i \text{ do not depend on } \beta_0) \\
&= -1 && \text{(subtract)}
\end{aligned}$$

Plugging the inner derivative back in:

$$\begin{aligned}
\frac{\partial \text{RSS}}{\partial \beta_0} &= \sum_{i=1}^n 2(y_i - \beta_0 - \beta_x x_i)(-1) && \text{(inner derivative is } -1) \\
&= \sum_{i=1}^n -2(y_i - \beta_0 - \beta_x x_i) && \text{(multiply 2 by } -1) \\
&= -2 \sum_{i=1}^n (y_i - \beta_0 - \beta_x x_i) && \text{(constant factor out of the sum)} \\
&= -2 \left( \sum_{i=1}^n y_i - \sum_{i=1}^n \beta_0 - \sum_{i=1}^n \beta_x x_i \right) && \text{(split the sum)} \\
&= -2 \left( \sum_{i=1}^n y_i - n\beta_0 - \sum_{i=1}^n \beta_x x_i \right) && \text{(sum of a constant)} \\
&= -2 \left( \sum_{i=1}^n y_i - n\beta_0 - \beta_x \sum_{i=1}^n x_i \right) && \text{(constant factor out of the sum)} \\
&= -2 \left( n\bar{y} - n\beta_0 - \beta_x \sum_{i=1}^n x_i \right) && \left( \sum_{i=1}^n y_i = n\bar{y} \right) \\
&= -2(n\bar{y} - n\beta_0 - \beta_x n\bar{x}) && \left( \sum_{i=1}^n x_i = n\bar{x} \right)
\end{aligned}$$

**Exercise 0.7** (Derivative of RSS with respect to the slope). Find the partial derivative of the simple linear regression residual sum of squares  $\text{RSS}(\beta_0, \beta_x)$  (Example 0.3) with respect to the slope,  $\partial \text{RSS} / \partial \beta_x$ .

**Solution**

*Solution 0.7.* Differentiate one operation at a time, starting from the definition (Example 0.3):

$$\begin{aligned}
\frac{\partial \text{RSS}}{\partial \beta_x} &= \frac{\partial}{\partial \beta_x} \sum_{i=1}^n (y_i - \beta_0 - \beta_x x_i)^2 && \text{(definition of RSS)} \\
&= \sum_{i=1}^n \frac{\partial}{\partial \beta_x} (y_i - \beta_0 - \beta_x x_i)^2 && \text{(derivative of a sum)} \\
&= \sum_{i=1}^n 2(y_i - \beta_0 - \beta_x x_i) \frac{\partial}{\partial \beta_x} (y_i - \beta_0 - \beta_x x_i) && \text{(chain rule)}
\end{aligned}$$

The inner derivative:

$$\begin{aligned}
\frac{\partial}{\partial \beta_x} (y_i - \beta_0 - \beta_x x_i) &= \frac{\partial y_i}{\partial \beta_x} - \frac{\partial \beta_0}{\partial \beta_x} - \frac{\partial (\beta_x x_i)}{\partial \beta_x} && \text{(derivative of a sum)} \\
&= 0 - 0 - x_i && (y_i \text{ and } \beta_0 \text{ do not depend on } \beta_x) \\
&= -x_i && \text{(subtract)}
\end{aligned}$$

Plugging the inner derivative back in:

$$\begin{aligned}
\frac{\partial \text{RSS}}{\partial \beta_x} &= \sum_{i=1}^n 2(y_i - \beta_0 - \beta_x x_i)(-x_i) && \text{(inner derivative is } -x_i) \\
&= \sum_{i=1}^n -2x_i(y_i - \beta_0 - \beta_x x_i) && \text{(rearrange the product)} \\
&= -2 \sum_{i=1}^n x_i(y_i - \beta_0 - \beta_x x_i) && \text{(constant factor out of the sum)}
\end{aligned}$$

**Exercise 0.8** (Solving the normal equations). Using Exercise 0.6 and Exercise 0.7, solve the normal equations of Example 0.6 for  $\beta_0$  and  $\beta_x$ , assuming  $S_{xx} > 0$  (Definition 0.12).

**Solution**

*Solution 0.8. First normal equation.* Set the derivative from Exercise 0.6 to zero, then isolate  $\beta_0$ :

$$\begin{aligned}
0 &= -2(n\bar{y} - n\beta_0 - \beta_x n\bar{x}) && \text{(first normal equation)} \\
0 &= n\bar{y} - n\beta_0 - \beta_x n\bar{x} && \text{(divide both sides by } -2) \\
0 &= \bar{y} - \beta_0 - \beta_x \bar{x} && \text{(divide both sides by } n) \\
\beta_0 &= \bar{y} - \beta_x \bar{x} && \text{(add } \beta_0 \text{ to both sides)}
\end{aligned}$$

**Second normal equation.** Substitute that intercept into the derivative from Exercise 0.7:

$$\begin{aligned}
\frac{\partial \text{RSS}}{\partial \beta_x} &= -2 \sum_{i=1}^n x_i (y_i - \beta_0 - \beta_x x_i) && \text{(derivative with respect to } \beta_x) \\
&= -2 \sum_{i=1}^n x_i (y_i - (\bar{y} - \beta_x \bar{x}) - \beta_x x_i) && \text{(substitute } \beta_0) \\
&= -2 \sum_{i=1}^n x_i (y_i - \bar{y} + \beta_x \bar{x} - \beta_x x_i) && \text{(remove the inner parentheses)} \\
&= -2 \sum_{i=1}^n x_i ((y_i - \bar{y}) - (\beta_x x_i - \beta_x \bar{x})) && \text{(group terms)} \\
&= -2 \sum_{i=1}^n x_i ((y_i - \bar{y}) - \beta_x (x_i - \bar{x})) && \text{(factor out } \beta_x)
\end{aligned}$$

To shorten the next steps, write  $d_i \stackrel{\text{def}}{=} (y_i - \bar{y}) - \beta_x (x_i - \bar{x})$ , so that  $\partial \text{RSS} / \partial \beta_x = -2 \sum_{i=1}^n x_i d_i$ . The  $d_i$  sum to zero, because deviations from a mean sum to zero (Exercise 0.4):

$$\begin{aligned}
\sum_{i=1}^n d_i &= \sum_{i=1}^n ((y_i - \bar{y}) - \beta_x (x_i - \bar{x})) && \text{(definition of } d_i) \\
&= \sum_{i=1}^n (y_i - \bar{y}) - \sum_{i=1}^n \beta_x (x_i - \bar{x}) && \text{(split the sum)} \\
&= \sum_{i=1}^n (y_i - \bar{y}) - \beta_x \sum_{i=1}^n (x_i - \bar{x}) && \text{(constant factor out of the sum)} \\
&= 0 - \beta_x \cdot 0 && \text{(deviations sum to zero)} \\
&= 0 - 0 && \text{(multiply)} \\
&= 0 && \text{(subtract)}
\end{aligned}$$

So subtracting  $\bar{x} \sum_{i=1}^n d_i$  from  $\sum_{i=1}^n x_i d_i$  does not change it:

$$\begin{aligned}
\sum_{i=1}^n x_i d_i &= \sum_{i=1}^n x_i d_i - \bar{x} \sum_{i=1}^n d_i && (\sum_{i=1}^n d_i = 0) \\
&= \sum_{i=1}^n x_i d_i - \sum_{i=1}^n \bar{x} d_i && (\text{constant factor into the sum}) \\
&= \sum_{i=1}^n (x_i - \bar{x}) d_i && (\text{combine the sums}) \\
&= \sum_{i=1}^n (x_i - \bar{x}) ((y_i - \bar{y}) - \beta_x (x_i - \bar{x})) && (\text{definition of } d_i) \\
&= \sum_{i=1}^n ((x_i - \bar{x})(y_i - \bar{y}) - \beta_x (x_i - \bar{x})^2) && (\text{distribute}) \\
&= \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) - \sum_{i=1}^n \beta_x (x_i - \bar{x})^2 && (\text{split the sum}) \\
&= \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) - \beta_x \sum_{i=1}^n (x_i - \bar{x})^2 && (\text{constant factor out of the sum}) \\
&= S_{xy} - \beta_x \sum_{i=1}^n (x_i - \bar{x})^2 && (\text{definition of } S_{xy}) \\
&= S_{xy} - \beta_x S_{xx} && (\text{definition of } S_{xx})
\end{aligned}$$

So  $\partial \text{RSS} / \partial \beta_x = -2(S_{xy} - \beta_x S_{xx})$ . Set it to zero and isolate  $\beta_x$ :

$$\begin{aligned}
0 &= -2(S_{xy} - \beta_x S_{xx}) && (\text{second normal equation}) \\
0 &= S_{xy} - \beta_x S_{xx} && (\text{divide both sides by } -2) \\
\beta_x S_{xx} &= S_{xy} && (\text{add } \beta_x S_{xx} \text{ to both sides}) \\
\beta_x &= \frac{S_{xy}}{S_{xx}} && (\text{divide both sides by } S_{xx} > 0)
\end{aligned}$$

So the normal equations have exactly one solution:  $\beta_x = S_{xy}/S_{xx}$  and  $\beta_0 = \bar{y} - \beta_x \bar{x}$ .

**Exercise 0.9** (Second derivatives of RSS). Find the matrix of second derivatives of  $\text{RSS}(\beta_0, \beta_x)$  (Example 0.3) with respect to  $\beta_0$  and  $\beta_x$ , and show that it is positive definite when  $S_{xx} > 0$  (Definition 0.12).

**Solution**

*Solution 0.9.* Differentiate each first derivative again.

**Derivatives of  $\partial \text{RSS} / \partial \beta_0$ .** By Exercise 0.6,  $\partial \text{RSS} / \partial \beta_0 = -2(n\bar{y} - n\beta_0 - \beta_x n\bar{x})$ . Differentiating with respect to  $\beta_0$ :

$$\begin{aligned}
\frac{\partial^2 \text{RSS}}{\partial \beta_0^2} &= -2 \frac{\partial}{\partial \beta_0} (n\bar{y} - n\beta_0 - \beta_x n\bar{x}) && \text{(constant factor)} \\
&= -2 \left( \frac{\partial(n\bar{y})}{\partial \beta_0} - \frac{\partial(n\beta_0)}{\partial \beta_0} - \frac{\partial(\beta_x n\bar{x})}{\partial \beta_0} \right) && \text{(derivative of a sum)} \\
&= -2(0 - n - 0) && \text{(differentiate each term)} \\
&= -2(-n) && \text{(drop the zeros)} \\
&= 2n && \text{(multiply)}
\end{aligned}$$

Differentiating with respect to  $\beta_x$ :

$$\begin{aligned}
\frac{\partial^2 \text{RSS}}{\partial \beta_x \partial \beta_0} &= -2 \frac{\partial}{\partial \beta_x} (n\bar{y} - n\beta_0 - \beta_x n\bar{x}) && \text{(constant factor)} \\
&= -2 \left( \frac{\partial(n\bar{y})}{\partial \beta_x} - \frac{\partial(n\beta_0)}{\partial \beta_x} - \frac{\partial(\beta_x n\bar{x})}{\partial \beta_x} \right) && \text{(derivative of a sum)} \\
&= -2(0 - 0 - n\bar{x}) && \text{(differentiate each term)} \\
&= -2(-n\bar{x}) && \text{(drop the zeros)} \\
&= 2n\bar{x} && \text{(multiply)}
\end{aligned}$$

**Derivatives of  $\partial \text{RSS} / \partial \beta_x$ .** By Exercise 0.7,  $\partial \text{RSS} / \partial \beta_x = -2 \sum_{i=1}^n x_i (y_i - \beta_0 - \beta_x x_i)$ . Differentiating with respect to  $\beta_0$ , and reusing the inner derivative  $-1$  from Exercise 0.6:

$$\begin{aligned}
\frac{\partial^2 \text{RSS}}{\partial \beta_0 \partial \beta_x} &= -2 \frac{\partial}{\partial \beta_0} \sum_{i=1}^n x_i (y_i - \beta_0 - \beta_x x_i) && \text{(constant factor)} \\
&= -2 \sum_{i=1}^n \frac{\partial}{\partial \beta_0} (x_i (y_i - \beta_0 - \beta_x x_i)) && \text{(derivative of a sum)} \\
&= -2 \sum_{i=1}^n x_i \frac{\partial}{\partial \beta_0} (y_i - \beta_0 - \beta_x x_i) && (x_i \text{ does not depend on } \beta_0) \\
&= -2 \sum_{i=1}^n x_i (-1) && \text{(inner derivative is } -1) \\
&= -2 \sum_{i=1}^n (-x_i) && \text{(multiply)} \\
&= -2 \left( - \sum_{i=1}^n x_i \right) && \text{(constant factor out of the sum)} \\
&= 2 \sum_{i=1}^n x_i && \text{(multiply)} \\
&= 2n\bar{x} && \left( \sum_{i=1}^n x_i = n\bar{x} \right)
\end{aligned}$$

Differentiating with respect to  $\beta_x$ , and reusing the inner derivative  $-x_i$  from Exercise 0.7:

$$\begin{aligned}
\frac{\partial^2 \text{RSS}}{\partial \beta_x^2} &= -2 \frac{\partial}{\partial \beta_x} \sum_{i=1}^n x_i (y_i - \beta_0 - \beta_x x_i) && \text{(constant factor)} \\
&= -2 \sum_{i=1}^n \frac{\partial}{\partial \beta_x} (x_i (y_i - \beta_0 - \beta_x x_i)) && \text{(derivative of a sum)} \\
&= -2 \sum_{i=1}^n x_i \frac{\partial}{\partial \beta_x} (y_i - \beta_0 - \beta_x x_i) && (x_i \text{ does not depend on } \beta_x) \\
&= -2 \sum_{i=1}^n x_i (-x_i) && \text{(inner derivative is } -x_i) \\
&= -2 \sum_{i=1}^n (-x_i^2) && \text{(multiply)} \\
&= -2 \left( - \sum_{i=1}^n x_i^2 \right) && \text{(constant factor out of the sum)} \\
&= 2 \sum_{i=1}^n x_i^2 && \text{(multiply)}
\end{aligned}$$

**The matrix.** Collecting the four second derivatives, the matrix of second derivatives is

$$\begin{pmatrix} 2n & 2n\bar{x} \\ 2n\bar{x} & 2 \sum_{i=1}^n x_i^2 \end{pmatrix}.$$

Its top-left entry,  $2n$ , is positive. Its determinant, using the expanded form of  $S_{xx}$  from Exercise 0.5, is

$$\begin{aligned}
\det \begin{pmatrix} 2n & 2n\bar{x} \\ 2n\bar{x} & 2 \sum_{i=1}^n x_i^2 \end{pmatrix} &= (2n) \left( 2 \sum_{i=1}^n x_i^2 \right) - (2n\bar{x})(2n\bar{x}) && (2 \times 2 \text{ determinant}) \\
&= 4n \sum_{i=1}^n x_i^2 - 4n^2 \bar{x}^2 && \text{(multiply)} \\
&= 4n \left( \sum_{i=1}^n x_i^2 - n\bar{x}^2 \right) && \text{(factor out } 4n) \\
&= 4n S_{xx} && \text{(expanded form of } S_{xx})
\end{aligned}$$

which is positive when  $S_{xx} > 0$ . A symmetric  $2 \times 2$  matrix with a positive top-left entry and a positive determinant is positive definite.

**Theorem 0.5** (Closed-form OLS estimates). *Suppose  $S_{xx} > 0$  (Definition 0.12), that is, not all  $x_i$  are equal. Then the OLS estimates (Example 0.5) are unique, and*

$$\hat{\beta}_x = \frac{S_{xy}}{S_{xx}}, \quad \hat{\beta}_0 = \bar{y} - \hat{\beta}_x \bar{x}.$$

### **i** Proof

*Proof.* RSS is differentiable, so any point that minimizes it solves the [normal equations](#). By [Exercise 0.8](#), the normal equations have exactly one solution,  $\beta_x = S_{xy}/S_{xx}$  and  $\beta_0 = \bar{y} - \beta_x \bar{x}$ . By [Exercise 0.9](#), the matrix of second derivatives of RSS is positive definite at every point, so RSS is strictly convex and its only stationary point is its unique global minimum. So the OLS estimates exist, are unique, and equal that solution.  $\square$

The same estimates follow from a derivation in vector notation, which treats  $(\beta_0, \beta_x)$  as a single vector instead of differentiating with respect to each component separately.

**Definition 0.13** (Covariate vector). In simple linear regression ([Definition 0.8](#)), the **covariate vector**  $\tilde{x}_i$  of observation  $i$  is its covariate value  $x_i$ , preceded by a 1 for the intercept:

$$\tilde{x}_i \stackrel{\text{def}}{=} \begin{pmatrix} 1 \\ x_i \end{pmatrix}.$$

**Exercise 0.10** (Mean as a dot product). Show that in simple linear regression ([Definition 0.8](#)), the conditional mean  $\mu(x_i)$  is the dot product of the covariate vector  $\tilde{x}_i$  ([Definition 0.13](#)) and the coefficient vector  $\tilde{\beta} = (\beta_0, \beta_x)^\top$  ([Example 0.6](#)):  $\mu(x_i) = \tilde{x}_i \cdot \tilde{\beta}$ .

### Solution

*Solution 0.10.* Start from the dot product:

$$\begin{aligned} \tilde{x}_i \cdot \tilde{\beta} &= \begin{pmatrix} 1 \\ x_i \end{pmatrix} \cdot \tilde{\beta} && \text{(definition of } \tilde{x}_i) \\ &= \begin{pmatrix} 1 \\ x_i \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_x \end{pmatrix} && \text{(definition of } \tilde{\beta}, \text{ written as a column)} \\ &= 1 \cdot \beta_0 + x_i \beta_x && \text{(definition of the dot product)} \\ &= \beta_0 + x_i \beta_x && (1 \cdot a = a) \\ &= \beta_0 + \beta_x x_i && \text{(reorder the factors)} \\ &= \mu(x_i) && \text{(definition of } \mu(x)) \end{aligned}$$

So  $\mu(x_i) = \tilde{x}_i \cdot \tilde{\beta}$ .

**Lemma 0.1** (The conditional mean is a dot product). *In simple linear regression (Definition 0.8), the conditional mean of observation  $i$  is the dot product of its covariate vector (Definition 0.13) and the coefficient vector  $\tilde{\beta}$  (Example 0.6):*

$$\mu(x_i) = \tilde{x}_i \cdot \tilde{\beta}.$$

### **i** Proof

*Proof.* This is the solution to Exercise 0.10. □

**Example 0.7** (Residual sum of squares in vector notation). In simple linear regression, take  $\tilde{\theta} = \tilde{\beta}$  and substitute  $\mu(x_i) = \tilde{x}_i \cdot \tilde{\beta}$  (Lemma 0.1) into each residual of the residual sum of squares (Definition 0.9). The residual sum of squares is then a function of the coefficient vector  $\tilde{\beta}$ , the same function as  $\text{RSS}(\beta_0, \beta_x)$  in Example 0.3:

$$\text{RSS}(\tilde{\beta}) = \sum_{i=1}^n (y_i - \tilde{x}_i \cdot \tilde{\beta})^2.$$

**Exercise 0.11** (Gradient of RSS in vector notation). Find the gradient of the residual sum of squares in vector notation (Example 0.7),  $\frac{\partial}{\partial \tilde{\beta}} \text{RSS}(\tilde{\beta})$ , by differentiating with respect to the vector  $\tilde{\beta}$  directly, rather than one component at a time.

### Solution

*Solution 0.11.* Differentiate one operation at a time, starting from  $\text{RSS}(\tilde{\beta})$  in vector notation (Example 0.7):

$$\begin{aligned} \frac{\partial}{\partial \tilde{\beta}} \text{RSS}(\tilde{\beta}) &= \frac{\partial}{\partial \tilde{\beta}} \sum_{i=1}^n (y_i - \tilde{x}_i \cdot \tilde{\beta})^2 && \text{(RSS in vector form)} \\ &= \sum_{i=1}^n \frac{\partial}{\partial \tilde{\beta}} (y_i - \tilde{x}_i \cdot \tilde{\beta})^2 && \text{(derivative of a sum)} \\ &= \sum_{i=1}^n 2(y_i - \tilde{x}_i \cdot \tilde{\beta}) \frac{\partial}{\partial \tilde{\beta}} (y_i - \tilde{x}_i \cdot \tilde{\beta}) && \text{(chain rule)} \end{aligned}$$

For the inner derivative, we need the gradient of a linear function. For any constant vector  $\tilde{a} = (a_0, a_x)^\top$ :

$$\begin{aligned}
\frac{\partial}{\partial \tilde{\beta}} \tilde{a} \cdot \tilde{\beta} &= \frac{\partial}{\partial \tilde{\beta}} (a_0 \beta_0 + a_x \beta_x) && \text{(write out the dot product)} \\
&= \left( \frac{\partial}{\partial \beta_0} (a_0 \beta_0 + a_x \beta_x), \frac{\partial}{\partial \beta_x} (a_0 \beta_0 + a_x \beta_x) \right) && \text{(definition of the gradient)} \\
&= \begin{pmatrix} a_0 + 0 \\ 0 + a_x \end{pmatrix} && \text{(derivative of each term)} \\
&= \begin{pmatrix} a_0 \\ a_x \end{pmatrix} && \text{(add zero)} \\
&= \tilde{a} && \text{(definition of } \tilde{a} \text{)}
\end{aligned}$$

With  $\tilde{a} = \tilde{x}_i$ :

$$\begin{aligned}
\frac{\partial}{\partial \tilde{\beta}} (y_i - \tilde{x}_i \cdot \tilde{\beta}) &= \frac{\partial}{\partial \tilde{\beta}} y_i - \frac{\partial}{\partial \tilde{\beta}} \tilde{x}_i \cdot \tilde{\beta} && \text{(derivative of a difference)} \\
&= \tilde{0} - \frac{\partial}{\partial \tilde{\beta}} \tilde{x}_i \cdot \tilde{\beta} && (y_i \text{ does not depend on } \tilde{\beta}) \\
&= \tilde{0} - \tilde{x}_i && \text{(gradient of a linear function)} \\
&= -\tilde{x}_i && \text{(subtract from zero)}
\end{aligned}$$

Plugging the inner derivative back in:

$$\begin{aligned}
\frac{\partial}{\partial \tilde{\beta}} \text{RSS}(\tilde{\beta}) &= \sum_{i=1}^n 2(y_i - \tilde{x}_i \cdot \tilde{\beta})(-\tilde{x}_i) && \text{(inner derivative is } -\tilde{x}_i \text{)} \\
&= \sum_{i=1}^n 2(-\tilde{x}_i)(y_i - \tilde{x}_i \cdot \tilde{\beta}) && \text{(reorder the scalar and vector factors)} \\
&= \sum_{i=1}^n (-2)\tilde{x}_i(y_i - \tilde{x}_i \cdot \tilde{\beta}) && (2(-\tilde{x}_i) = -2\tilde{x}_i) \\
&= -2 \sum_{i=1}^n \tilde{x}_i(y_i - \tilde{x}_i \cdot \tilde{\beta}) && \text{(constant factor out of the sum)}
\end{aligned}$$

**Exercise 0.12** (Normal equations in vector notation). Define the matrix and vector

$$A \stackrel{\text{def}}{=} \sum_{i=1}^n \tilde{x}_i \tilde{x}_i^\top, \quad \tilde{c} \stackrel{\text{def}}{=} \sum_{i=1}^n \tilde{x}_i y_i.$$

Using Exercise 0.11, show that the [normal equations](#) can be written as  $A\tilde{\beta} = \tilde{c}$ .

## Solution

*Solution* 0.12. Distribute the sum in the gradient from Exercise 0.11:

$$\begin{aligned}
 \frac{\partial}{\partial \tilde{\beta}} \text{RSS}(\tilde{\beta}) &= -2 \sum_{i=1}^n \tilde{x}_i (y_i - \tilde{x}_i \cdot \tilde{\beta}) && \text{(gradient of RSS)} \\
 &= -2 \sum_{i=1}^n (\tilde{x}_i y_i - \tilde{x}_i (\tilde{x}_i \cdot \tilde{\beta})) && \text{(distribute } \tilde{x}_i) \\
 &= -2 \left( \sum_{i=1}^n \tilde{x}_i y_i - \sum_{i=1}^n \tilde{x}_i (\tilde{x}_i \cdot \tilde{\beta}) \right) && \text{(sum of differences)}
 \end{aligned}$$

Each term of the second sum regroups as a matrix times  $\tilde{\beta}$ :

$$\begin{aligned}
 \tilde{x}_i (\tilde{x}_i \cdot \tilde{\beta}) &= \tilde{x}_i (\tilde{x}_i^\top \tilde{\beta}) && \text{(dot product as a transpose product)} \\
 &= (\tilde{x}_i \tilde{x}_i^\top) \tilde{\beta} && \text{(matrix multiplication is associative)}
 \end{aligned}$$

So the gradient is

$$\begin{aligned}
 \frac{\partial}{\partial \tilde{\beta}} \text{RSS}(\tilde{\beta}) &= -2 \left( \tilde{c} - \sum_{i=1}^n \tilde{x}_i (\tilde{x}_i \cdot \tilde{\beta}) \right) && \text{(definition of } \tilde{c}) \\
 &= -2 \left( \tilde{c} - \sum_{i=1}^n (\tilde{x}_i \tilde{x}_i^\top) \tilde{\beta} \right) && \text{(regroup each term)} \\
 &= -2 \left( \tilde{c} - \left( \sum_{i=1}^n \tilde{x}_i \tilde{x}_i^\top \right) \tilde{\beta} \right) && \text{(factor } \tilde{\beta} \text{ out of the sum)} \\
 &= -2 (\tilde{c} - A \tilde{\beta}) && \text{(definition of } A)
 \end{aligned}$$

The normal equations set this gradient equal to  $\tilde{0}$ . Solving for  $A \tilde{\beta}$  one operation at a time:

$$\begin{aligned}
 -2 (\tilde{c} - A \tilde{\beta}) &= \tilde{0} && \text{(normal equations)} \\
 \tilde{c} - A \tilde{\beta} &= \tilde{0} && \text{(divide both sides by } -2) \\
 \tilde{c} &= A \tilde{\beta} && \text{(add } A \tilde{\beta} \text{ to both sides)} \\
 A \tilde{\beta} &= \tilde{c} && \text{(swap the sides)}
 \end{aligned}$$

**Exercise 0.13** (Solving the normal equations in vector notation). Assuming  $S_{xx} > 0$  (Definition 0.12), show that  $A$  from Exercise 0.12 is invertible, solve  $A \tilde{\beta} = \tilde{c}$  for  $\tilde{\beta}$ , and compare the result with Exercise 0.8.

## Solution

*Solution* 0.13. Write out the entries of each term of  $A$  and  $\tilde{c}$ , then add them up:

$$\begin{aligned}
 A &= \sum_{i=1}^n \tilde{x}_i \tilde{x}_i^\top && \text{(definition of } A) \\
 &= \sum_{i=1}^n \begin{pmatrix} 1 & x_i \\ x_i & x_i^2 \end{pmatrix} && \text{(outer product of } \tilde{x}_i) \\
 &= \begin{pmatrix} \sum_{i=1}^n 1 & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{pmatrix} && \text{(matrices add entrywise)} \\
 &= \begin{pmatrix} n & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{pmatrix} && \text{(sum of a constant)} \\
 &= \begin{pmatrix} n & n\bar{x} \\ n\bar{x} & \sum_{i=1}^n x_i^2 \end{pmatrix} && \left(\sum_{i=1}^n x_i = n\bar{x}\right) \\
 \\
 \tilde{c} &= \sum_{i=1}^n \tilde{x}_i y_i && \text{(definition of } \tilde{c}) \\
 &= \sum_{i=1}^n \begin{pmatrix} y_i \\ x_i y_i \end{pmatrix} && \text{(scalar times a vector)} \\
 &= \begin{pmatrix} \sum_{i=1}^n y_i \\ \sum_{i=1}^n x_i y_i \end{pmatrix} && \text{(vectors add entrywise)} \\
 &= \begin{pmatrix} n\bar{y} \\ \sum_{i=1}^n x_i y_i \end{pmatrix} && \left(\sum_{i=1}^n y_i = n\bar{y}\right)
 \end{aligned}$$

The determinant of  $A$  is

$$\begin{aligned}
 \det A &= n \sum_{i=1}^n x_i^2 - (n\bar{x})(n\bar{x}) && \text{(determinant of a } 2 \times 2 \text{ matrix)} \\
 &= n \sum_{i=1}^n x_i^2 - n^2 \bar{x}^2 && \text{(multiply)} \\
 &= n \left( \sum_{i=1}^n x_i^2 - n\bar{x}^2 \right) && \text{(factor out } n) \\
 &= nS_{xx} && \text{(expansion of } S_{xx})
 \end{aligned}$$

where the last step uses Exercise 0.5. Since  $n > 0$  and  $S_{xx} > 0$ ,  $\det A > 0$ , so  $A$  is invertible, and  $A\tilde{\beta} = \tilde{c}$  has exactly one solution,  $A^{-1}\tilde{c}$ .

To simplify  $A^{-1}\tilde{c}$ , we need two expansions of uncentered sums. Adding  $n\bar{x}^2$  to both sides of Exercise 0.5 gives  $\sum_{i=1}^n x_i^2 = S_{xx} + n\bar{x}^2$ . The cross-product sum expands the same way:

$$\begin{aligned}
S_{xy} &= \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) && \text{(definition of } S_{xy}\text{)} \\
&= \sum_{i=1}^n (x_i y_i - x_i \bar{y} - \bar{x} y_i + \bar{x} \bar{y}) && \text{(expand the product)} \\
&= \sum_{i=1}^n x_i y_i - \sum_{i=1}^n x_i \bar{y} - \sum_{i=1}^n \bar{x} y_i + \sum_{i=1}^n \bar{x} \bar{y} && \text{(sum of a sum)} \\
&= \sum_{i=1}^n x_i y_i - \bar{y} \sum_{i=1}^n x_i - \bar{x} \sum_{i=1}^n y_i + \sum_{i=1}^n \bar{x} \bar{y} && \text{(constant factor out of the sum)} \\
&= \sum_{i=1}^n x_i y_i - \bar{y} \sum_{i=1}^n x_i - \bar{x} \sum_{i=1}^n y_i + \sum_{i=1}^n \bar{x} \bar{y} && \text{(constant factor out of the sum)} \\
&= \sum_{i=1}^n x_i y_i - \bar{y} \sum_{i=1}^n x_i - \bar{x} \sum_{i=1}^n y_i + n \bar{x} \bar{y} && \text{(sum of a constant)} \\
&= \sum_{i=1}^n x_i y_i - \bar{y} (n \bar{x}) - \bar{x} \sum_{i=1}^n y_i + n \bar{x} \bar{y} && \left( \sum_{i=1}^n x_i = n \bar{x} \right) \\
&= \sum_{i=1}^n x_i y_i - \bar{y} (n \bar{x}) - \bar{x} (n \bar{y}) + n \bar{x} \bar{y} && \left( \sum_{i=1}^n y_i = n \bar{y} \right) \\
&= \sum_{i=1}^n x_i y_i - n \bar{x} \bar{y} - n \bar{x} \bar{y} + n \bar{x} \bar{y} && \text{(reorder the factors)} \\
&= \sum_{i=1}^n x_i y_i - n \bar{x} \bar{y} && \text{(combine like terms)}
\end{aligned}$$

Adding  $n \bar{x} \bar{y}$  to both sides gives  $\sum_{i=1}^n x_i y_i = S_{xy} + n \bar{x} \bar{y}$ . Now compute the solution:

$$\begin{aligned}
A^{-1}\tilde{c} &= \frac{1}{\det A} \begin{pmatrix} \sum_{i=1}^n x_i^2 & -n\bar{x} \\ -n\bar{x} & n \end{pmatrix} \tilde{c} && \text{(inverse of a } 2 \times 2 \text{ matrix)} \\
&= \frac{1}{nS_{xx}} \begin{pmatrix} \sum_{i=1}^n x_i^2 & -n\bar{x} \\ -n\bar{x} & n \end{pmatrix} \tilde{c} && (\det A = nS_{xx}) \\
&= \frac{1}{nS_{xx}} \begin{pmatrix} \sum_{i=1}^n x_i^2 & -n\bar{x} \\ -n\bar{x} & n \end{pmatrix} \begin{pmatrix} n\bar{y} \\ \sum_{i=1}^n x_i y_i \end{pmatrix} && \text{(substitute } \tilde{c} \text{)} \\
&= \frac{1}{nS_{xx}} \begin{pmatrix} \sum_{i=1}^n x_i^2 \cdot n\bar{y} + (-n\bar{x}) \sum_{i=1}^n x_i y_i \\ (-n\bar{x}) \cdot n\bar{y} + n \sum_{i=1}^n x_i y_i \end{pmatrix} && \text{(matrix times a vector)} \\
&= \frac{1}{nS_{xx}} \begin{pmatrix} n\bar{y} \sum_{i=1}^n x_i^2 - n\bar{x} \sum_{i=1}^n x_i y_i \\ -n^2 \bar{x} \bar{y} + n \sum_{i=1}^n x_i y_i \end{pmatrix} && \text{(multiply out the products)} \\
&= \frac{1}{nS_{xx}} n \begin{pmatrix} \bar{y} \sum_{i=1}^n x_i^2 - \bar{x} \sum_{i=1}^n x_i y_i \\ -n\bar{x} \bar{y} + \sum_{i=1}^n x_i y_i \end{pmatrix} && \text{(factor } n \text{ out of the vector)} \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y} \sum_{i=1}^n x_i^2 - \bar{x} \sum_{i=1}^n x_i y_i \\ -n\bar{x} \bar{y} + \sum_{i=1}^n x_i y_i \end{pmatrix} && \text{(cancel } n \text{)} \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y} \sum_{i=1}^n x_i^2 - \bar{x} \sum_{i=1}^n x_i y_i \\ \sum_{i=1}^n x_i y_i - n\bar{x} \bar{y} \end{pmatrix} && \text{(reorder the terms of the second entry)} \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}(S_{xx} + n\bar{x}^2) - \bar{x} \sum_{i=1}^n x_i y_i \\ \sum_{i=1}^n x_i y_i - n\bar{x} \bar{y} \end{pmatrix} && \left( \sum_{i=1}^n x_i^2 = S_{xx} + n\bar{x}^2 \right) \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}(S_{xx} + n\bar{x}^2) - \bar{x}(S_{xy} + n\bar{x} \bar{y}) \\ (S_{xy} + n\bar{x} \bar{y}) - n\bar{x} \bar{y} \end{pmatrix} && \left( \sum_{i=1}^n x_i y_i = S_{xy} + n\bar{x} \bar{y} \right) \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}S_{xx} + \bar{y}n\bar{x}^2 - \bar{x}S_{xy} - \bar{x}n\bar{x} \bar{y} \\ S_{xy} + n\bar{x} \bar{y} - n\bar{x} \bar{y} \end{pmatrix} && \text{(distribute)} \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}S_{xx} + n\bar{x}^2 \bar{y} - \bar{x}S_{xy} - n\bar{x} \bar{x} \bar{y} \\ S_{xy} + n\bar{x} \bar{y} - n\bar{x} \bar{y} \end{pmatrix} && \text{(reorder the factors)} \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}S_{xx} + n\bar{x}^2 \bar{y} - \bar{x}S_{xy} - n\bar{x}^2 \bar{y} \\ S_{xy} + n\bar{x} \bar{y} - n\bar{x} \bar{y} \end{pmatrix} && (\bar{x}\bar{x} = \bar{x}^2) \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}S_{xx} - \bar{x}S_{xy} \\ S_{xy} + n\bar{x} \bar{y} - n\bar{x} \bar{y} \end{pmatrix} && \text{(first entry: cancel } \pm n\bar{x}^2 \bar{y} \text{)} \\
&= \frac{1}{S_{xx}} \begin{pmatrix} \bar{y}S_{xx} - \bar{x}S_{xy} \\ S_{xy} \end{pmatrix} && \text{(second entry: cancel } \pm n\bar{x} \bar{y} \text{)} \\
&= \begin{pmatrix} \bar{y}S_{xx}/S_{xx} - \bar{x}S_{xy}/S_{xx} \\ S_{xy}/S_{xx} \end{pmatrix} && \text{(multiply each entry by } 1/S_{xx} \text{)} \\
&= \begin{pmatrix} \bar{y} - \bar{x}S_{xy}/S_{xx} \\ S_{xy}/S_{xx} \end{pmatrix} && \text{(cancel } S_{xx} \text{)}
\end{aligned}$$

Its entries are  $\beta_0 = \bar{y} - \beta_x \bar{x}$  and  $\beta_x = S_{xy}/S_{xx}$ , the same solution as Exercise 0.8.

**Exercise 0.14** (Hessian of RSS in vector notation). Find the Hessian  $\frac{\partial}{\partial \tilde{\beta}} \frac{\partial}{\partial \tilde{\beta}^\top} \text{RSS}(\tilde{\beta})$ , and show that it is positive definite when  $S_{xx} > 0$  (Definition 0.12).

### Solution

*Solution* 0.14. Differentiate the gradient  $-2(\tilde{c} - A\tilde{\beta})$  from Exercise 0.12 with respect to  $\tilde{\beta}^\top$ .  $\tilde{c}$  does not depend on  $\tilde{\beta}$ . Each entry of  $A\tilde{\beta}$  is the dot product of a row of  $A$  with  $\tilde{\beta}$ , so by the gradient of a linear function (Exercise 0.11), the derivative of  $A\tilde{\beta}$  with respect to  $\tilde{\beta}^\top$  is  $A$ . So:

$$\begin{aligned}
 \frac{\partial}{\partial \tilde{\beta}} \frac{\partial}{\partial \tilde{\beta}^\top} \text{RSS}(\tilde{\beta}) &= \frac{\partial}{\partial \tilde{\beta}^\top} (-2(\tilde{c} - A\tilde{\beta})) && \text{(differentiate the gradient)} \\
 &= -2 \frac{\partial}{\partial \tilde{\beta}^\top} (\tilde{c} - A\tilde{\beta}) && \text{(constant factor)} \\
 &= -2 \left( \frac{\partial}{\partial \tilde{\beta}^\top} \tilde{c} - \frac{\partial}{\partial \tilde{\beta}^\top} A\tilde{\beta} \right) && \text{(derivative of a difference)} \\
 &= -2 \left( \mathbf{0} - \frac{\partial}{\partial \tilde{\beta}^\top} A\tilde{\beta} \right) && (\tilde{c} \text{ does not depend on } \tilde{\beta}) \\
 &= -2(\mathbf{0} - A) && \text{(derivative of a linear function)} \\
 &= -2(-A) && \text{(subtract from zero)} \\
 &= 2A && \text{(multiply)} \\
 &= 2 \sum_{i=1}^n \tilde{x}_i \tilde{x}_i^\top && \text{(definition of } A)
 \end{aligned}$$

To show it is positive definite, take any vector  $\tilde{v} = (v_0, v_x)^\top \neq \tilde{0}$ :

$$\begin{aligned}
\tilde{v}^\top (2A)\tilde{v} &= 2 \tilde{v}^\top A \tilde{v} && \text{(constant factor)} \\
&= 2 \tilde{v}^\top \left( \sum_{i=1}^n \tilde{x}_i \tilde{x}_i^\top \right) \tilde{v} && \text{(definition of } A) \\
&= 2 \sum_{i=1}^n \tilde{v}^\top (\tilde{x}_i \tilde{x}_i^\top) \tilde{v} && \text{(matrix multiplication distributes over the sum)} \\
&= 2 \sum_{i=1}^n (\tilde{v}^\top \tilde{x}_i)(\tilde{x}_i^\top \tilde{v}) && \text{(matrix multiplication is associative)} \\
&= 2 \sum_{i=1}^n (\tilde{v} \cdot \tilde{x}_i)(\tilde{x}_i \cdot \tilde{v}) && \text{(transpose products as dot products)} \\
&= 2 \sum_{i=1}^n (\tilde{v} \cdot \tilde{x}_i)^2 && \text{(dot product is symmetric)} \\
&= 2 \sum_{i=1}^n (v_0 + v_x x_i)^2 && \text{(write out the dot product)} \\
&\geq 0 && \text{(sum of squares)}
\end{aligned}$$

Equality would need  $v_0 + v_x x_i = 0$  for every  $i$ . If  $v_x \neq 0$ , that would force every  $x_i = -v_0/v_x$ , contradicting  $S_{xx} > 0$ ; if  $v_x = 0$ , it would force  $v_0 = 0$ , contradicting  $\tilde{v} \neq \tilde{0}$ . So the inequality is strict, and the Hessian is positive definite at every  $\tilde{\beta}$ .

**Theorem 0.6** (OLS estimates in vector notation). *Suppose  $S_{xx} > 0$  (Definition 0.12). Then the matrix  $A = \sum_{i=1}^n \tilde{x}_i \tilde{x}_i^\top$  is invertible, and the vector of OLS estimates (Definition 0.10) is*

$$\hat{\tilde{\beta}} = A^{-1} \tilde{c}, \quad \tilde{c} = \sum_{i=1}^n \tilde{x}_i y_i.$$

*Its entries are the estimates of Theorem 0.5.*

### **i** Proof

*Proof.* RSS is differentiable, so any point that minimizes it solves the [normal equations](#). By Exercise 0.12, the normal equations are  $A\tilde{\beta} = \tilde{c}$ . By Exercise 0.13,  $A$  is invertible, so their only solution is  $A^{-1}\tilde{c}$ , whose entries are those of Theorem 0.5. By Exercise 0.14, the Hessian of RSS is positive definite at every  $\tilde{\beta}$ , so RSS is strictly convex and  $A^{-1}\tilde{c}$  is its unique global minimum.  $\square$

**Exercise 0.15** (OLS slope and the correlation). Assuming  $S_{xx} > 0$  and  $S_{yy} > 0$  (Definition 0.12), write the OLS slope  $\hat{\beta}_x$  from Theorem 0.5 in terms of the [Pearson correlation coefficient](#)  $r$  and the [sample standard deviations](#)  $s_x$  and  $s_y$  of the  $x_i$  and the  $y_i$ .

#### Solution

*Solution 0.15.* In terms of the centered sums (Definition 0.12),

$$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}, \quad s_x = \sqrt{\frac{S_{xx}}{n-1}}, \quad s_y = \sqrt{\frac{S_{yy}}{n-1}}.$$

So:

$$\begin{aligned} r \frac{s_y}{s_x} &= \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} \cdot \frac{\sqrt{S_{yy}/(n-1)}}{\sqrt{S_{xx}/(n-1)}} && \text{(substitute } r, s_x, s_y) \\ &= \frac{S_{xy}}{\sqrt{S_{xx}}\sqrt{S_{yy}}} \cdot \frac{\sqrt{S_{yy}/(n-1)}}{\sqrt{S_{xx}/(n-1)}} && \text{(square root of a product)} \\ &= \frac{S_{xy}}{\sqrt{S_{xx}}\sqrt{S_{yy}}} \cdot \frac{\sqrt{S_{yy}}/\sqrt{n-1}}{\sqrt{S_{xx}}/\sqrt{n-1}} && \text{(square root of a quotient)} \\ &= \frac{S_{xy}}{\sqrt{S_{xx}}\sqrt{S_{yy}}} \cdot \frac{\sqrt{S_{yy}}}{\sqrt{S_{xx}}} && \text{(cancel the factors of } \sqrt{n-1}) \\ &= \frac{S_{xy}\sqrt{S_{yy}}}{\sqrt{S_{xx}}\sqrt{S_{yy}}\sqrt{S_{xx}}} && \text{(multiply the fractions)} \\ &= \frac{S_{xy}}{\sqrt{S_{xx}}\sqrt{S_{xx}}} && \text{(cancel } \sqrt{S_{yy}}, \text{ which is positive)} \\ &= \frac{S_{xy}}{S_{xx}} && (\sqrt{S_{xx}}\sqrt{S_{xx}} = S_{xx}) \\ &= \hat{\beta}_x && \text{(closed-form OLS slope)} \end{aligned}$$

**Corollary 0.2** (OLS slope in terms of the correlation). If  $S_{xx} > 0$  and  $S_{yy} > 0$  (Definition 0.12), then

$$\hat{\beta}_x = r \frac{s_y}{s_x},$$

where  $r$  is the [Pearson correlation coefficient](#) and  $s_x$  and  $s_y$  are the [sample standard](#)

deviations of the  $x_i$  and the  $y_i$ .

### Proof

*Proof.* This is the solution to Exercise [0.15](#). □

### Video lecture

Hutchinson's [Linear Regression \(pt1\)](#) (17 min) covers fitting a linear regression by least squares, framed as machine learning (Hutchinson, n.d.). The login for the video site is posted [on Canvas](#).

## Fitting a simple linear regression in R

**Example 0.8** (Regression of fasting glucose on BMI in HERS). The OLS estimates from Theorem [0.5](#), and the slope from Corollary [0.2](#), for the participants with a BMI measurement:

```
hers_bmi <- hers |> dplyr::filter(!is.na(BMI))
x <- hers_bmi$BMI
y <- hers_bmi$glucose
b1 <- sum((x - mean(x)) * (y - mean(y))) / sum((x - mean(x))^2)
c(
  intercept = mean(y) - b1 * mean(x),
  slope = b1,
  slope_from_r = cor(x, y) * sd(y) / sd(x)
)
#>      intercept      slope slope_from_r
#>      60.07405      1.82218      1.82218
```

`lm()` gives the same estimates:

```

slr_fit <- lm(glucose ~ BMI, data = hers)
summary(slr_fit)
#>
#> Call:
#> lm(formula = glucose ~ BMI, data = hers)
#>
#> Residuals:
#>      Min       1Q   Median       3Q      Max
#> -81.55 -18.98 -10.35   3.76 190.81
#>
#> Coefficients:
#>              Estimate Std. Error t value Pr(>|t|)
#> (Intercept)   60.074      3.565    16.9   <2e-16 ***
#> BMI           1.822      0.122    14.9   <2e-16 ***
#> ---
#> Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
#>
#> Residual standard error: 35.5 on 2756 degrees of freedom
#> (5 observations deleted due to missingness)
#> Multiple R-squared:  0.0743, Adjusted R-squared:  0.074
#> F-statistic: 221 on 1 and 2756 DF, p-value: <2e-16

```

Figure 4 shows the data with the fitted line.

```
ggplot2::ggplot(hers_bmi, ggplot2::aes(x = BMI, y = glucose)) +
  ggplot2::geom_point(alpha = 0.2) +
  ggplot2::geom_abline(
    intercept = coef(slr_fit)[[1]],
    slope = coef(slr_fit)[["BMI"]],
    color = "firebrick"
  ) +
  ggplot2::labs(x = "BMI (kg/m^2)", y = "Fasting glucose (mg/dL)")
```




Figure 4: Fasting glucose against BMI in HERS, with the OLS line

The estimated slope is  $\hat{\beta}_{\text{BMI}} = 1.82$  mg/dL per  $\text{kg}/\text{m}^2$ : mean fasting glucose is about 1.8 mg/dL higher among participants whose BMI is 1  $\text{kg}/\text{m}^2$  higher. The t statistic for the slope equals the correlation test statistic of Example 0.1.

### The coefficient of determination

**Definition 0.14** (Total sum of squares). The **total sum of squares** of  $y_1, \dots, y_n$  is

$$\text{TSS} \stackrel{\text{def}}{=} \sum_{i=1}^n (y_i - \bar{y})^2.$$

**Example 0.9** (Total sum of squares of three values). For  $y = 1, 2, 2$ ,  $\bar{y} = 5/3$ , so

$$\begin{aligned} \text{TSS} &= \left(1 - \frac{5}{3}\right)^2 + 2\left(2 - \frac{5}{3}\right)^2 && \text{(definition)} \\ &= \frac{4}{9} + \frac{2}{9} && \text{(square the deviations)} \\ &= \frac{2}{3} && \text{(arithmetic)} \end{aligned}$$

**Definition 0.15** (Coefficient of determination). The **coefficient of determination** of a model fitted with parameter estimate  $\hat{\theta}$  is one minus the ratio of its **residual sum of squares** at  $\hat{\theta}$  to the **total sum of squares** of its outcomes:

$$R^2 \stackrel{\text{def}}{=} 1 - \frac{\text{RSS}(\hat{\theta})}{\text{TSS}}.$$

*Remark 0.6* (Interpreting the coefficient of determination).  $R^2$  is often described as the proportion of the variation in  $Y$  explained by the regression on  $X$ .

**Exercise 0.16** (Residual sum of squares of the OLS fit). Assuming  $S_{xx} > 0$  (Definition 0.12), write the **residual sum of squares** at the OLS estimates (Theorem 0.5),  $\text{RSS}(\hat{\beta}_0, \hat{\beta}_x) = \sum_{i=1}^n r_i^2$ , in terms of  $S_{xx}$ ,  $S_{yy}$  and  $S_{xy}$ .

Solution

*Solution 0.16.* The **fitted values** of the OLS fit are  $\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_x x_i$ , so each **residual** is

$$\begin{aligned} r_i &= y_i - \hat{y}_i && \text{(definition of a residual)} \\ &= y_i - (\hat{\beta}_0 + \hat{\beta}_x x_i) && \text{(substitute } \hat{y}_i) \\ &= y_i - (\bar{y} - \hat{\beta}_x \bar{x} + \hat{\beta}_x x_i) && \text{(substitute } \hat{\beta}_0) \\ &= y_i - \bar{y} + \hat{\beta}_x \bar{x} - \hat{\beta}_x x_i && \text{(distribute the minus sign)} \\ &= (y_i - \bar{y}) - (\hat{\beta}_x x_i - \hat{\beta}_x \bar{x}) && \text{(group terms)} \\ &= (y_i - \bar{y}) - \hat{\beta}_x (x_i - \bar{x}) && \text{(factor out } \hat{\beta}_x) \end{aligned}$$

Squaring and summing:

$$\begin{aligned}
\sum_{i=1}^n r_i^2 &= \sum_{i=1}^n ((y_i - \bar{y}) - \hat{\beta}_x(x_i - \bar{x}))^2 && \text{(substitute } r_i\text{)} \\
&= \sum_{i=1}^n ((y_i - \bar{y})^2 - 2\hat{\beta}_x(x_i - \bar{x})(y_i - \bar{y}) + \hat{\beta}_x^2(x_i - \bar{x})^2) && \text{(expand the square)} \\
&= \sum_{i=1}^n (y_i - \bar{y})^2 - \sum_{i=1}^n 2\hat{\beta}_x(x_i - \bar{x})(y_i - \bar{y}) + \sum_{i=1}^n \hat{\beta}_x^2(x_i - \bar{x})^2 && \text{(sum of a sum)} \\
&= \sum_{i=1}^n (y_i - \bar{y})^2 - 2\hat{\beta}_x \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) + \sum_{i=1}^n \hat{\beta}_x^2(x_i - \bar{x})^2 && \text{(constant factor out of the sum)} \\
&= \sum_{i=1}^n (y_i - \bar{y})^2 - 2\hat{\beta}_x \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) + \hat{\beta}_x^2 \sum_{i=1}^n (x_i - \bar{x})^2 && \text{(constant factor out of the sum)} \\
&= S_{yy} - 2\hat{\beta}_x \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) + \hat{\beta}_x^2 \sum_{i=1}^n (x_i - \bar{x})^2 && \text{(definition of } S_{yy}\text{)} \\
&= S_{yy} - 2\hat{\beta}_x S_{xy} + \hat{\beta}_x^2 \sum_{i=1}^n (x_i - \bar{x})^2 && \text{(definition of } S_{xy}\text{)} \\
&= S_{yy} - 2\hat{\beta}_x S_{xy} + \hat{\beta}_x^2 S_{xx} && \text{(definition of } S_{xx}\text{)} \\
&= S_{yy} - 2 \frac{S_{xy}}{S_{xx}} S_{xy} + \left( \frac{S_{xy}}{S_{xx}} \right)^2 S_{xx} && \text{(substitute } \hat{\beta}_x = S_{xy}/S_{xx}\text{)} \\
&= S_{yy} - 2 \frac{S_{xy}^2}{S_{xx}} + \left( \frac{S_{xy}}{S_{xx}} \right)^2 S_{xx} && \text{(multiply)} \\
&= S_{yy} - 2 \frac{S_{xy}^2}{S_{xx}} + \frac{S_{xy}^2}{S_{xx}^2} S_{xx} && \text{(square of a quotient)} \\
&= S_{yy} - 2 \frac{S_{xy}^2}{S_{xx}} + \frac{S_{xy}^2}{S_{xx}} && \text{(cancel one factor of } S_{xx}\text{)} \\
&= S_{yy} - \frac{S_{xy}^2}{S_{xx}} && \text{(combine like terms)}
\end{aligned}$$

**Exercise 0.17** ( $R^2$  and the correlation). Assuming  $S_{xx} > 0$  and  $S_{yy} > 0$  (Definition 0.12), use Exercise 0.16 to write the **coefficient of determination**  $R^2$  of the OLS fit in terms of the **Pearson correlation coefficient**  $r$ , and show that  $0 \leq R^2 \leq 1$ .

#### Solution

*Solution 0.17.* The **total sum of squares** is  $S_{yy}$  (Definition 0.12), and Exercise 0.16 gives the residual sum of squares, so:

$$\begin{aligned}
R^2 &= 1 - \frac{\text{RSS}(\hat{\theta})}{\text{TSS}} && \text{(definition of } R^2\text{)} \\
&= 1 - \frac{\text{RSS}(\hat{\beta}_0, \hat{\beta}_x)}{\text{TSS}} && \text{(the OLS estimate is } (\hat{\beta}_0, \hat{\beta}_x)\text{)} \\
&= 1 - \frac{\text{RSS}(\hat{\beta}_0, \hat{\beta}_x)}{S_{yy}} && \text{(TSS is } S_{yy}\text{)} \\
&= 1 - \frac{S_{yy} - S_{xy}^2/S_{xx}}{S_{yy}} && \text{(residual sum of squares of the OLS fit)} \\
&= 1 - \left( \frac{S_{yy}}{S_{yy}} - \frac{S_{xy}^2/S_{xx}}{S_{yy}} \right) && \text{(split the fraction)} \\
&= 1 - \left( 1 - \frac{S_{xy}^2/S_{xx}}{S_{yy}} \right) && (S_{yy}/S_{yy} = 1) \\
&= 1 - 1 + \frac{S_{xy}^2/S_{xx}}{S_{yy}} && \text{(distribute the minus sign)} \\
&= \frac{S_{xy}^2/S_{xx}}{S_{yy}} && (1 - 1 = 0) \\
&= \frac{S_{xy}^2}{S_{xx}S_{yy}} && \text{(divide a fraction by } S_{yy}\text{)} \\
&= \left( \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} \right)^2 && \text{(write as a square)} \\
&= r^2 && (r = S_{xy}/\sqrt{S_{xx}S_{yy}})
\end{aligned}$$

Since  $-1 \leq r \leq 1$  ([range of the correlation coefficient](#)),  $0 \leq r^2 \leq 1$ .

**Theorem 0.7** ( $R^2$  of a simple linear regression). *For the OLS fit of a simple linear regression, with  $S_{xx} > 0$  and  $S_{yy} > 0$  (Definition 0.12),  $R^2$  is the square of the [Pearson correlation coefficient](#)  $r$  of the  $x_i$  and  $y_i$ , and so lies between 0 and 1:*

$$R^2 = r^2, \quad 0 \leq R^2 \leq 1.$$

### **i** Proof

*Proof.* This is the solution to Exercise 0.17, which uses the residual sum of squares from Exercise 0.16. □

**Example 0.10** ( $R^2$  for the regression of glucose on BMI in HERS). For the fit in Example 0.8,  $R^2$  computed from Definition 0.15, the square of the Pearson correlation, and `lm()`'s value agree:

```
fit <- lm(glucose ~ BMI, data = hers)
y <- fit$model$glucose
c(
  r_squared = 1 - sum(residuals(fit)^2) / sum((y - mean(y))^2),
  cor_squared = cor(fit$model$BMI, y)^2,
  lm = summary(fit)$r.squared
)
#>    r_squared cor_squared      lm
#>  0.0743455  0.0743455  0.0743455
```

BMI accounts for only about 7% of the variation in baseline fasting glucose.

## Further reading

[Linear Models Overview](#) covers linear regression in depth, including inference for the coefficients and multiple predictors. Vittinghoff et al. (2012) cover linear regression in Chapter 4.

## References

- Hogg, Robert V., Elliot A. Tanis, and Dale L. Zimmerman. 2019. *Probability and Statistical Inference*. Tenth edition. Pearson.
- Hutchinson, Brian. n.d. *DATA 471/571 (Machine Learning) and CSCI 481/581 (Deep Learning) Video Lectures*. Western Washington University. Accessed September 28, 2026. [https://facultyweb.cs.wwu.edu/~hutchib2/video\\_lectures/data371/](https://facultyweb.cs.wwu.edu/~hutchib2/video_lectures/data371/).
- Vittinghoff, Eric, David V Glidden, Stephen C Shiboski, and Charles E McCulloch. 2012. *Regression Methods in Biostatistics: Linear, Logistic, Survival, and Repeated Measures Models*. 2nd ed. Springer. <https://doi.org/10.1007/978-1-4614-1353-0>.