

# Copyright Notice

These slides are distributed under the Creative Commons License.

[DeepLearning.ai](#) makes these slides available for educational purposes. You may not use or distribute these slides for commercial purposes. You may make copies of these slides and use or distribute them for educational purposes as long as you cite [DeepLearning.AI](#) as the source of the slides.

For the rest of the details of the license, see <https://creativecommons.org/licenses/by-sa/2.0/legalcode>



DeepLearning.AI

# Math for Machine Learning

---

## Linear algebra - Week 4

# W4 Lesson 1




DeepLearning.AI

## Determinants and Eigenvectors



---

### Machine learning motivation


# Principal Component Analysis



# Principal Component Analysis




# Principal Component Analysis



# Principal Component Analysis

- Reduce dimensions (columns) of dataset
- Preserve as much information as possible



species	culmen_length_mm	culmen_depth_mm	flipper_length_mm	body_mass_g	PC1	PC2	species
Adelie	40.6	17.2	187.0	3475.0	1.353843	-0.422253	Adelie
Adelie	38.9	17.8	181.0	3625.0	1.760446	-0.350965	Adelie
Adelie	35.7	16.9	185.0	3150.0	2.005766	-1.113797	Adelie
Gentoo	50.0	15.3	220.0	5550.0	-2.585758	0.061768	Gentoo
Adelie	34.5	18.1	187.0	2900.0	2.438111	-0.786227	Adelie

# What to expect?



Linear transformation



Characterize your transformation

# What to expect?



Linear transformation




Singular / Non-singular



Characterize your transformation

Non-singular

🍎	banana
3	1
1	2



Singular

🍎	banana
1	1
2	2

# What to expect?



Linear transformation

1




2

3

Determinant as area



Characterize your transformation



# What to expect?



Linear transformation

1

2

3

Properties of determinant



Characterize your transformation

$$\begin{matrix} 3 & 1 \\ 1 & 2 \end{matrix}$$

Det = 5

$$\begin{matrix} 1 & 1 \\ -2 & 1 \end{matrix}$$

Det = 3

$$\begin{matrix} 1 & 4 \\ -3 & 3 \end{matrix}$$

Det = 15  
 $= 5 \cdot 3$

$$\begin{matrix} 3 & 1 \\ 1 & 2 \end{matrix}$$

Det = 5

$$\begin{matrix} 0.4 & -0.2 \\ -0.2 & 0.6 \end{matrix}$$

$\begin{matrix} 3 & 1 \\ 1 & 2 \end{matrix}^{-1} = \frac{1}{5} \begin{matrix} 0.4 & -0.2 \\ -0.2 & 0.6 \end{matrix}$

# What to expect?



Linear transformation



Singular / Non-singular



Determinant as area





Properties of determinant






Characterize your transformation





# What to expect?






# What to expect?



# What to expect?



# What to expect?



$$\begin{matrix} 2 & 1 & 1 \\ 0 & 3 & 0 \end{matrix} = \begin{matrix} 2 \\ 0 \end{matrix}$$

$(1,0) \rightarrow (2,0)$

$$A v_1 = \lambda_1 v_1$$

# What to expect?



Linear transformation

1

Singular / Non-singular

2

Determinant as area

3

Properties of determinant



Characterize your transformation



1

Basis

2


Span

3

Eigenvalues and  
Eigenvectors



PCA



2 dimensions



1 dimension






DeepLearning.AI

## Determinants and Eigenvectors


---

**Singularity and rank of linear transformations**

# Non-singular transformation




# Singular transformation




A diagram illustrating a singular transformation. It shows a 2x2 matrix multiplication:  $\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ . The result is a 2x1 vector where both elements are 2. Above the multiplication, there are two icons: an apple and a banana.

$(0,0) \rightarrow (0,0)$   
 $(1,0) \rightarrow (1,2)$   
 $(0,1) \rightarrow (1,2)$   
 $(1,1) \rightarrow (2,4)$




# Singular transformation



A diagram illustrating matrix multiplication. On the left, a 2x2 matrix with entries 0 and 0 is multiplied by a 2x1 column vector with entries  $a$  and  $b$ . The result is a 2x1 column vector with entries 0 and 0. This represents a singular transformation where the input space is collapsed into a single dimension.


$$\begin{matrix} 0 & 0 \\ 0 & 0 \end{matrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$



# Singular and non-singular transformations


Non-singular

3	1
1	2




Singular

1	1
2	2



Singular

0	0
0	0



# Rank of linear transformations

Rank 2

3	1
1	2



3	1
1	2

1	1
2	2

Rank 1

1	1
2	2



0	0
0	0




0	0
0	0

Rank 0

0	0
0	0



0	0
0	0







DeepLearning.AI

# Determinants and Eigenvectors

---


**Determinant as an area**

# Determinant as an area




$$\text{Det} = 3 \cdot 2 - 1 \cdot 1$$

$$\text{Det} = 5$$




# Determinant as an area





1	1
2	2

$$\text{Det} = 1 \cdot 2 - 1 \cdot 2$$

$$\text{Det} = 0$$




# Determinant as an area



$$\text{Det} = 0 \cdot 0 - 0 \cdot 0$$

$$\text{Det} = 0$$




# Determinant as an area

Non-singular

3	1
1	2

Determinant = 5




Area = 5

Singular

1	1
2	2

Determinant = 0




Area = 0

Singular

0	0
0	0

Determinant = 0



Area = 0

# Negative determinants?

	
3	1
1	2

$$\text{Det} = 3 \cdot 2 - 1 \cdot 1$$



$$\text{Det} = 5$$

	
1	3
2	1

$$\text{Det} = 1 \cdot 1 - 3 \cdot 2$$


$$\text{Det} = -5$$

# Determinant as an area





$$\text{Det} = 1 \cdot 1 - 3 \cdot 2$$

$$\text{Det} = -5$$




# Determinant as an area



$$\text{Det} = 1 \cdot 1 - 3 \cdot 2$$

$$\text{Det} = -5$$

Negative





DeepLearning.AI

## Determinants and Eigenvectors

---

### Determinant of a product

# Determinant of a product

3	1
1	2

5	2
1	2

=

16	8
7	6

$$\det = 5$$

$$3 \cdot 2 - 1 \cdot 1$$

$$\det = 8$$

$$5 \cdot 2 - 2 \cdot 1$$


$$\det = 40$$

$$16 \cdot 6 - 8 \cdot 7$$

# Determinant of a product


$$\det(AB) = \det(A) \det(B)$$

# Determinant of a product




$$\begin{array}{|c|c|} \hline 3 & 1 \\ \hline 1 & 2 \\ \hline \end{array}$$

Det = 5




Area blows up by 5

# Determinant of a product




$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline -2 & 1 \\ \hline \end{array}$$

Det = 3



Area blows up by 3

# Determinant of a product



# Quiz

- The product of a singular and a non-singular matrix (in any order) is:
  - Singular
  - Non-singular
  - Could be either one

# Solution

- If A is non-singular and B is singular, then  $\det(AB) = \det(A) \times \det(B) = 0$ , since  $\det(B) = 0$ . Therefore  $\det(AB) = 0$ , so AB is **singular**.


# When one factor is zero

$$5 \cdot 0 = 0$$

# When one factor is singular...


Non-singular	Singular	Singular
$\begin{matrix} 3 & 1 \\ 1 & 2 \end{matrix}$	$\begin{matrix} 1 & 2 \\ 1 & 2 \end{matrix}$	$\begin{matrix} 4 & 8 \\ 3 & 6 \end{matrix}$
$\text{Det} = 5$	$\text{Det} = 0$	$\text{Det} = 0$

# If one factor is singular...




3	1
1	2

Det = 5



1	2
1	2

Det = 0



Area blows up by 0



DeepLearning.AI

## Determinants and Eigenvectors

---

### Determinant of inverse

# Quiz

- Find the determinants of the following matrices

0.4	-0.2
-0.2	0.6

0.25	-0.25
-0.125	0.625

# Solution

$$\text{Det} \begin{array}{|c|c|} \hline 0.4 & -0.2 \\ \hline -0.2 & 0.6 \\ \hline \end{array} = (0.4)(0.6) - (-0.2)(-0.2) = 0.2$$

$$\text{Det} \begin{array}{|c|c|} \hline 0.25 & -0.25 \\ \hline -0.125 & 0.625 \\ \hline \end{array} = (0.25)(0.625) - (-0.125)(-0.25) = 0.125$$

# Determinant of an inverse

$$\begin{vmatrix} 3 & 1 \\ 1 & 2 \end{vmatrix}^{-1} = \begin{vmatrix} 0.4 & -0.2 \\ -0.2 & 0.6 \end{vmatrix}$$

$$\det = 5$$

$$5^{-1} = 0.2$$

$$\begin{vmatrix} 5 & 2 \\ 1 & 2 \end{vmatrix}^{-1} = \begin{vmatrix} 0.25 & -0.25 \\ -0.125 & 0.625 \end{vmatrix}$$

$$\det = 8$$

$$8^{-1} = 0.125$$

$$\begin{vmatrix} 1 & 2 \\ 1 & 2 \end{vmatrix} = \begin{vmatrix} ? & ? \\ ? & ? \end{vmatrix}$$

$$\det = 0$$

$$0^{-1} = ???$$

# Determinant of an inverse

$$\det(A^{-1}) = \frac{1}{\det(A)}$$

# Why?

$$\det(AB) = \det(A) \det(B)$$

Why is this?

$$\det(A^{-1}) = \frac{1}{\det(A)}$$

$$\begin{aligned} \det(AA^{-1}) &= \det(A) \det(A^{-1}) \\ \det(I) &= \det(A) \det(A^{-1}) \\ 1 &= \det(A) \det(A^{-1}) \\ 1 &= \frac{1}{\det(A)} \end{aligned}$$

# Determinant of the identity matrix

$$\det \begin{array}{|cc|} \hline 1 & 0 \\ 0 & 1 \\ \hline \end{array} = 1 \cdot 1 - 0 \cdot 0 = 1$$

$$\det(I) = 1$$

# W4 Lesson 2




DeepLearning.AI

# Determinants and Eigenvectors


---

## Bases


# Bases




3	1
1	2




# Bases




3	1
1	2





# Bases




Bases




# Bases




# Bases




**Bases**




# Bases




**Bases**



# What is not a basis?



Not bases






DeepLearning.AI

# Determinants and Eigenvectors


---

## Span


# Span




# Span




# Span




# Span



# Span





# Is this a basis?




No


# Is this a basis for something?



# A basis is a minimal spanning set




Basis




Not a basis

# A basis is a minimal spanning set




Basis




Not a basis

# Number of elements in the basis is the dimension




Dimensions: 1  
1 element in the basis




Dimensions: 2  
2 elements in the basis

# Linearly independent and linearly dependent vectors




Linearly independent

# Linearly independent and linearly dependent vectors




Linearly independent




Linearly independent

# Linearly independent and linearly dependent vectors




Linearly independent




Linearly independent


# Linearly independent and linearly dependent vectors



Linearly independent




Linearly independent




Linearly dependent

# Linearly independent and linearly dependent vectors




Linearly independent




Linearly independent


# Linearly independent and linearly dependent vectors



Linearly independent




Linearly independent




Linearly dependent


# Linearly independent and linearly dependent vectors



Linearly independent




Linearly independent




Linearly dependent


# Linearly independent and linearly dependent vectors



Linearly independent




Linearly independent




Linearly dependent


# Linearly independent and linearly dependent vectors



Linearly independent




Linearly independent




Linearly dependent

# Let's see how to check for linear dependence




$$\alpha + \beta =$$

# Let's see how to check for linear dependence



# Let's see how to check for linear dependence




$$\alpha v_1 + \beta v_2 = v_3$$
$$\alpha \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \beta \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$$

- 1
- 2

$$-\alpha + 2\beta = -5$$
$$\alpha + \beta = 3$$

# Let's see how to check for linear dependence



$$\alpha v_1 + \beta v_2 = v_3$$
$$\alpha \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \beta \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$$

$v_3$  is a linear combination  
of  $v_1$  and  $v_2$

$$\begin{array}{l} 1 \\ 2 \end{array} \quad \begin{aligned} -\alpha + 2\beta &= -5 \\ \alpha + \beta &= 3 \end{aligned}$$

$$\begin{array}{l} 1 \\ 2 \end{array} \quad \begin{array}{l} + \\ 3\beta = -2 \end{array} \rightarrow \beta = -\frac{2}{3}$$
$$\alpha - \frac{2}{3} = 3 \rightarrow \alpha = \frac{11}{3}$$

# Quiz

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

# Solution

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

1                  -1                  =                  Linearly dependent

# Solution

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} - 1 \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Linearly dependent

# Solution

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Not a basis!

Linearly independent

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

Linearly independent

$$\begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Linearly independent

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$


# Basis: a formal definition

A basis is a set of vectors that:


- Spans a vector space
- Is linearly independent




Not all sets of N vectors are a basis  
for N-dimensional space




Spans a line  
Linearly independent  
Is a basis



Spans the plane  
Linearly independent  
Is a basis



Spans a line  
Linearly dependent  
Not a basis



Spans the plane  
Linearly dependent  
Not a basis




DeepLearning.AI

# Determinants and Eigenvectors

---


## Eigenbasis

# Basis




$$\begin{matrix} 2 & 1 & 0 \\ 0 & 3 & 1 \end{matrix} = \begin{matrix} 1 \\ 3 \end{matrix}$$

$$\begin{aligned} (1,0) &\rightarrow (2,0) \\ (0,1) &\rightarrow (1,3) \end{aligned}$$




# Eigenbasis




$$\begin{matrix} 2 & 1 & 1 \\ 0 & 3 & 1 \end{matrix} = \begin{matrix} 3 \\ 3 \end{matrix}$$

$$\begin{array}{l} \times 2 \\ \hline (1,0) \rightarrow (2,0) \\ (1,1) \rightarrow (3,3) \end{array}$$




# Eigenbasis



$$\begin{matrix} 2 & 1 & 3 \\ 0 & 3 & 2 \end{matrix} = \begin{matrix} 8 \\ 6 \end{matrix}$$

$(3,2) \rightarrow (8,6)$






DeepLearning.AI

## Determinants and Eigenvectors

---

# Eigenvalues and Eigenvectors


# Eigenvalues and eigenvectors



$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 3 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & -1 \\ 0 & 3 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 6 \end{bmatrix} = \lambda \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$



# Eigenvalues and eigenvectors

$$\begin{matrix} 2 & 1 & 1 \\ 0 & 3 & 0 \end{matrix} = \begin{matrix} 2 & 1 \\ 0 & 0 \end{matrix}$$

8 multiplications

2 multiplications

Matrix Multiplication  
More work

$$A v_1$$

Scalar Multiplication  
Less work

$$\lambda_1 v_1$$

First pair

$$\begin{matrix} 2 & 1 & 1 \\ 0 & 3 & 1 \end{matrix} = \begin{matrix} 3 & 1 \\ 1 & 1 \end{matrix}$$

$$A v_2$$


$$\lambda_2 v_2$$

Second pair


Eigenvalues

Eigenvectors


# Eigenvalues and eigenvectors




# Eigenvalues and eigenvectors




# Eigenvalues and eigenvectors



# Eigenvalues and eigenvectors



# Eigenvalues and eigenvectors



# Eigenvalues and eigenvectors

- $Av = \lambda v$  for each eigenvector / eigenvalue
- Eigenvectors: direction of stretch
- Eigenvalues: how much stretch
- **Eigenbasis:** the set of a matrix's eigenvectors, can be arranged as a matrix with one eigenvector in each column
- Save work and characterize a transformation





DeepLearning.AI

## Determinants and Eigenvectors


---

**Calculating eigenvalues and eigenvectors**



# Finding eigenvalues




# Finding eigenvalues



# Finding eigenvalues



# Finding eigenvalues



# Finding eigenvalues

If  $\lambda$  is an eigenvalue:

$$\begin{array}{|c|c|c|} \hline 2 & 1 & x \\ \hline 0 & 3 & y \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline \lambda & 0 & x \\ \hline 0 & \lambda & y \\ \hline \end{array}$$

For infinitely many (x,y)

$$\begin{array}{|c|c|c|} \hline 2-\lambda & 1 & x \\ \hline 0 & 3-\lambda & y \\ \hline \end{array} = \begin{array}{|c|c|} \hline 0 \\ \hline 0 \\ \hline \end{array}$$

Has infinitely many solutions

$$\det \begin{array}{|c|c|} \hline 2-\lambda & 1 \\ \hline 0 & 3-\lambda \\ \hline \end{array} = 0$$

Characteristic polynomial

$$(2 - \lambda)(3 - \lambda) - 1 \cdot 0 = 0$$

$$\lambda = 2$$

$$\lambda = 3$$

# Finding eigenvectors

Eigenvalues:  $\lambda = 2$   
 $\lambda = 3$

Solve the equations

$$\begin{array}{|c|c|c|} \hline 2 & 1 & x \\ \hline 0 & 3 & y \\ \hline \end{array} = 2 \begin{array}{|c|c|} \hline x \\ \hline y \\ \hline \end{array}$$
$$2x + y = 2x$$
$$0x + 3y = 2y$$
$$x = 1$$
$$y = 0$$
$$\begin{array}{|c|} \hline 1 \\ \hline 0 \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline 2 & 1 & x \\ \hline 0 & 3 & y \\ \hline \end{array} = 3 \begin{array}{|c|c|} \hline x \\ \hline y \\ \hline \end{array}$$
$$2x + y = 3x$$
$$0x + 3y = 3y$$
$$x = 1$$
$$y = 1$$
$$\begin{array}{|c|} \hline 1 \\ \hline 1 \\ \hline \end{array}$$

# Quiz

- Find the eigenvalues and eigenvectors of this matrix:

9	4
4	3

# Solution

- Eigenvalues: 11, 1
- Eigenvectors: (2,1), (-1,2)

9	4
4	3

- The characteristic polynomial is

$$\det \begin{array}{|cc|} \hline 9-\lambda & 4 \\ 4 & 3-\lambda \\ \hline \end{array} = (9 - \lambda)(3 - \lambda) - 4 \cdot 4 = 0$$

- Which factors as  $\lambda^2 - 12\lambda + 11 = (\lambda - 11)(\lambda - 1)$

- The solutions are  $\lambda = 11$   
 $\lambda = 1$

# Finding eigenvalues

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{bmatrix}$$

$$\lambda I = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

**Characteristic polynomial:**  $\det(A - \lambda I) = 0$

$$\det \begin{bmatrix} 2 - \lambda & 1 & -1 \\ 1 & -\lambda & -3 \\ -1 & -3 & -\lambda \end{bmatrix} = 0$$

$$(2 - \lambda)\lambda^2 + 3 + 3 - 9(2 - \lambda) + \lambda + \lambda = -\lambda^3 + 2\lambda^2 + 11\lambda - 12 = 0$$

$$-(\lambda + 3)(\lambda - 1)(\lambda - 4) = 0$$

**Eigenvalues:**  $-3, 1, 4$

# Finding eigenvalues

$$A = \begin{matrix} & \begin{matrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{matrix} \end{matrix} \quad \text{Eigenvalues: } -3, 1, 4$$

$$Av = \lambda v$$

$$\underbrace{\begin{matrix} 2 & 1 & -1 & x_1 \\ 1 & 0 & -3 & x_2 \\ -1 & -3 & 0 & x_3 \end{matrix}}_{= 4} = 4 \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix}$$
$$\begin{matrix} 2x_1 + x_2 - x_3 \\ x_1 - 3x_3 \\ -x_1 - 3x_2 \end{matrix} = \begin{matrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{matrix}$$

# Finding eigenvalues

$$A = \begin{array}{|ccc|} \hline & 2 & 1 & -1 \\ & 1 & 0 & -3 \\ & -1 & -3 & 0 \\ \hline \end{array}$$

$$Av = \lambda v$$

$$\underbrace{\begin{array}{|ccc|c|} \hline & 2 & 1 & -1 & x_1 \\ & 1 & 0 & -3 & x_2 \\ & -1 & -3 & 0 & x_3 \\ \hline \end{array}}_{\begin{array}{l} 2x_1 + x_2 - x_3 \\ x_1 - 3x_3 \\ -x_1 - 3x_2 \end{array}} = 4 \begin{array}{|c|} \hline x_1 \\ x_2 \\ x_3 \\ \hline \end{array}$$

**Eigenvalues:**  $-3, 1, 4$

$$\begin{array}{rcl} 2x_1 + x_2 - x_3 & = & 4x_1 \\ x_1 - 3x_3 & = & 4x_2 \\ -x_1 - 3x_2 & = & 4x_3 \end{array}$$

$$\begin{array}{rcl} R_1 & -2x_1 + x_2 - x_3 & = 0 \\ R_2 & x_1 - 4x_2 - 3x_3 & = 0 \\ R_3 & -x_1 - 3x_2 - 4x_3 & = 0 \end{array}$$

$$\begin{array}{rcl} R_2 + R_3 & & 3R_1 + R_3 \\ -7x_2 - 7x_3 = 0 & & -7x_1 - 7x_3 = 0 \\ x_2 = -x_3 & & x_1 = -x_3 \end{array}$$

$$\begin{array}{l} x_1 = k \\ x_2 = k \\ x_3 = -k \end{array}$$

infinite solutions  
of this form

$$\begin{array}{ll} x_1 = 1 & x_1 = 2 \\ x_2 = 1 & x_2 = 2 \\ x_3 = -1 & x_3 = -2 \end{array}$$

this works! so does this!

**Eigenvector:**

$$\begin{array}{|c|} \hline 1 \\ 1 \\ -1 \\ \hline \end{array}$$

# Finding eigenvalues

$$A = \begin{matrix} \begin{array}{|c|c|c|} \hline 2 & 1 & -1 \\ \hline 1 & 0 & -3 \\ \hline -1 & -3 & 0 \\ \hline \end{array} \end{matrix}$$

**Eigenvalues**       $\lambda_1 = 4$        $\lambda_2 = 1$        $\lambda_3 = -3$

**Eigenvectors**

$\begin{matrix} 1 \\ 1 \\ -1 \end{matrix}$	$\begin{matrix} 0 \\ 1 \\ 1 \end{matrix}$	$\begin{matrix} 2 \\ -1 \\ 1 \end{matrix}$
--	---	--

# Note on dimensions

Eigenvalues → Determinant → Square Matrix

9	4
4	3



9	4	5
4	3	-2





DeepLearning.AI

# Determinants and Eigenvectors

---

**On the number of  
eigenvectors**

# Number of eigenvectors

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{bmatrix}$$

3 by 3 matrix

?

3 distinct eigenvalues

?

3 distinct eigenvectors

Eigenvalues

$$\lambda_1 = 4 \quad \lambda_2 = 1 \quad \lambda_3 = -3$$



Eigenvectors

$$\begin{array}{ccc} \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} & \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} & \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} \end{array}$$

# Repeated eigenvalues - Example 1

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{pmatrix}$$

Characteristic polynomial =  $\det(A - \lambda I) = \det$

$$\begin{pmatrix} 2 - \lambda & 0 & 0 \\ 1 & 4 - \lambda & 0.5 \\ 0 & 0 & 2 - \lambda \end{pmatrix}$$

$$(2 - \lambda)^2(4 - \lambda) + 0 + 0 - 0 - 0 - 0 = 0$$

Eigenvalues: 4, 2, 2 Repeated eigenvalue

# Repeated eigenvalues - Example 1

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 4$$

$$Av = 4v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 0 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{pmatrix}} = 4 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{pmatrix}$$

# Repeated eigenvalues - Example 1

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 4$$

$$Av = 4v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 0 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{matrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{matrix}} = 4 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{pmatrix}$$

$$\begin{aligned} 2x_1 &= 4x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 4x_2 \\ 2x_3 &= 4x_3 \end{aligned}$$

$$\begin{aligned} -2x_1 &= 0 \\ -x_1 - 0.5x_3 &= 0 \\ -2x_3 &= 0 \\ \rightarrow x_1 &= 0 \\ x_2 &= \text{any number} \\ \rightarrow x_3 &= 0 \end{aligned}$$

Eigenvector

$$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

# Repeated eigenvalues - Example 1

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 2$$

$$Av = 2v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 0 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{pmatrix}} = 2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{pmatrix}$$

# Repeated eigenvalues - Example 1

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 2$$

$$Av = 2v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 0 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{matrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{matrix}} = 2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{pmatrix}$$

$$\begin{aligned} 2x_1 &= 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 2x_2 \\ 2x_3 &= 2x_3 \end{aligned}$$

$$\begin{aligned} 0 &= 0 \\ -x_1 + 2x_2 - 0.5x_3 &= 0 \\ 0 &= 0 \end{aligned}$$

$$x_1 = 2x_2 - 0.5x_3$$

$$\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$$

$$\begin{aligned} x_1 &= 2 \\ x_2 &= 1 \\ x_3 &= 0 \end{aligned}$$

$$\begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$$

$$\begin{aligned} x_1 &= 1 \\ x_2 &= 1 \\ x_3 &= 2 \end{aligned}$$

Point in different directions  
Different eigenvectors

# Repeated eigenvalues - Example 1

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{pmatrix}$$

Eigenvalues       $\lambda_1 = 4$        $\lambda_2 = 2$        $\lambda_3 = 2$

Eigenvectors

$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$	$\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$	$\begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$
---	---	---

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix}$$

# Repeated eigenvalues - Example 2

$$A = \begin{matrix} \begin{array}{|ccc|} \hline & 2 & 0 \\ & -1 & 4 \\ & 4 & 0 \\ \hline & 0 & -0.5 \\ & 2 & \\ \hline \end{array} \end{matrix}$$

Characteristic polynomial =  $\det(A - \lambda I)$  =  $\det$

$$\begin{matrix} \begin{array}{|ccc|} \hline & 2-\lambda & 0 & 0 \\ & 1 & 4-\lambda & 0.5 \\ & -4 & 0 & 2-\lambda \\ \hline \end{array} \end{matrix}$$

$$(2 - \lambda)^2(4 - \lambda) + 0 + 0 - 0 - 0 - 0$$

Eigenvalues: 4, 2, 2    Repeated eigenvalue

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 4$$

$$Av = 4v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 4 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{pmatrix}} = 4 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{pmatrix}$$

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 4$$

$$Av = 4v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 4 & 0 & 2 & x_3 \end{pmatrix}}_{2x_1} = 4 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{pmatrix}$$

$$\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{pmatrix}$$

$$\begin{aligned} 2x_1 &= 4x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 4x_2 \\ 4x_1 + 2x_3 &= 4x_3 \end{aligned}$$

$$\begin{aligned} -2x_1 &= 0 \\ -x_1 - 0.5x_3 &= 0 \\ 4x_1 - 2x_3 &= 0 \end{aligned}$$

$$x_1 = 0 \quad x_3 = 0 \quad x_2 = \text{any number}$$

$$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad \begin{aligned} x_1 &= 0 \\ x_2 &= 1 \\ x_3 &= 0 \end{aligned}$$

Same as before!

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 2$$

$$Av = 2v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 4 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{pmatrix}} = 2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{pmatrix}$$

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 2$$

$$Av = 2v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 4 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{pmatrix}} = 2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{pmatrix}$$

$$\begin{aligned} 2x_1 &= 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 2x_2 \\ 4x_1 + 2x_3 &= 2x_3 \end{aligned}$$

$$\begin{aligned} 0 &= 0 & 0 \\ -x_1 + 2x_2 - 0.5x_3 &= 0 & k \\ 4x_1 &= 0 & 4k \end{aligned}$$

$$x_1 = 0 \qquad x_3 = 4x_2$$

$$\begin{array}{c|c} \begin{matrix} 0 \\ 1 \\ 4 \end{matrix} & \begin{matrix} x_1 = 0 \\ x_2 = 1 \\ x_3 = 4 \end{matrix} \\ \hline \begin{matrix} 0 \\ 0.5 \\ 2 \end{matrix} & \begin{matrix} x_1 = 0 \\ x_2 = 0.5 \\ x_3 = 2 \end{matrix} \end{array}$$

$$\begin{pmatrix} 0 \\ k \\ 4k \end{pmatrix}$$

On the same line  
Same eigenvector

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix} \quad \text{Eigenvalue: } 2$$

$$Av = 2v$$

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 & x_1 \\ -1 & 4 & -0.5 & x_2 \\ 4 & 0 & 2 & x_3 \end{pmatrix}}_{\begin{pmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{pmatrix}} = 2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{pmatrix}$$

$$\begin{aligned} 2x_1 &= 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 2x_2 \\ 4x_1 + 2x_3 &= 2x_3 \end{aligned}$$

$$\begin{aligned} 0 &= 0 \\ -x_1 + 2x_2 - 0.5x_3 &= 0 \\ 4x_1 &= 0 \end{aligned}$$

$$x_1 = 0 \qquad x_3 = 4x_2$$

$$\begin{pmatrix} 0 \\ k \\ 4k \end{pmatrix}$$

# Repeated eigenvalues - Example 2

$$A = \begin{pmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{pmatrix}$$

Eigenvalues

$$\lambda_1 = 4$$

$$\lambda_2 = 2$$

$$\lambda_3 = 2$$

Eigenvectors

$$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix}$$

Can't create an eigenbasis  
from this matrix

# Summary

a	b
c	d

Eigenvalues

$$\lambda_1, \lambda_2$$

a	b	c
d	e	f
g	h	i

$$\lambda_1, \lambda_2, \lambda_3$$

If  $\lambda_1 \neq \lambda_2$   2 eigenvectors  
(2 different directions)

If  $\lambda_1 = \lambda_2$    
1 eigenvector  
(1 direction)  
2 eigenvectors  
(2 different directions)

If  $\lambda_1 \neq \lambda_2 \neq \lambda_3$   3 eigenvectors  
(3 different directions)

If  $\lambda_1 = \lambda_2 \neq \lambda_3$    
2 eigenvectors  
(2 different directions)  
 3 eigenvectors  
(3 different directions)

If  $\lambda_1 = \lambda_2 = \lambda_3$    
1 eigenvector  
(1 direction)  
 2 eigenvectors  
(2 different directions)  
 3 eigenvectors  
(3 different directions)



DeepLearning.AI


## Determinants and Eigenvectors

---

**Dimensionality reduction  
and projection**


# Dimensionality Reduction

- Reduce dimensions (# of columns) of dataset
- Preserve as much information as possible



# Dimensionality Reduction

- Leads to smaller datasets
- Easier to visualize




Customer Age	Account Age	Days Since Login	Total Purchases	Total \$ Spent
23	1 month	10 days	1	\$100
71	45 months	2 days	5	\$150
54	30 months	15 days	2	\$70
36	22 months	12 days	4	\$210

Easy approach - just delete columns  
Loses valuable information


# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6



# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6



# Projections


x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

1
1

=

$$(1 + 1)$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

1
1

$$= (1 + 1)$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$(1+1)/\sqrt{2}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

Norm of 1

$$\frac{1}{\sqrt{2}} = \frac{1}{\left\| \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\|_2}$$

$$(1 + 1) / \sqrt{2}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ 1 \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6) \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2) \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2)/\sqrt{2} \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2)/\sqrt{2} \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

$$\begin{matrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2)/\sqrt{2} \\ (-1.3-0.6) \end{matrix}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

Final coordinates

$$\begin{aligned} & (1+1)/\sqrt{2} \\ & (1.2+1.6)/\sqrt{2} \\ & (-0.5+0.2)/\sqrt{2} \\ & (-1.3-0.6)/\sqrt{2} \end{aligned}$$




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix} 1 \\ 1 \end{matrix} \frac{1}{\sqrt{2}} =$$

Final coordinates

1.4142
1.9799
-0.2121
-1.344




# Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{2}} =$$

Final coordinates

1.4142
1.9799
-0.2121
-1.344



# Projections

To project a matrix  $A$  onto a vector  $v$

$$A_P = A \frac{v}{\|v\|_2}$$

$r \times 1$        $r \times c$        $c \times 1$

# Projections

To project a matrix  $A$  onto vectors  $v_1$  and  $v_2$

$$A_P = A \underbrace{\begin{bmatrix} \frac{v_1}{\|v_1\|_2} & \frac{v_2}{\|v_2\|_2} \end{bmatrix}}_{V}$$

$r \times 2$        $r \times c$        $c \times 2$

# Projections

To project a matrix  $A$  onto vectors  $v_1$  and  $v_2$

$$A_P = A \underbrace{\begin{bmatrix} \frac{v_1}{\|v_1\|_2} & \frac{v_2}{\|v_2\|_2} \end{bmatrix}}_{V}$$

$r \times 2$        $\boxed{r} \times c$        $c \times \boxed{2}$

# Projections

To project a matrix  $A$  onto vectors  $v_1$  and  $v_2$

$$A_P = \textcolor{brown}{A} \textcolor{green}{V}$$

$$r \times 2 \quad r \times c \quad c \times 2$$




DeepLearning.AI

## Determinants and Eigenvectors


---

## Motivating PCA


# Dimensionality Reduction




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)





# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)



# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)



# Benefits of Dimensionality Reduction

- Easier dataset to manage
- PCA reduces dimensions while minimizing information loss
- Simpler visualization





DeepLearning.AI


## Determinants and Eigenvectors

---


## Variance and covariance

# Mean


“The average of the data”




# Variance




# Variance



# Variance



# Variance



# Variance

$$Variance(x) = \frac{1}{n - 1} \sum_{i=1}^n (x_i - Mean(x))^2 = 16$$

	$x_i$	$x_i - Mean(x)$	$(x_i - Mean(x))^2$
1	10	1	1
2	4	-5	25
3	11	2	4
4	14	5	25
5	6	-3	9

→ 64

$$Mean(x) = 9$$

# Variance


$$Variance(x) = \frac{1}{n-1} \sum_{i=1}^n (x_i - \text{Mean}(x))^2$$

$$Var(x) \quad \mu_x$$

“The average squared distance from the mean”


# Variance

$$Var(y) = \frac{1}{n-1} \sum_{i=1}^n (y_i - \mu_Y)^2$$




$$Var(x) = \frac{1}{n-1} \sum_{i=1}^n (x_i - \mu_x)^2$$

# Problem



Negative covariance

Solution: Covariance




Positive covariance

# Covariance


“Take the average”

$$Cov(x, y) = \frac{1}{n - 1} \sum_{i=1}^n (x_i - \mu_x)(y_i - \mu_y)$$


$$Var(x) = \frac{1}{n - 1} \sum_{i=1}^n (x_i - \mu_x)^2$$



negative covariance



covariance zero  
(or very small)




positive covariance


# Covariance

$$Cov(x, y) = \frac{1}{n - 1} \sum_{i=1}^n (x_i - \mu_x)(y_i - \mu_y)$$


“The direction of the relationship between two variables”



negative covariance



covariance zero  
(or very small)



positive covariance




DeepLearning.AI

## Determinants and Eigenvectors


---

### The covariance matrix


# Covariance matrix



$$\begin{bmatrix} & \\ & \end{bmatrix}$$




$$\begin{bmatrix} & \\ & \end{bmatrix}$$




$$\begin{bmatrix} & \\ & \end{bmatrix}$$


# Covariance matrix



$$\begin{bmatrix} 3 & \\ & 1 \end{bmatrix}$$




$$\begin{bmatrix} 3 & \\ & 1 \end{bmatrix}$$




$$\begin{bmatrix} 3 & \\ & 1 \end{bmatrix}$$


# Covariance matrix



$$\begin{bmatrix} 3 & -2 \\ -2 & 1 \end{bmatrix}$$




$$\begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$$



$$\begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix}$$

# Covariance matrix



$$C = \begin{bmatrix} x & y \\ Cov(x, x) & Cov(x, y) \\ Cov(y, x) & Cov(y, y) \end{bmatrix}$$

$$Cov(x, x) = Var(x)$$

# Covariance matrix

$$\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix}$$

# Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \quad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$C = \frac{1}{n - 1} (\mathbf{x} - \boldsymbol{\mu})^T (\mathbf{x} - \boldsymbol{\mu})$$

# Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \quad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$C = \frac{1}{n - 1}(A - \mu)^T(A - \mu)$$

# Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \quad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$\begin{aligned} C &= \frac{1}{n-1}(A - \mu)^T(A - \mu) = \frac{1}{n-1} \left( \begin{array}{c|c} & - \\ \hline - & \end{array} \right)^T \left( \begin{array}{c|c} & - \\ \hline - & \end{array} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \end{aligned}$$

# Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \quad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &\quad \boxed{2} \times n \qquad \qquad \qquad n \times \boxed{2} \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C &= \frac{1}{n-1}(A - \mu)^T(A - \mu) = \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &\quad (x_1 - \mu_x)(x_1 - \mu_x) + (x_2 - \mu_x)(x_2 - \mu_x) + \dots + (x_n - \mu_x)(x_n - \mu_x) \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C &= \frac{1}{n-1}(A - \mu)^T(A - \mu) = \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &\quad \sum_{i=1}^n (x_i - \mu_x)^2 = \text{Var}(x) \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C &= \frac{1}{n-1}(A - \mu)^T(A - \mu) = \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & \\ & \ddots \\ & & Var(y) \end{bmatrix} \\ &\quad \frac{1}{n-1} \sum_{i=1}^n (x_i - \mu_x)^2 = Var(x) \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} = \begin{bmatrix} \textcolor{red}{Var(x)} \\ \vdots \end{bmatrix} \end{aligned}$$

$(x_1 - \mu_x)(y_1 - \mu_y) + (x_2 - \mu_x)(y_2 - \mu_y) + \dots + (x_n - \mu_x)(y_n - \mu_y)$

# Covariance matrix

$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} = \begin{bmatrix} \text{Var}(x) \\ \text{Var}(y) \end{bmatrix} \\ &\quad \sum_{i=1}^n (x_i - \mu_x)(y_i - \mu_y) = \text{Cov}(x, y) \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \sum_{i=1}^n (x_i - \mu_x)(y_i - \mu_y) = Cov(x, y) \end{aligned}$$

# Covariance matrix


$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x \\ x_2 - \mu_x \\ \vdots \\ x_n - \mu_x \end{bmatrix} \begin{bmatrix} y_1 - \mu_y \\ y_2 - \mu_y \\ \vdots \\ y_n - \mu_y \end{bmatrix} = \begin{bmatrix} \textcolor{red}{Var(x)} & \textcolor{green}{Cov(x,y)} \\ \textcolor{green}{Cov(y,x)} & \textcolor{blue}{Var(y)} \end{bmatrix} \end{aligned}$$

# Covariance matrix

$$\begin{aligned} C = \frac{1}{n-1}(A - \mu)^T(A - \mu) &= \frac{1}{n-1} \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right)^T \left( \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix} \right) \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \\ &= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & x_2 - \mu_x & \dots & x_n - \mu_n \\ y_1 - \mu_y & y_2 - \mu_y & \dots & y_n - \mu_y \end{bmatrix} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} = \begin{bmatrix} \textcolor{red}{Var(x)} & \textcolor{green}{Cov(x,y)} \\ \textcolor{green}{Cov(y,x)} & \textcolor{blue}{Var(y)} \end{bmatrix} \end{aligned}$$

# Matrix formula

$$A - \mu = \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \quad C = \frac{1}{n-1}(A - \mu)^T(A - \mu)$$



$A$

$x$	$y$
10	5
12	3
6	9
6	4
5	11
14	2
8	1
3	13

$A - \mu$

$x - \mu_x$	$y - \mu_y$
2	-1
4	-3
-2	3
-2	-2
-3	5
6	-4
0	-5
-5	8

$(A - \mu)^T$

$$\frac{1}{7}$$

$x - \mu_x$	2	4	-2	-2	-3	6	0	-5
$y - \mu_y$	-1	-3	3	-2	5	-4	-5	8

$A - \mu$

$x - \mu_x$	$y - \mu_y$
2	-1
4	-3
-2	3
-2	-2
-3	5
6	-4
0	-5
-5	8

$$\mu_x = 8 \quad \mu_y = 6$$

$$C = \begin{bmatrix} 14 & -11.86 \\ -11.86 & 19.71 \end{bmatrix}$$

# Matrix formula

$$A = \begin{bmatrix} x_1 & y_1 & z_1 \\ \vdots & \vdots & \vdots \\ x_n & y_n & z_n \end{bmatrix} \quad C = \frac{1}{n-1} (A - \mu)^T (A - \mu)$$

1. Arrange data with a different feature in each column
2. Calculate column averages
3. Subtract each average from their respective column to generate  $A - \mu$
4.  $\frac{1}{n-1} (A - \mu)^T (A - \mu)$  gives the covariance matrix  $C$




DeepLearning.AI

## Determinants and Eigenvectors


---

### PCA - Overview


# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)




# Principal Component Analysis (PCA)



# Principal Component Analysis (PCA)



# Principal Component Analysis (PCA)




$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

Eigenvalues (magnitude)

Eigenvectors (direction)

The matrix  $C$  is shown as a 2x2 matrix with eigenvalues 9 and 4 circled in cyan. To the right of the matrix are two pinkish-red arrows representing eigenvectors. The first arrow is horizontal and the second is diagonal. Below the matrix is a yellow trophy icon with the number 11 next to it.

# Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$


$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Eigenvectors  
(direction)

$$11$$

Eigenvalues  
(magnitude)

# Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$


$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Eigenvectors  
(direction)

$$11$$

Eigenvalues  
(magnitude)

# Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$


$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Eigenvectors  
(direction)


$$11$$

Eigenvalues  
(magnitude)


# Principal Component Analysis (PCA)





# Principal Component Analysis (PCA)




# PCA: Principal Component Analysis





# PCA: Principal Component Analysis




Eigenvalues/vectors




# PCA: Principal Component Analysis



Eigenvalues/vectors



# PCA: Principal Component Analysis






DeepLearning.AI

## Determinants and Eigenvectors


---

### PCA - Why it works


# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works





# PCA: Why It Works




# PCA: Why It Works




# PCA: Why It Works



# PCA: Why It Works



# PCA: Why It Works





DeepLearning.AI

## Determinants and Eigenvectors

---

**PCA - Mathematical  
formulation**

# PCA Mathematical formulation

You have  $n$  observations of 5 variables ( $x_1, x_2, x_3, x_4, x_5$ )

Goal: Reduce to 2 variables

1 Create matrix

$$X = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{15} \\ x_{21} & x_{22} & \dots & x_{25} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \dots & x_{n5} \end{bmatrix}$$

5 variables

*n Observations*

2 Center the data

$$X - \mu = \begin{bmatrix} x_{11} - \mu_1 & x_{12} - \mu_2 & \dots & x_{15} - \mu_5 \\ x_{21} - \mu_1 & x_{22} - \mu_2 & \dots & x_{25} - \mu_5 \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} - \mu_1 & x_{n2} - \mu_2 & \dots & x_{n5} - \mu_5 \end{bmatrix}$$

# PCA Mathematical formulation

You have  $n$  observations of 5 variables  $(x_1, x_2, x_3, x_4, x_5)$

Goal: Reduce to 2 variables

3

Calculate Covariance Matrix

$$C = \frac{1}{n-1}(X - \mu)^T(X - \mu) = \begin{bmatrix} Var(X_1) & Cov(X_1, X_2) & Cov(X_1, X_3) & Cov(X_1, X_4) & Cov(X_1, X_5) \\ Cov(X_1, X_2) & Var(X_2) & Cov(X_2, X_3) & Cov(X_2, X_4) & Cov(X_2, X_5) \\ Cov(X_1, X_3) & Cov(X_2, X_3) & Var(X_3) & Cov(X_3, X_4) & Cov(X_3, X_5) \\ Cov(X_1, X_4) & Cov(X_2, X_4) & Cov(X_3, X_4) & Var(X_4) & Cov(X_4, X_5) \\ Cov(X_1, X_5) & Cov(X_2, X_5) & Cov(X_3, X_5) & Cov(X_4, X_5) & Var(X_5) \end{bmatrix}$$

# PCA Mathematical formulation

You have  $n$  observations of 5 variables ( $x_1, x_2, x_3, x_4, x_5$ )

Goal: Reduce to 2 variables

4

Calculate Eigenvectors  
and Eigenvalues

5

Create Projection Matrix

6

Project Centered Data

Big	$\lambda_1$	$v_1$
	$\lambda_2$	$v_2$
	$\lambda_3$	$v_3$
	$\lambda_4$	$v_4$
Small	$\lambda_5$	$v_5$

$$V = \begin{bmatrix} & \\ \frac{1}{\|v_1\|_2} & \frac{1}{\|v_2\|_2} \\ & \end{bmatrix}$$

$$X_{PCA} = (X - \mu)V$$



DeepLearning.AI

## Determinants and Eigenvectors

---

## Conclusion