

Most common food-drug interaction

Drugs	Food	Interaction
WARFARIN	High-protein diet	raise serum albumin levels, decrease in international normalized ratio (INR).
	Vegetables containing vitamin k	decrease warfarin activity
	Cooked onions	increase warfarin activity
	Cranberry juice	elevated INR without bleeding in elderly patient.
	Leafy green vegetables	thromboembolic complications may develop.
MONOAMNINE OXIDASES	Tyramine-containing food	hypertensive crisis
PROPRANOLOL	Rich protein food	serum level may be increased
CELIPROLOL	Orange juice	the intestinal absorption is inhibited
ACES INHIBITORS	Empty stomach	absorption is increased
CA2 CHANNEL	Grape fruit juice	increases the bioavailability
ANTIBIOTICS	with milk products	reduced bioavailability
ACETAMINOPHEN	Pectin	delays its absorption and onset
NSAIDS	Garlic	Reduce blood clotting.
THEOPHYLINE	High-fat meal and grape fruit juice	increase bioavailability
	Caffeine	increases the risk of drug toxicity

ESOMEPRAZOLE	High-fat meal	reduce bioavailability
CIMETIDINE, RUPATADINE	with food (any type)	increase bioavailability
CYCLOSERINE	High-fat meal	decrease the serum concentration
ESOMEPRAZOLE	High-fat meal	reduce bioavailability
CIMETIDINE, RUPATADINE	with food (any type)	increase bioavailability
ACARBOSE	at start of each meal	maximum effectiveness
MERCAPTOPURINE	Cow's milk	reduce bioavailability
LEVOTHYROXINE	Grapefruit juice	delay the absorption
GLIMEPIRIDE	with breakfast	Increase bioavailability