

Medical Diets

NPO

- Aka “nothing by mouth”
- No food or water!
- Before some diagnostic tests and most surgeries

CLEAR LIQUID

- Common post-surgery
- Ok foods: gelatin, hard candy, broth, coffee and tea, popsicles

FULL LIQUID

- Post-surgery and when patient can tolerate more
- Ok foods: ice cream, milk, pudding, fruit and vegetable juice

SOFT LIQUID

- For those with difficulty chewing and swallowing
- Food needs to be mashed or chopped like mashed potatoes

THICKENED LIQUIDS

- Fluids are thickened with a thickening agent
- For difficulty swallowing, stroke/CVA, old age

FLUID-RESTRICTED

- Common in heart or renal failure patients
- **1000mL, 1500mL, or 2000mL max fluids** allowed per 24 hours

SODIUM-RESTRICTED

- For hypertension, heart failure, renal disease, heart disease

FAT-RESTRICTED

- For malabsorption diseases: pancreatitis, gallbladder disease, GERD
- Fat causes release of enzymes/bile that exacerbates disease

RENAL

- For acute glomerulonephritis, nephrotic syndrome, acute and chronic renal failure
- Low amounts of sodium, protein, potassium, or fluids

CARDIAC

- For hypertension, heart failure, CAD
- Low fat, cholesterol, sodium
- High in fruits, veggies and whole grains

LOW TYRAMINE

- Avoid high tyramine foods when taking MAOIs (psych meds) and some TB meds
- Tyramine foods: aged cheese, cured meat, coffee, chocolate, alcohol
- Can cause a hypertensive crisis if taken together

DIABETIC

- Well balanced meals. Ensure pt takes in carbs/protein prior to exercise
- Low sugar and carbs

LOW FIBER

- For inflammatory bowel diseases
- Okay: cooked vegetables, fruits, white breads, meats
- Avoid: beans, legumes, whole grains, raw vegetables, fruits or their juices, nuts, seeds

HIGH FIBER

- For constipation
- Increase fruits, beans, veggies, wheat, bran, fluids

GLUTEN FREE

- For celiac disease and gluten sensitivity
- No **BROW**: barley, rye, oats, wheat; instead, eat corn, rice or millet

LIVER/CIRRHOSIS

- Low in protein, sodium-restricted, fluid-restricted → to decrease the amount of ammonia produced from protein waste

LOW PURINE /URIC ACID

- For kidney stones, gout, uric acid retention
- Okay: spinach, asparagus, peas, fruit juice, cereal, egg, fat free milk, cottage cheese
- Avoid: protein/meat, some fish, beer/wine, sweetbreads

HIGH CALORIE /HIGH PROTEIN

- For debilitating diseases: cancer, burns, and COPD
- Pt's need the extra calories because their body burns off calories quickly