


# Long-term outcomes of eversion and conventional carotid endarterectomy: A multicenter clinical trial

Vascular  
2022, Vol. 0(0) 1–8  
© The Author(s) 2022  
Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: 10.1177/17085381221084803  
[journals.sagepub.com/home/vas](https://journals.sagepub.com/home/vas)  


Yuri V Belov<sup>1</sup>, Anton N Kazantsev<sup>2</sup>, Roman A Vinogradov<sup>3</sup> and Alexander V Korotkikh<sup>4</sup> 

## Abstract

**Aim:** To compare the long-term results of eversion (ECEA) and conventional carotid endarterectomy (CCEA).

**Methods:** We designed a retrospective, multicenter study which included 25,106 patients who underwent ECEA ( $n = 18,362$ ) or CCEA ( $n = 6744$ ). The duration of follow-up was  $124.7 \pm 53.8$  months.

**Results:** In the postoperative period, none of the interventions showed clear benefits reducing the frequency of complications: fatal outcome (ECEA: 0.19%,  $n = 36$ ; CCEA: 0.17%,  $n = 12$ ; OR = 1.1, 95% CI = 0.57–2.11,  $p = 0.89$ ), myocardial infarction (ECEA: 0.15%,  $n = 28$ ; CCEA: 0.13%,  $n = 9$ ;  $p = 0.87$ ; OR = 1.14; 95% CI = 0.53–2.42); acute cerebrovascular accident (CVA) (Group I: 0.33%,  $n = 62$ ; Group II: 0.4%,  $n = 27$ ;  $p = 0.53$ ; OR = 0.84; 95% CI = 0.53–1.32); bleeding with acute haematoma appearance in the area of intervention (Group I: 0.39%,  $n = 73$ ; Group II: 0.41%,  $n = 28$ ;  $p = 0.93$ ; OR = 0.95; 95% CI = 0.61–1.48); internal carotid artery (ICA) thrombosis (Group I: 0.05%,  $n = 11$ ; Group II: 0.07%,  $n = 5$ ; OR = 0.80, 95% CI = 0.28–2.32,  $p = 0.90$ ). During the long-term follow-up, ECEA was associated with lower frequency of fatal outcome (ECEA: 2.7%,  $n = 492$ ; CCEA: 9.1%,  $n = 616$ ; OR = 0.27; 95% CI = 0.24–0.3,  $p < 0.0001$ ), cerebrovascular death (ECEA: 1.0%,  $n = 180$ ; CCEA: 5.5%,  $n = 371$ ; OR = 0.17, 95% CI = 0.14–0.21,  $p < 0.0001$ ), non-fatal ischaemic stroke (ECEA: 0.62%,  $n = 114$ ; CCEA: 7.0%,  $n = 472$ ; OR = 0.08; 95% CI = 0.06–0.1,  $p < 0.0001$ ); repeated revascularization because of  $>60\%$  restenosis (ECEA: 1.6%,  $n = 296$ ; CCEA: 12.6%,  $n = 851$ ; OR = 0.11, 95% CI = 0.09–0.12,  $p < 0.0001$ ), and combined endpoint (ECEA: 2.2%,  $n = 397$ ; CCEA: 13.2%,  $n = 888$ ; OR = 0.14; 95% CI = 0.12–1.16,  $p < 0.0001$ ).

**Conclusion:** ECEA is beneficial over CCEA in a long term.

## Keywords

eversion carotid endarterectomy, conventional carotid endarterectomy, patch angioplasty, multicenter study, adverse outcomes, long-term follow-up

## Introduction

For  $>50$  years, both eversion (ECEA) and conventional carotid endarterectomy (CCEA) have been the ‘gold standard’ of cerebral revascularization, a surgical intervention aimed on treatment of internal carotid artery (ICA) stenosis [1–5]. ECEA is performed by cutting off the ICA from the bifurcation with the further endarterectomy after artery eversion [6,7]. In contrast, CCEA implies a longitudinal arteriotomy from the common carotid artery to the ICA [8,9]. In this surgical modality, plaque excision is followed by a patch angioplasty for closing the wound and preventing thrombosis [8–10]. These patches are fabricated from autologous saphenous veins, bovine or porcine pericardium, or poly (tetrafluoroethylene) [8,11–13] all having nearly equal

efficiency [12–15]. Currently, the surgeon is fully responsible for the choice of the intervention and the patch, while needing to take into account both clinical scenario and healthcare setting [16–18].

<sup>1</sup>Petrovsky National Research Centre of Surgery, Moscow, Russia

<sup>2</sup>Alexander Clinic, St. Petersburg, Russia

<sup>3</sup>Ochapovsky Regional Clinical Hospital #1, Krasnodar, Russia

<sup>4</sup>Cardiac Surgery Clinic, Amur State Medical Academy, Blagoveshchensk, Russia

## Corresponding author:

Anton N Kazantsev, MD, Alexander Hospital, Prospekt Solidarnosti, 4, St. Petersburg, Russia, 193231.

Email: [dr.antonio.kazantsev@mail.ru](mailto:dr.antonio.kazantsev@mail.ru)

Short- and long-term efficiency of ECEA and CCEA remains a matter of debate<sup>[19–22]</sup>. While several studies showed equal efficiency of these interventions, others demonstrated the inferior outcomes upon CCEA as patch angioplasty is associated with higher incidence of ICA restenosis and subsequent stroke<sup>[20–24]</sup>. However, current national guidelines authorise both of the indicated techniques, largely due to the lack of relevant multicenter studies in Russian Federation<sup>[18]</sup>. This leads to an unacceptably large number of preventable adverse outcomes, as currently 144 specialised clinics in Russia perform  $\approx 20,000$  CEAs annually<sup>[25]</sup>.

Here, we designed and performed a retrospective multicenter study to compare in-hospital and long-term outcomes of ECEA and CCEA. We found that albeit short-term efficiency of these interventions is equal, ECEA is associated with a significant reduction in adverse outcomes, including cerebrovascular death and repeated revascularization, in a long term.

## Materials and Methods

This retrospective multicenter study enrolled 25,106 patients who underwent ECEA ( $n = 18,362$ ) or CCEA ( $n = 6,744$ ) from 1 February 2006 to 1 September 2021. In CCEA, we used 3,187 (47.2%) xenopericardial patches, 2,354 (34.9%) poly (tetrafluoroethylene) patches and 1,203 (17.8%) patches fabricated from autologous veins. A 6/0 polypropylene suture was used to create anastomosis. To assess brain tolerance to ischaemia, we measured retrograde pressure and installed a temporary bypass if it was  $<60\%$  from the systemic blood pressure.

Criteria of inclusion were indications for CEA according to the current [Appendix I](#). Criteria of exclusion were:

- concomitant diseases limiting patient observation in the long term;
- contralateral neuropathy of glossopharyngeal and/or hypoglossal nerve;
- contralateral vocal cord paresis;
- severe neurological deficit;
- acute ischaemic stroke;
- acute coronary syndrome;
- bilateral severe ICA stenosis.

The severity of coronary atherosclerosis was calculated after the coronary angiography using the SYNTAX score. The duration of follow-up was  $124.7 \pm 53.8$  months. Patient data have been regularly collected by a phone survey and during the follow-up examination at a specialised healthcare facility. Among the adverse outcomes were

- fatal outcome;
- all-cause death;
- cerebrovascular death;
- cerebrovascular accident;
- non-fatal ischaemic stroke;
- ICA thrombosis;
- ICA restenosis  $\geq 60\%$  requiring repeated revascularization;
- myocardial infarction;
- bleeding of type  $\geq 3b$  (Bleeding Academic Research Consortium scale);
- combined endpoint (for the hospital observation: fatal outcome + myocardial infarction + cerebrovascular accident; for the long-term follow-up: cerebrovascular death + non-fatal ischaemic stroke + myocardial infarction).

A written informed consent was provided by all study participants after receiving a full explanation of the study. The investigation was carried out in accordance with the Good Clinical Practice and a latest revision of Declaration of Helsinki (2013).

Sample distribution was assessed using Kolmogorov–Smirnov test. For descriptive statistics, data were represented by the proportions or by the median, 25th and 75th percentiles, and range. Two independent groups were compared by Pearson’s chi-squared test with Yates’s correction for continuity or by Mann–Whitney  $U$  test. Calculation of odds ratio and respective 95% confidence intervals was conducted employing a respective MedCalc online calculator (MedCalc Software). Kaplan–Meier method was used to create survival curves during the long-term follow-up while log-rank test has been employed to compare survival distributions.  $p$  values  $\leq 0.05$  were regarded as statistically significant.

## Results

The samples were comparable regarding all clinicopathological features ([Table 1](#)). Elderly and male patients were predominant ([Table 1](#)). Around 20% of patients suffered from stable angina or had a past medical history of cerebrovascular accident ([Table 1](#)). Other comorbid conditions included diabetes mellitus, chronic obstructive pulmonary disease, and chronic kidney disease ([Table 1](#)). Severity of ICA stenosis and coronary atherosclerosis, as well as frequency of temporary bypass placement and duration of ICA clamping, did not differ significantly between the groups ([Table 2](#)).

Postoperatively, none of the interventions showed clear benefits reducing the frequency of complications: fatal outcome (ECEA: 0.19%,  $n = 36$ ; CCEA: 0.17%,  $n = 12$ ;

**Table 1.** Clinicopathological features of patient groups.

Features	Groups		p
	ECEA <i>n</i> = 18,362	CCEA <i>n</i> = 6,744	
Age, years (M ± m)	66.3 ± 6.1	65.1 ± 5.5	0.72
Male gender, <i>n</i> (%)	13646 (74.3)	4982 (73.9)	0.48
Cardiac fibrosis, <i>n</i> (%)	1953 (10.6)	720 (10.7)	0.94
Stable angina CCS class I-II	4067 (22.1)	1491 (22.1)	0.95
Diabetes mellitus, <i>n</i> (%)	1230 (6.7)	426 (6.3)	0.29
Chronic obstructive pulmonary disease, <i>n</i> (%)	512 (2.8)	193 (2.8)	0.78
Chronic kidney disease, <i>n</i> (%)	558 (3.0)	217 (3.2)	0.49
Peripheral atherosclerosis, <i>n</i> (%)	2461 (13.4)	854 (12.6)	0.13
Past medical history of myocardial revascularization, <i>n</i> (%)	2860 (15.6)	1059 (15.7)	0.82
Past medical history of stroke, <i>n</i> (%)	3752 (20.4)	1361 (20.2)	0.67

ECEA, eversion carotid endarterectomy; CCEA, conventional carotid endarterectomy.

**Table 2.** Angiographic and intraoperative features of patients.

	Groups		
Features	ECEA <i>n</i> = 18,362	CCEA <i>n</i> = 6,744	p
Angiographic features			
ICA stenosis, %	81.5 ± 5.3	83.3±6.0	0.7
SYNTAX score, M ± m	6.5 ± 2.1	5.9 ± 2.2	0.9
Intraoperative features			
Temporary bypass, <i>n</i> (%)	849 (4.6)	321 (4.7)	0.67
Duration of ICA clamping, minutes	24.5±3.8	25.6±4.5	0.2

ECEA, eversion carotid endarterectomy; CCEA, conventional carotid endarterectomy; ICA, internal carotid artery.

OR = 1.1, 95% CI = 0.57–2.11, *p* = 0.89), myocardial infarction (ECEA: 0.15%, *n* = 28; CCEA: 0.13%, *n* = 9; OR = 1.14, 95% CI = 0.53–2.42, *p* = 0.87); cerebrovascular accident (ECEA: 0.33%, *n* = 62; CCEA: 0.40%, *n* = 27; OR = 0.84, 95% CI = 0.53–1.32, *p* = 0.53); bleeding of type ≥ 3b (ECEA: 0.39%, *n* = 73; CCEA: 0.41%, *n* = 28; OR = 0.95, 95% CI = 0.61–1.48, *p* = 0.93); ICA thrombosis (ECEA: 0.05%, *n* = 11; CCEA: 0.07%, *n* = 5; OR = 0.80, 95% CI = 0.28–2.32, *p* = 0.90, Table 3).

ICA thrombosis (*n* = 15) was always accompanied by ischaemic stroke, being caused by (neo)intimal detachment behind the endarterectomy area (*n* = 10) or by arterial dissection upon temporary bypass placement (*n* = 5). These patients underwent urgent open thromboendarterectomy, ultimately leading to the neurological recovery in 11 out of 15 cases. Myocardial infarction (*n* = 37) was triggered by coronary artery thrombosis or in-stent restenosis and was successfully treated by percutaneous coronary intervention in all cases.

Among the cerebrovascular accidents (*n* = 89) were ischaemic stroke (*n* = 40), transient ischaemic attack (*n* = 29), haemorrhagic stroke (*n* = 12), and mixed stroke (*n* = 8). Fatal outcomes (*n* = 48) were caused by a cerebrovascular



accident (*n* = 43) or myocardial infarction (*n* = 5). The prevalence of a combined endpoint was ≈1%.

During the long-term follow-up, ECEA was associated with a lower frequency of fatal outcome (ECEA: 2.7%, *n* = 492; CCEA: 9.1%, *n* = 616; OR = 0.27; 95% CI = 0.24–0.3, *p* < 0.0001), cerebrovascular death (ECEA: 1.0%, *n* = 180; CCEA: 5.5%, *n* = 371; OR = 0.17, 95% CI = 0.14–0.21, *p* < 0.0001), non-fatal ischaemic stroke (ECEA: 0.62%, *n* = 114; CCEA: 7.0%, *n* = 472; OR = 0.08; 95% CI = 0.06–0.1, *p* < 0.0001); repeated revascularization because of >60% restenosis (ECEA: 1.6%, *n* = 296; CCEA: 12.6%, *n* = 851; OR = 0.11, 95% CI = 0.09–0.12, *p* < 0.0001), and combined endpoint (ECEA: 2.2%, *n* = 397; CCEA: 13.2%, *n* = 888; OR = 0.14; 95% CI = 0.12–1.16, *p* < 0.0001, Table 3). Cerebrovascular death – (Figure 1), cerebrovascular accident – (Figure 2), restenosis (≥60%, Figure 3, and combined endpoint-free survival (Figure 4) was also higher upon ECEA as compared with CCEA. Notably, ≥ 50% of all adverse outcomes after CCEA were recorded within 1 year of follow-up and were provoked by neointimal hyperplasia, probably triggered by a patch implantation.

**Table 3.** In-hospital and long-term outcomes.

Features	Groups		p	OR	95% CI
	ECEA n = 18,362	CCEA n = 6,744			
In-hospital outcomes					
Fatal outcome, n (%)	36 (0.19)	12 (0.17)	0.89	1.1	0.57–2.11
Myocardial infarction, n (%)	28 (0.15)	9 (0.13)	0.87	1.14	0.53–2.42
Cerebrovascular accident, n (%)	62 (0.33)	27 (0.4)	0.53	0.84	0.53–1.32
Bleeding of type ≥ 3b (BARC scale), n (%)	73 (0.39)	28 (0.41)	0.93	0.95	0.61–1.48
ICA thrombosis, n (%)	11 (0.05)	5 (0.07)	0.9	0.8	0.28–2.32
Combined endpoint, n (%)	126 (0.68)	48 (0.71)	0.89	0.96	0.69–1.34
Long-term outcomes					
All-cause death, n (%)	492 (2.7)	616 (9.1)	<0.0001	0.27	0.24–0.3
Cerebrovascular death, n (%)	180 (1.0)	371 (5.5)	<0.0001	0.17	0.14–0.21
Myocardial infarction, n (%)	103 (0.56)	45 (0.66)	0.37	0.84	0.59–1.19
Non-fatal ischaemic stroke, n (%)	114 (0.62)	472 (7.0)	<0.0001	0.08	0.06–0.1
Repeated revascularization because of ≥60% internal carotid artery restenosis, n (%)	296 (1.6)	851 (12.6)	<0.0001	0.11	0.09–0.12
Combined endpoint, n (%)	397 (2.2)	888 (13.2)	<0.0001	0.14	0.12–0.16


ECEA, eversion carotid endarterectomy; CCEA, conventional carotid endarterectomy; ICA, internal carotid artery; BARC, Bleeding Academic Research Consortium; OR, odds ratio; CI, confidence interval.

**Figure 1.** Cerebrovascular death-free survival.**Figure 2.** Cerebrovascular accident-free survival.


## Discussion

The reasons behind ICA restenosis upon CEA performance can be of local (e.g. patch-associated neointimal hyperplasia), systemic (e.g. arterial hypertension, severe dyslipidaemia, diabetes mellitus and smoking), and iatrogenic (e.g. technical errors) origin.

Excessive expansion of carotid bifurcation often results in the alterations of the flow pattern which becomes turbulent, entailing thrombosis and neointimal hyperplasia [26–30]. This, the inferiority of CCEA revealed in our trial concurs with those obtained in other studies<sup>[5,20,31]</sup>. A personalised computer simulation approach to patch selection to match arterial geometry remains the only option



**Figure 3.** Restenosis ( $\geq 60\%$ )-free survival.



**Figure 4.** Combined endpoint-free survival.

for the prevention of CCEA-associated complications which, however, remains rarely available<sup>[26–30]</sup>. Successful performance of virtual CCEA involving the implantation of a tailored patch provides a technical rationale for this treatment modality<sup>[26–30]</sup>. Albeit current guidelines do not specify the benefits of either ECEA or virtually assisted CCEA<sup>[18]</sup>, we suggest that cerebral revascularization should become increasingly personalised.

Systemic risk factors for the development of restenosis affect CEA outcomes regardless of intervention type<sup>[32–36]</sup>. Most of them are modifiable, yet it has been shown that

genetic background also affects the long-term outcomes<sup>[37–41]</sup>. Additional supervision of high-risk patients might permit early detection of ICA restenosis and timely repeated revascularization, preventing cerebrovascular accidents and death<sup>[37–41]</sup>.

Iatrogenic complications include narrowing of the arterial lumen as a result of the imposition of secondary sutures to prevent bleeding, crude patch-patient mismatch<sup>[42,43]</sup>, or residual stenosis because of incomplete plaque excision or intimal detachment after the blood flow restoration<sup>[44,45]</sup>. Unfortunately, approaches to predict a behaviour of (neo) intima behind the endarterectomy area upon restarting the blood flow are still lacking<sup>[42–45]</sup>.

To summarise, better long-term results of ECEA and its lower costs in comparison with CCEA prompt to recommend it as an only appropriate surgical option. Although ICA restenosis is a multifactorial condition and immense efforts are required to minimise the adverse outcomes, replacement of CCEA with ECEA would probably be a significant step on this route.

## Conclusion

ECEA is beneficial over CCEA in a long term.

## Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## References

1. Kazantsev AN, Tarasov RS, Burkov NN, et al. Carotid endarterectomy three-year follow-up in a single-center registry. *Angiol Vas Surg* 2018; 2417085381221084803(3): 101–108.
2. Vinogradov RA and Zakeryaev AB. Features of surgery of the internal carotid arteries with a combination of critical stenosis and pathological tortuosity. *Bulletin of the A.N. Baklava RAMS. Cardiovascular diseases* 2018; 1917085381221084803(3): 305–312. DOI: [10.24022/1810-0694-2018-19-3-305-312](https://doi.org/10.24022/1810-0694-2018-19-3-305-312)
3. Vinokurov IA, Yasnopolskaya NV, Hasanov EN, et al. Carotid endarterectomy in the early period after systemic thrombolysis in acute cerebrovascular accident. *Cardiol Cardiovascular Surg* 2021; 1417085381221084803(5): 370–375. DOI: [10.17116/kardio202114051370](https://doi.org/10.17116/kardio202114051370)
4. Zavaruev AV. Prevention of ischemic brain injury in surgery of the brachiocephalic arteries. *Cardiol Cardiovascular Surg* 2022; 1417085381221084803(5): 376–379. DOI: [10.17116/kardio202114051376](https://doi.org/10.17116/kardio202114051376)



5. Kazantsev AN, Chernykh KP, Leader RY, et al. Glomus-sparing carotid endarterectomy according to A.N. Kazantsev. Hospital and mid-term results. *Circulatory Pathol Cardiac Surg* 2020; 2417085381221084803(3): 70–79. DOI: [10.21688/1681-3472-2020-3-70-79](https://doi.org/10.21688/1681-3472-2020-3-70-79)
6. Kazantsev AN, Chernykh KP, Zarkua NE, et al. Eversion carotid endarterectomy transposition of the internal carotid artery over the hypoglossal nerve. *Surgery* 2021; 617085381221084803: 63–71. DOI: [10.17116/hirurgia202106163](https://doi.org/10.17116/hirurgia202106163)
7. Kazantsev AN, Vinogradov RA, Erofeev AA, et al. Extended lesion of the internal carotid artery: Six types of reconstruction. Results of a multicenter study. *Cardiol Cardiovascular Surg* 2021; 1417085381221084803(5): 354–369. DOI: [10.17116/kardio202114051354](https://doi.org/10.17116/kardio202114051354)
8. Kazantsev AN, Chernyavsky MA, Vinogradov RA, et al. Implantation of a long biological patch during classical carotid endarterectomy for extended atherosclerotic lesions. Long-term results. *Bulletin of Transplantol Artificial Organs* 2021; 2317085381221084803(1): 112–124. DOI: [10.15825/1995-1191-2021-1-112-124](https://doi.org/10.15825/1995-1191-2021-1-112-124)
9. Vachev AN, Prozhoga MG and Dmitriev OV. Influence of systemic hemodynamics on retrograde pressure in the internal carotid artery during carotid endarterectomy. *Clin Physiol Blood Circulation* 2019; 1617085381221084803(4): 293–298. DOI: [10.24022/1814-6910-2019-16-4-293-298](https://doi.org/10.24022/1814-6910-2019-16-4-293-298)
10. Kazantsev AN, Chernykh KP, Zarkua NE, et al. A new method of glomus-sparing carotid endarterectomy according to A. N. Kazantsev: Cutting off the internal carotid artery at the site from the external and common carotid artery. *Russian J Cardiol* 2020; 2517085381221084803(8): 10–17. DOI: [10.15829/1560-4071-2020-3851](https://doi.org/10.15829/1560-4071-2020-3851)
11. Belov YV, Lysenko AV, Lednev PV, et al. Application of decellularized xenopericardium patch in brachiocephalic artery surgery. *Cardiol Cardiovascular Surg* 2018; 1117085381221084803(2): 31–34. DOI: [10.17116/kardio201811231-34](https://doi.org/10.17116/kardio201811231-34)
12. Bockeria LA, Abdulgasanov RAO, Shogenov MA, et al. Infection of synthetic patches and prostheses of brachiocephalic arteries: Diagnostic and treatment methods. *Breast and Cardiovascular Surg* 2020; 6217085381221084803(6): 494–505. DOI: [10.24022/0236-2791-2020-626-494-505](https://doi.org/10.24022/0236-2791-2020-626-494-505)
13. Darvish NAMA, Shogenov MA, Abdulgasanov RAO, et al. Surgical treatment methods for atherosclerotic lesions of the carotid arteries. *Breast and Cardiovascular Surg* 2020; 6217085381221084803(5): 405–416. DOI: [10.24022/0236-2791-2020-62-5-405-416](https://doi.org/10.24022/0236-2791-2020-62-5-405-416)
14. Yarikov AV, Fraerman AP, Mukhin AS, et al. The history of the development of carotid artery surgery. *Neurosurg Neurol Kazakhstan* 2019; 5617085381221084803(3): 78–90.
15. Fokin AA and Kuvatov AV. Immediate and long-term results of carotid endarterectomy with dilatation patch repair. Proceedings of higher educational institutions. *Ural region* 2013; 217085381221084803: 161–166.
16. Kazantsev AN, Sultanov RV, Burkov NN, et al. Long-term results of surgical and conservative treatment of patients with occlusive-stenotic lesions of the carotid arteries. *Surgery* 2020; 117085381221084803: 67–73. DOI: [10.17116/hirurgia202001167](https://doi.org/10.17116/hirurgia202001167)
17. Kazantsev AN, Tarasov RS, Burkov NN, et al. Hospital results of percutaneous coronary intervention and carotid endarterectomy in hybrid and phased modes. *Angiol Vascular Surg* 2019; 2517085381221084803(1): 101–107. DOI: [10.33529/angio2019114](https://doi.org/10.33529/angio2019114)
18. National guidelines for the management of patients with brachiocephalic artery disease. *Angiol Vascular Surg*. 2013; 1917085381221084803(2): 4–68
19. Dudanov IP, Abuazab BS and Akhmetov VV. Comparative characteristics of classical and eversion endarterectomy of the carotid arteries. *Res Pract Med* 2017; 417085381221084803(3): 8–15. DOI: [10.17709/2409-2231-2017-4-3-1](https://doi.org/10.17709/2409-2231-2017-4-3-1)
20. Kazantsev AN, Burkov NN, Anufriev AI, et al. Medium-term results of carotid endarterectomy in patients with varying degrees of contralateral lesion. *Cardiol Cardiovascular Surg* 2020; 1317085381221084803(2): 95–103. DOI: [10.17116/kardio20201302195](https://doi.org/10.17116/kardio20201302195)
21. Gavrilenko AV, Kuklin AV, Al-Yousef NN, et al. Meta-analysis of the results of eversion carotid endarterectomy and endarterectomy with plastic patch. *Angiol Vascular Surg* 2020; 2617085381221084803(1): 176–183. DOI: [10.33529/ANGIO202012](https://doi.org/10.33529/ANGIO202012)
22. Kazantsev AN, Chernykh KP, Leader RY, et al. Comparative results of classical and eversion carotid endarterectomy. *Cardiol Cardiovascular Surg* 2020; 1317085381221084803(6): 550–555. DOI: [10.17116/kardio202013061550](https://doi.org/10.17116/kardio202013061550)
23. Alekryan BG, Pokrovsky AV, Zotikov AE, et al. Results of various treatment strategies for patients with combined lesions of the internal carotid and coronary arteries. *Endovascular Surg* 2021; 817085381221084803(2): 144–153. DOI: [10.24183/2409-4080-2021-8-2-144-153](https://doi.org/10.24183/2409-4080-2021-8-2-144-153)
24. Pokrovsky AV, Beloyartsev DF, Adyrkhaev ZA, et al. Does the method of carotid reconstruction affect the immediate results of the intervention? *Angiol Vascular Surg* 2012; 1817085381221084803(3): 81–91
25. Pokrovsky AV and Golovyuk AL. The state of vascular surgery in the Russian Federation in 2018. *Angiol Vascular Surg App* 2018; 2517085381221084803(2): 1–40.
26. Kazantsev AN, Vinogradov RA, Zakharov YN, et al. Prediction of restenosis after carotid endarterectomy by computer simulation. *Emerg Medical Care J Them. N.V. Sklifosovsky* 2021; 1017085381221084803(2): 401–407. DOI: [10.23934/2223-9022-2021-10-2-401-407](https://doi.org/10.23934/2223-9022-2021-10-2-401-407)
27. Kazantsev AN, Burkov NN, Zakharov YN, et al. Personalized brain revascularization: A method of computer modeling of the reconstruction area for carotid endarterectomy. *Surgery* 2020; 617085381221084803: 71–75. DOI: [10.17116/hirurgia202006171](https://doi.org/10.17116/hirurgia202006171)

28. Kazantsev AN, Burkov NN, Borisov VG, et al. Computer modeling of hemodynamic parameters in the bifurcation of the carotid arteries after carotid endarterectomy. *Angiol Vascular Surg* 2019; 2517085381221084803(3): 107–112. DOI: [10.33529/ANGIO2019311](https://doi.org/10.33529/ANGIO2019311)
29. Kazantsev AN, Bogomolova AV, Burkov NN, et al. Morphology of restenosis after classical carotid endarterectomy with diepoxy-treated xenopericardium patch. *Cardiol Cardiovascular Surg* 2020;1317085381221084803(1):68–71. DOI: [10.17116/kardio202013011168](https://doi.org/10.17116/kardio202013011168)
30. Kazantsev AN, Chernykh KP, Zarkua NE, et al. “Chik-chirik” carotid endarterectomy. Bulletin NTSSSH them. A.N. Bakuleva RAMS. *Cardiovascular Diseases* 2020; 2117085381221084803(4): 414–428. DOI: [10.24022/1810-0694-2020-21-4-414-428](https://doi.org/10.24022/1810-0694-2020-21-4-414-428)
31. Kazantsev AN, Tarasov RS, Burkov NN, et al. Predictors of long-term complications after carotid endarterectomy. *Surgery* 2019; 617085381221084803: 20–25. DOI: [10.17116/hirurgia201906120](https://doi.org/10.17116/hirurgia201906120)
32. Bazylev VV, Shmatkov MG and Morozov ZA. Predictors of complications in the early postoperative period after carotid artery stenting and carotid endarterectomy. *Diagnostic and Interventional Radiol* 2019; 1317085381221084803(2): 44–50. DOI: [10.25512/DIR.2019.13.2.05](https://doi.org/10.25512/DIR.2019.13.2.05)
33. Anufriev PL, Tanashyan MM, Gulevskaya TS, et al. Morphological markers of the main pathogenetic variants of ischemic strokes in cerebral atherosclerosis. *An Clin Exp Neurol* 2018; 1217085381221084803(4): 16–22
34. Vinogradov RA, Kosenkov AN, Vinokurov IA, et al. “Dumb” ischemic foci in the brain after revascularization of the brachiocephalic arteries. *Bulletin of the National Medical and Surgical Center. N.I. Pirogov* 2017; 1217085381221084803(2): 52–54
35. Kazantsev AN. Multicenter study: Results of classical and eversional carotid endarterectomy with and without type 2 diabetes mellitus. *Russian J Cardiol* 2021; 2617085381221084803(S7): 7
36. Zybin AV, Timina IE, Pokrovsky AV, et al. Immediate results of carotid endarterectomy in patients receiving statin therapy. *Russian Med Biol Bulletin Named After Acad I.P. Pavlova* 2017; 2517085381221084803(3): 1189–1414. DOI: [10.23888/PAVLOVJ20173404-414](https://doi.org/10.23888/PAVLOVJ20173404-414)
37. Nazarenko MS, Sleptcov AA, Lebedev IN, et al. Genomic structural variations for cardiovascular and metabolic comorbidity. *Sci Rep* 2017; 717085381221084803: 41268. DOI: [10.1038/srep41268](https://doi.org/10.1038/srep41268)
38. Nazarenko MS, Markov AV, Koroleva YA, et al. Identification of differentially methylated genes potentially associated with atherosclerosis in humans. *Russian J Cardiol* 2017; 2217085381221084803(10): 42–48. DOI: [10.15829/1560-4071-2017-10-42-48](https://doi.org/10.15829/1560-4071-2017-10-42-48)
39. Sharysh DV, Markov AV, Sleptsov AA, et al. Demethylation of line-1 retrotransposon is associated with the destabilization of atherosclerotic plaques. *Med Genetics* 2020; 202017085381221084803: 19. DOI: [10.25557/2073-7998.2020.05.50-51](https://doi.org/10.25557/2073-7998.2020.05.50-51)
40. Sleptsov AA, Nazarenko MS, Zaitseva AV, et al. Structural variability of the apobec3a-apobec3b locus in patients with carotid atherosclerosis in combination with acute cerebrovascular accidents and type 2 diabetes mellitus. *Med Genetics* 2020; 202017085381221084803: 19. DOI: [10.25557/2073-7998.2020.05.48-49](https://doi.org/10.25557/2073-7998.2020.05.48-49)
41. Sleptsov AA, Nazarenko MS, Zaitseva AV, et al. Somatic mosaicism and structural variability of the gbp3 gene in atherosclerosis. *Atherosclerosis* 2019; 1517085381221084803(4): 46–51. DOI: [10.15372/ATER20190404](https://doi.org/10.15372/ATER20190404)
42. Guselnikova YI, Leader RY, Kazantsev AN, et al. Characteristics of restenosis and progression of atherosclerosis after carotid endarterectomy in the long-term follow-up period. *Complex Prob Cardiovascular Dis* 2019; 817085381221084803(S3–2): 17.
43. Tarasov RS, Kazantsev AN, Anufriev AI, et al. Surgical factors of restenosis of the internal carotid artery after carotid endarterectomy. *Cardiol Cardiovascular Surg* 2018; 1117085381221084803(4): 47–53. DOI: [10.17116/kardio201811447](https://doi.org/10.17116/kardio201811447)
44. Bakhmetyev AS, Chekhonatskaya ML, Dvoenko OG, et al. The incidence of residual stenosis after carotid endarterectomy. *Bulletin Med Internet Conf* 2017; 717085381221084803(4): 683
45. Cherednichenko YV. Endovascular treatment of acute tandem occlusion of the left internal carotid and middle cerebral arteries as a complication of carotid endarterectomy. *Endovascular Neurorengenosurg* 2019; 3017085381221084803(4): 68–78. DOI: [10.26683/2304-9359-2019-4\(30\)-68-78](https://doi.org/10.26683/2304-9359-2019-4(30)-68-78)

## Appendix I

Criteria for CEA of current Russian national guidelines for the management of patients with diseases brachiocephalic arteries (download link [angiolsurgery.org/recommendations/2013/recommendations\\_brachiocephalic.pdf](https://angiolsurgery.org/recommendations/2013/recommendations_brachiocephalic.pdf)).

Recommendations for treatment tactics depending on the neurological symptoms and the degree of stenosis of the carotid arteries:

1. Surgical treatment of stenosis of the carotid arteries is absolutely indicated in symptomatic patients with stenosis of more than 60% (NASCET) [A], if the anticipated rate of periprocedural stroke or mortality is less than 3% for patients with TIA and less than 5% for patients with stroke. The overall mortality rate should not exceed 2%. Carotid endarterectomy is contraindicated in symptomatic patients with stenosis less than 50% [A].

2. It is possible to perform carotid endarterectomy in symptomatic patients with ICA stenosis from 50 to 60%, taking into account the morphological instability of the atherosclerotic plaque (ulceration, haemorrhage into the plaque, intimal flotation and parietal thrombus).
3. It is advisable to perform carotid endarterectomy within 2 weeks from the onset of the last episode of stroke with TIA or minor strokes (mRs < 3), 6–8 weeks after major strokes. Carotid endarterectomy can be performed within days after TIA [B].
4. Carotid endarterectomy can be recommended for asymptomatic patients with stenosis from 70 to 99% if the operational risk is less than 3% [A].

Recommendations for choosing a method of treatment depending on an associated pathology:

1. CEA can be performed in high-risk patients with rates of stroke, death and heart complications within acceptable standards [B].
2. For asymptomatic patients with ‘exceptionally’ high risk (multiple comorbidities at the same time), the best drug therapy may be the optimal choice over invasive procedures [C].
3. CAS is associated with a high risk of embolization in elderly patients of old age. CEA can be performed in elderly patients without increasing the risk of embolization and with an acceptable incidence of neurological and cardiac complications [C].
4. CAS should not be offered to asymptomatic high-risk patients if the likelihood of perioperative stroke and mortality is greater than 3% [C].

On the choice of the method of treatment depending on the features of the vascular and local anatomy and the structure of the atherosclerotic plaque:

1. CAS is indicated in the following cases:
  - restenosis after a previously performed CEA,
  - contralateral paresis of the cranial nerves, previous surgical intervention on the organs of the neck,
  - radiation therapy of the neck organs, the spread of atherosclerotic lesions to the intracranial parts of the ICA.
2. CAS can be recommended in case of high the location of the CCA bifurcation.
3. CAS should be performed with extreme caution with ICA stenosis more than 90% and morphologically unstable atherosclerotic ICA plaque. In these situations, the use of a proximal protection is advisable.
4. CAS is not recommended:
  - in patients with calcified atherosclerotic plaques,
  - with a combination of stenosis with pathological tortuosity of the ICA,
  - with widespread atherosclerotic lesions of the aortic arch and its branches (this is possible only in centres with high surgical activity and a documented low perioperative risk of stroke and death).

## Appendix A

### Abbreviations and acronyms

ECEA	–	eversion carotid endarterectomy
CCEA	–	conventional carotid endarterectomy
ICA	–	internal carotid artery
BARC	–	Bleeding Academic Research Consortium