


Indhold

Indhold	1
1 Fundamental Particles	2
1.1 Units	2
2 Scattering and Decay	3
2.1 Special Relativity	3
2.2 Two-body decay	3
Three-body decay	4
2.3 Decay Amplitudes	4
2.4 Scattering	5
3 Appendix	6
3.1 Four-Vector	6
Contravariant and Covariant Vectors	6
Scalar Product	6
Squared Norm of a Four-Vector	6
The Four-Momentum	7
Lorentz Transformations	7

1 Fundamental Particles



1.1 Units

To simplify calculations we switch to natural units, where $c = \hbar = 1$, $k = 1/4\pi$, and therefore all units are in powers of eV. We can then say that

$$(NU) = \hbar^a c^b (SI)$$

and with that go between systems. Some good conversions to remember is:

$$\hbar c \approx 197 \text{ MeV} \cdot \text{fm}, \quad (1.17)$$

$$\hbar \approx 6.58 \cdot 10^{-16} \text{ eV} \cdot \text{s}. \quad (1.18)$$

2 Scattering and Decay

2.1 Special Relativity

Covariance is a prescription on the form of the physical laws that ensures compliance with the axioms of relativity:

(A1) The physical laws retain the same functional form in all inertial frames.

(A2) The speed of light in vacuum is the same for all inertial observers.

The Four-vector for location is

$$x^\mu = (ct, x, y, z), \quad (2.19)$$

and the transformed vector is:

$$\begin{pmatrix} x'_0 \\ x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} \gamma & \beta\gamma & 0 & 0 \\ \beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \\ x_3 \end{pmatrix}, \quad (2.21)$$

$$x'^\mu = \Lambda^\mu_\nu x^\nu. \quad (2.22)$$

The velocity and acceleration four vectors are:

$$\begin{aligned} v^\mu &= \gamma(c, v_x, v_y, v_z), \\ a^\mu &= \gamma(\dot{\gamma}c, \dot{\gamma}\mathbf{v} + \gamma\mathbf{a}). \end{aligned}$$

The four-momentum is:

$$p^\mu = (m\gamma, m\gamma\mathbf{v}) = (E, \mathbf{p}), \quad (2.40)$$

We can use this to calculate energies in particle interactions, since we know there is conservation of momentum.

$$\sum_{i=1}^{N_i} p_i^\mu = \sum_{j=1}^{N_f} p_j^\mu. \quad (2.43)$$

Some helpful equations are:

$$(p^\mu)^2 = (p^0)^2 - |\mathbf{p}|^2, \quad (2.57)$$

$$E = p^0 = (m^2 + |\mathbf{p}|^2)^{1/2}. \quad (2.58)$$

2.2 Two-body decay

Creation or destruction of particles can only happen if kinematics of Special relativity and of laws of the standard model are conserved. For decays of one particle a into two particles, the momentum must be conserved:

$$p_a^\mu = p_1^\mu + p_2^\mu, \quad (2.64)$$

if we look at the rest frame of a , we get that

$$p_2^2 = (p_a - p_1)^2 = p_a^2 + p_1^2 - 2p_a \cdot p_1, \quad (2.65)$$

$$m_2^2 = m_a^2 + m_1^2 - 2p_a^\mu \cdot p_{1\mu}. \quad (2.66)$$

Here we used that the norm of a single particles momentum is it's rest mass. Because we are in the rest frame of a

$$p_a^\mu p_{1\mu} = g_{\mu\nu} p_a^\mu p_1^\nu = m_a E_1, \quad (2.67)$$

and with this we have:

$$E_1 = \frac{m_a^2 + m_1^2 - m_2^2}{2m_a}, \quad E_2 = \frac{m_a^2 + m_2^2 - m_1^2}{2m_a}. \quad (2.68)$$

Three-body decay

In a three-body decay $a \rightarrow 1 + 2 + 3$, we have too many variables, and can therefore only say something about the minimum and maximum energy of particle 1.

$$m_1 \leq E_1^{CM} \leq \frac{m_a^2 + m_1^2 - (m_2 + m_3)^2}{2m_a}. \quad (2.71)$$

2.3 Decay Amplitudes

In a general term we can use the **relativistic golden rule** for unstable particles that decay into N_f particles. We use this rule to calculate the transition rate, which is the probability per unit time that the particle in the initial state decay into the final state:

$$\Gamma_{i \rightarrow f} = \frac{1}{2E_a} \int \prod_{i=1}^{N_f} \frac{d^4 p_i}{(2\pi)^4} |\mathcal{M}|^2 (2\pi)^4 \delta^4 \left(p_a - \sum_{j=1}^{N_f} p_j \right) \times (2\pi) \delta(p_i^2 - m_i^2). \quad (2.90)$$

The decay Lorentz transforms as

$$\Gamma'_{i \rightarrow f} = \Gamma_{i \rightarrow f} / \gamma. \quad (2.94)$$

And is related to the lifetime of a particle by

$$\tau = 1/\Gamma. \quad (2.96)$$

But most particles decay in multiple nodes and we therefore have to generalize as:

$$\tau = \frac{1}{\Gamma} = \frac{1}{\sum_{i=1}^N \Gamma_i}. \quad (2.101)$$

The **Branching Ratio** of node i is defined as:

$$\text{BR}(i) = \frac{\Gamma_i}{\Gamma}. \quad (2.102)$$

2.4 Scattering

A measurement for scattering in QM is the **differential cross section**, which is the ratio between the number of particles that are scattered in a solid angle between Ω and $\Omega + d\Omega$ per unit time divided by the flux (the rate of incoming particles per unit surface). Or in other words the likelihood that a particle is scattered at an angle Ω .

$$\frac{d\sigma}{d\Omega}(E, \Omega) = \frac{1}{F} \frac{dN_s}{d\Omega}. \quad (2.113)$$

3 Appendix

3.1 Four-Vector

A four-vector a^μ is an object in Minkowski space-time with four components. It is written as:

$$a^\mu = (a^0, a^1, a^2, a^3) = (a^0, \mathbf{a})$$

where a^0 is the time-like component and $\mathbf{a} = (a^1, a^2, a^3)$ is the vector of the space-like components.

Contravariant and Covariant Vectors

A four-vector can be in two forms:

- **Contravariant** (upper index): $a^\mu = (a^0, a^1, a^2, a^3)$
- **Covariant** (lower index): $a_\mu = (a_0, a_1, a_2, a_3)$

The transformation between them uses the Minkowski metric tensor $g_{\mu\nu}$.

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad (2.31)$$

The rule to lower an index is $a_\mu = g_{\mu\nu}a^\nu$. This results in:

$$a_\mu = (a^0, -a^1, -a^2, -a^3) = (a^0, -\mathbf{a})$$

Scalar Product

The scalar product of two four-vectors a^μ and b^μ is a Lorentz invariant scalar, denoted as $a \cdot b$ or $a_\mu b^\mu$.

$$a \cdot b = a_\mu b^\mu = g_{\mu\nu}a^\mu b^\nu \quad (1)$$

Using the Einstein summation convention, this expands to:

$$a \cdot b = a^0 b^0 - a^1 b^1 - a^2 b^2 - a^3 b^3 = a^0 b^0 - \mathbf{a} \cdot \mathbf{b}$$

Squared Norm of a Four-Vector

The squared norm of a four-vector a^μ is its scalar product with itself, denoted a^2 .

$$a^2 = a_\mu a^\mu = (a^0)^2 - (a^1)^2 - (a^2)^2 - (a^3)^2 = (a^0)^2 - |\mathbf{a}|^2 \quad (2)$$

The sign of the squared norm classifies the four-vector:

- $a^2 > 0$: **Time-like**
- $a^2 < 0$: **Space-like**
- $a^2 = 0$: **Light-like** or **null**

The Four-Momentum

The four-momentum p^μ is a key four-vector in relativistic kinematics. In natural units ($c = 1$), it is defined as:

$$p^\mu = (E, \mathbf{p}) = (m\gamma, m\gamma\mathbf{v}) \quad (2.40)$$

where $E = m\gamma$ is the relativistic energy, $\mathbf{p} = m\gamma\mathbf{v}$ is the relativistic three-momentum, m is the rest mass, and $\gamma = 1/\sqrt{1-v^2}$ is the Lorentz factor.

The squared norm of the four-momentum is the squared rest mass, an invariant:

$$p^2 = p_\mu p^\mu = E^2 - |\mathbf{p}|^2 = m^2$$

This yields the energy-momentum relation:

$$E = \sqrt{|\mathbf{p}|^2 + m^2} \quad (2.58)$$

Lorentz Transformations

A four-vector a^μ transforms between inertial frames via a Lorentz transformation matrix Λ_ν^μ .

$$a'^\mu = \Lambda_\nu^\mu a^\nu$$

For a boost with velocity v along the x^1 -axis, the matrix is:

$$\Lambda_\nu^\mu = \begin{pmatrix} \gamma & -\beta\gamma & 0 & 0 \\ -\beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

where $\beta = v$ in natural units. Applying this to the four-momentum $p^\mu = (E, p_x, p_y, p_z)$ gives:

$$E' = \gamma(E - \beta p_x)$$

$$p'_x = \gamma(p_x - \beta E)$$

$$p'_y = p_y$$

$$p'_z = p_z$$