

The Chain Rule

The chain rule expresses the relation between the conditional entropy and the joint/marginal entropies of the variables involved. We first state and prove the chain rule for two random variables, and then generalize it to n variables.

Proposition: Chain Rule

Let X and Y be random variables. Then

$$H(XY) = H(X) + H(Y|X).$$

Proof hint

We encourage you to try to prove this for yourself. As a starting point, write out the definition of $H(XY)$, and rewrite the terms $P_{XY}(x, y)$ into a conditional form in order to relate it to $H(Y|X)$.

Show full proof

The chain rule is a matter of rewriting:

$$\begin{aligned} H(XY) &= - \sum_{x,y} P_{XY}(x, y) \log P_{XY}(x, y) \\ &= - \sum_{x,y} P_{XY}(x, y) \log (P_X(x) P_{Y|X}(y|x)) \\ &= - \sum_{x,y} P_{XY}(x, y) \log P_X(x) - \sum_{x,y} P_{XY}(x, y) \log P_{Y|X}(y|x) \\ &= - \sum_x P_X(x) \log P_X(x) - \sum_{x,y} P_{XY}(x, y) \log P_{Y|X}(y|x) \\ &= - \sum_x P_X(x) \log P_X(x) - \sum_x P_X(x) \sum_y P_{Y|X}(y|x) \log P_{Y|X}(y|x) \\ &= H(X) + H(Y|X). \end{aligned}$$

This was to be shown.

The chain rule immediately results in the so-called 'independence bound':

Corollary: Subadditivity (independence bound)

$$H(XY) \leq H(X) + H(Y).$$

Equality holds iff X and Y are independent.

Proof

$$H(XY) = H(X) + H(Y|X) \leq H(X) + H(Y),$$

where the equality is due to the chain rule and the inequality is due to the upper bound on the conditional entropy that $H(Y|X) \leq H(Y)$ (and equal iff X and Y are independent).

We can naturally generalize the definition of conditional entropy by applying it to the conditional distribution $P_{XY|\mathcal{A}}$; this results in $H(X|Y, \mathcal{A})$, the entropy of X given Y and conditioned on the event \mathcal{A} . Since the entropy is a function of the *distribution* of a random variable, the chain rule also holds when conditioning on an event \mathcal{A} . Furthermore, it holds that

$$H(X|YZ) = \sum_z P_Z(z) H(X|Y, Z=z),$$

which is straightforward to verify. With this observation, we see that the chain rule generalizes as follows.

Corollary: generalized chain rule

Let X, Y and Z be random variables. Then

$$H(XY|Z) = H(X|Z) + H(Y|XZ).$$

Inductively applying the (generalized) chain rule implies that for any sequence X_1, \dots, X_n of random variables:

$$H(X_1 \cdots X_n) = H(X_1) + H(X_2|X_1) + \cdots + H(X_n|X_{n-1} \cdots X_1).$$

Combining this with the upper bound on the conditional entropy, we see that subadditivity generalizes to

$$H(X_1 \cdots X_n) \leq \sum_{i=1}^n H(X_i).$$