

The title

First Author¹ & Ernst-August Doelle^{1,2}

¹ Wilhelm-Wundt-University

² Konstanz Business School

Author Note

Add complete departmental affiliations for each author here. Each new line herein must be indented, like this line.

Enter author note here.

The authors made the following contributions. First Author: Conceptualization, Writing - Original Draft Preparation, Writing - Review & Editing; Ernst-August Doelle: Writing - Review & Editing, Supervision.

Correspondence concerning this article should be addressed to First Author, Postal address. E-mail: my@email.com

Abstract

One or two sentences providing a **basic introduction** to the field, comprehensible to a scientist in any discipline.

Keywords: keywords

Word count: X

The title

Results

The aggregated data-set describes 193,318 observations of daily physical activity and sleep from 24,752 unique participants. Table 1 shows demographic information for all participants. Observations were not uniform across the days of the week $\chi^2_{(6)} = 108.75$, $p = < .001$. There were fewer observations on Monday ($z = -8.92$) and Sunday ($z = -3.54$); and more on Wednesday ($z = 4.00$). A table of study characteristics can be found in supplementary materials.

The effects of physical activity volume on sleep

We estimated the effects of physical activity on sleep (RQ1) using mixed-effects models. We estimated the effect of physical activity volume on sleep by age, and the results are presented in Table 2 and Figure 1. There was no meaningful relationship between physical activity volume and sleep duration. However, we observed a curvilinear relationships between physical activity volume and sleep efficiency, onset, and regularity, all of which interacted with age. Sleep efficiency improved with greater physical activity volume, but improvements tapered off for older individuals. Physical activity volume and sleep onset had a positive association for younger individuals, but a negative association for older individuals, where sleep onset was reduced among those with the highest physical activity. There was a strong positive association between physical activity volume and sleep regularity, which was strongest among older participants. For participants aged 35 years and above, this link weakened among those with a physical activity volume greater than two standard deviations.

The effects of physical activity intensity on sleep

We estimated how physical activity intensity affects sleep across different age groups. We present the results controlling for sex, SES, and BMI, in Table 2 and Figure 2. We found

that higher physical activity intensity is directly proportional to longer sleep duration and better sleep efficiency. In the case of older participants, physical activity intensity had a U-shaped relationship with sleep onset, meaning that individuals with very low or very high physical activity intensity had longer sleep onset. We also found a strong link between physical activity intensity and improved sleep regularity, which weakened at higher intensity levels.

The effects of sleep duration on physical activity

We estimated the effect of sleep duration on physical activity by age Results, controlling for sex, SES, and BMI are presented in Table 3 and Figure 3. As age increases, both physical activity volume and intensity decrease. We found a subtle inverted U-shaped relationship between average sleep duration and physical activity volume, where the highest volume of physical activity was linked to average sleep duration.

The effects of sleep efficiency on physical activity

We estimated the effect of sleep efficiency on physical activity by age. Results, controlling for sex, SES, and BMI are presented in Table 3 and Figure 4. We did not find a relationship between physical activity volume and sleep efficiency. However, there was a subtle U-shaped relationship where individuals with above-average sleep efficiency engaged in more intense physical activity.

The effects of sleep onset on physical activity

We estimated the effect of sleep onset on physical activity by age Results, controlling for sex, SES, and BMI are presented in Table 3 and Figure 5. There were strong U-shaped relationships where average sleep onset was linked to the highest levels of physical activity

volume and intensity. The U-shaped relationship between sleep onset and physical activity volume attenuated for older participants.

The effects of sleep regularity on physical activity

We estimated the effect of sleep regularity on physical activity by age. Results, controlling for sex, SES, and BMI are presented in Table 3 and Figure 6. There was a U-shaped relationship between sleep regularity and physical activity volume. Participants with below-average sleep regularity tended to have average physical activity volume. Increases in regularity above the average were linked to greater physical activity volume. There was a strong linear relationship between sleep regularity and physical activity intensity which slightly attenuated with age. Greater sleep regularity was associated with greater physical activity the following day.

Table 1
Participant characteristics

Characteristic	Age group				66+ years
	2-11 years	12-18 years	19-35 years	36-65 years	
Numeric variables					
N	10,065	3,431	373	4,187	6,615
Valid weartime hours	21.77 (3.03)	21.20 (3.97)	21.06 (3.46)	22.57 (2.49)	23.78 (1.03)
PA volume	43.68 (25.56)	42.28 (16.83)	41.37 (11.44)	39.80 (13.35)	31.23 (10.39)
PA intensity	-2.11 (0.19)	-2.20 (0.19)	-2.35 (0.20)	-2.50 (0.19)	-2.67 (0.22)
Sleep duration	429.14 (120.91)	374.98 (131.77)	334.95 (108.40)	399.97 (70.13)	397.38 (74.34)
Sleep efficiency	0.78 (0.12)	0.81 (0.15)	0.87 (0.10)	0.87 (0.07)	0.86 (0.06)
Sleep onset	20.89 (1.14)	22.17 (2.03)	24.03 (1.73)	23.66 (1.37)	24.41 (1.32)
Sleep regularity	54.58 (12.72)	54.07 (13.28)	53.13 (13.30)	59.11 (11.76)	54.68 (12.59)
Age	9.58 (1.36)	13.73 (2.40)	22.56 (6.04)	57.77 (6.91)	72.22 (4.74)
BMI	17.98 (3.28)	20.43 (4.01)	24.77 (5.94)	26.99 (4.92)	27.08 (4.41)
Accelerometer Wear Location					
Hip	7,487 (74.39%)	1,321 (38.50%)	-	-	-
Wrist	2,578 (25.61%)	2,110 (61.50%)	373 (100.00%)	4,187 (100.00%)	6,615 (100.00%)
Region					
Africa	910 (9.04%)	175 (5.10%)	-	-	-
Asia	635 (6.31%)	95 (2.77%)	-	-	-
Europe	3,002 (29.83%)	1,238 (36.08%)	107 (28.69%)	3,772 (90.09%)	6,612 (99.95%)
North america	1,342 (13.33%)	85 (2.48%)	-	-	-
Oceania	1,798 (17.86%)	694 (20.23%)	146 (39.14%)	415 (9.91%)	3 (0.05%)
South america	2,378 (23.63%)	1,144 (33.34%)	120 (32.17%)	-	-
Season					

Table 1 continued

Characteristic	Age group			
	2-11 years	12-18 years	19-35 years	36-65 years
Autumn	3,310 (32.89%)	821 (23.93%)	34 (9.12%)	1,003 (23.96%)
Spring	2,320 (23.05%)	1,718 (50.07%)	272 (72.92%)	1,145 (27.35%)
Summer	1,260 (12.52%)	372 (10.84%)	30 (8.04%)	956 (22.83%)
Winter	3,175 (31.54%)	520 (15.16%)	37 (9.92%)	1,083 (25.87%)
Sex				
Female	5,259 (52.25%)	1,757 (51.21%)	232 (62.20%)	2,250 (53.74%)
Male	4,806 (47.75%)	1,674 (48.79%)	141 (37.80%)	1,937 (46.26%)
Sleep Conditions Reported				
Yes	49 (0.49%)	1 (0.03%)	-	268 (6.40%)
Socioeconomic Status				
Low	3,576 (35.53%)	898 (26.17%)	67 (17.96%)	1,180 (28.18%)
Medium	2,798 (27.80%)	1,228 (35.79%)	148 (39.68%)	1,681 (40.15%)
High	3,691 (36.67%)	1,305 (38.04%)	158 (42.36%)	1,326 (31.67%)

Note. For categorical variables the value is the count, and percentage. For numeric variables the value is the Mean and SD.

N = 24,752

Table 2

Physical activity predicting sleep controlling for SES, sex, and BMI.

Term	Physical activity volume (h)				Physical activity intensity (z)					
	β	[95% CI]	SE	t	p	β	[95% CI]	SE	t	p
<i>Sleep duration (z)</i>										
(Intercept)	0.16	[0.63, 0.96]	0.40	0.41	.685	0.17	[-0.15, 0.49]	0.16	1.06	.298
Log pa volume	-0.19	[-0.61, 0.22]	0.21	-0.91	.365	0.08	[0.04, 0.13]	0.02	3.44	.001
Age	-0.02	[-0.04, 0.00]	0.01	-1.53	.125	0.00	[-0.01, 0.00]	0.00	-0.79	.430
Log pa volume ²	0.05	[-0.01, 0.11]	0.03	1.74	.082	0.02	[-0.01, 0.04]	0.01	1.20	.229
Log pa volume × age	0.01	[0.00, 0.02]	0.01	1.55	.120	0.00	[0.00, 0.00]	0.00	-0.92	.358
Age × log pa volume ²	0.00	[0.00, 0.00]	0.00	-1.64	.102	0.00	[0.00, 0.00]	0.00	-1.14	.256
<i>Sleep efficiency (z)</i>										
(Intercept)	-1.81	[-2.60, -1.02]	0.40	-4.49	< .001	0.27	[-0.02, 0.55]	0.15	1.83	.070
Log pa volume	1.03	[0.60, 1.45]	0.22	4.77	< .001	0.04	[0.00, 0.09]	0.02	1.75	.080
Age	0.03	[0.01, 0.05]	0.01	2.48	.013	0.00	[0.00, 0.01]	0.00	0.87	.382
Log pa volume ²	-0.12	[-0.18, -0.06]	0.03	-4.07	< .001	-0.02	[-0.05, 0.00]	0.01	-1.63	.104
Log pa volume × age	-0.01	[-0.02, 0.00]	0.01	-1.84	.065	0.00	[0.00, 0.00]	0.00	-1.66	.097
Age × log pa volume ²	0.00	[0.00, 0.00]	0.00	1.38	.169	0.00	[0.00, 0.00]	0.00	0.91	.365
<i>Sleep onset (z)</i>										
(Intercept)	-0.86	[-1.50, -0.21]	0.33	-2.59	.010	0.04	[-0.30, 0.39]	0.18	0.24	.814
Log pa volume	0.54	[0.23, 0.85]	0.16	3.39	.001	-0.06	[-0.10, -0.03]	0.02	-3.59	< .001
Age	0.02	[0.01, 0.04]	0.01	2.87	.004	0.00	[0.00, 0.00]	0.00	-0.43	.670
Log pa volume ²	-0.08	[-0.12, -0.04]	0.02	-3.53	< .001	-0.02	[-0.04, 0.00]	0.01	-2.12	.034
Log pa volume × age	-0.01	[-0.02, 0.00]	0.00	-2.38	.017	0.00	[0.00, 0.00]	0.00	0.34	.730
Age × log pa volume ²	0.00	[0.00, 0.00]	0.00	1.73	.083	0.00	[0.00, 0.00]	0.00	2.39	.017
<i>Sleep regularity (z)</i>										

Table 2 continued

Term	Physical activity volume (ln)				Physical activity intensity (z)					
	β	[95% CI]	SE	t	p	β	[95% CI]	SE	t	p
(Intercept)	-2.50	[-3.24, -1.76]	0.38	-6.63	< .001	0.55	[0.28, 0.83]	0.14	3.97	< .001
Log pa volume	1.28	[0.89, 1.67]	0.20	6.42	< .001	0.27	[0.23, 0.32]	0.02	12.23	< .001
Age	-0.03	[-0.05, -0.01]	0.01	-3.15	.002	0.00	[-0.01, 0.00]	0.00	-1.14	.256
Log pa volume ²	-0.13	[-0.18, -0.07]	0.03	-4.51	< .001	-0.11	[-0.13, -0.08]	0.01	-8.08	< .001
Log pa volume × age	0.02	[0.01, 0.03]	0.01	3.53	< .001	0.00	[-0.01, 0.00]	0.00	-9.84	< .001
Age × log pa volume ²	0.00	[0.00, 0.00]	0.00	-3.73	< .001	0.00	[0.00, 0.00]	0.00	2.80	.005

Note. Adjusted for SES, sex, and BMI. Outcomes variables are listed in the column headers.

Sleep predicting physical activity controlling for SES, sex, and BMI

Term	Physical activity volume (ln)				Physical activity intensity (z)					
	β	[95% CI]	SE	t	p	β	[95% CI]	SE	t	p
Sleep duration (z)										
(Intercept)	4.31	[4.13, 4.49]	0.09	46.20	< .001	1.35	[1.09, 1.61]	0.13	10.07	< .001
Sleep duration	0.00	[0.01, 0.01]	0.01	-0.11	.910	-0.01	[-0.03, 0.01]	0.01	-0.91	.364
Age	-0.01	[-0.01, -0.01]	0.00	-8.44	< .001	-0.03	[-0.03, -0.02]	0.00	-10.18	< .001
Sleep duration ²	-0.01	[-0.01, 0.00]	0.00	-3.49	< .001	-0.02	[-0.03, -0.01]	0.00	-3.19	.001
Sleep duration × age	0.00	[0.00, 0.00]	0.00	-1.50	.133	0.00	[0.00, 0.00]	0.00	2.23	.026
Age × Sleep duration ²	0.00	[0.00, 0.00]	0.00	-1.36	.175	0.00	[0.00, 0.00]	0.00	1.20	.229
Sleep efficiency (z)										
(Intercept)	4.30	[4.12, 4.48]	0.09	46.15	< .001	1.33	[1.07, 1.60]	0.14	9.81	< .001
Sleep efficiency	0.02	[0.00, 0.03]	0.01	2.67	.008	0.02	[0.00, 0.05]	0.01	1.73	.084
Age	-0.01	[-0.01, -0.01]	0.00	-8.48	< .001	-0.03	[-0.03, -0.02]	0.00	-10.11	< .001
Sleep efficiency ²	0.00	[0.00, 0.00]	0.00	-0.17	.868	0.00	[0.00, 0.01]	0.00	0.62	.536
Sleep efficiency × age	0.00	[0.00, 0.00]	0.00	-1.94	.053	0.00	[0.00, 0.00]	0.00	-1.41	.157
Age × Sleep efficiency ²	0.00	[0.00, 0.00]	0.00	-0.40	.690	0.00	[0.00, 0.00]	0.00	-0.24	.808
Sleep onset (z)										
(Intercept)	4.31	[4.12, 4.49]	0.09	46.25	< .001	1.34	[1.08, 1.61]	0.14	9.92	< .001
Sleep onset	0.00	[-0.01, 0.02]	0.01	0.74	.462	0.02	[0.00, 0.04]	0.01	1.87	.062
Age	-0.01	[-0.01, -0.01]	0.00	-8.46	< .001	-0.03	[-0.03, -0.02]	0.00	-10.09	< .001
Sleep onset ²	-0.01	[-0.02, 0.00]	0.01	-1.27	.205	-0.01	[-0.03, 0.01]	0.01	-0.69	.491
Sleep onset × age	0.00	[0.00, 0.00]	0.00	0.18	.858	0.00	[0.00, 0.00]	0.00	-0.90	.369
Age × Sleep onset ²	0.00	[0.00, 0.00]	0.00	-0.46	.645	0.00	[0.00, 0.00]	0.00	1.07	.285
Sleep regularity (z)										

Table 3 continued

Term	Physical activity volume (h)				Physical activity intensity (z)					
	β	[95% CI]	SE	t	p	β	[95% CI]	SE	t	p
(Intercept)	4.27	[4.10, 4.45]	0.09	47.62	< .001	1.33	[1.07, 1.59]	0.13	9.95	< .001
Sleep regularity	0.08	[0.07, 0.09]	0.01	13.20	< .001	0.09	[0.07, 0.11]	0.01	7.64	< .001
Age	-0.01	[-0.01, -0.01]	0.00	-8.62	< .001	-0.03	[-0.03, -0.02]	0.00	-10.11	< .001
Sleep regularity ²	-0.01	[-0.01, 0.00]	0.00	-2.34	.020	0.00	[-0.02, 0.01]	0.01	-0.63	.528
Sleep regularity × age	0.00	[0.00, 0.00]	0.00	-3.13	.002	0.00	[0.00, 0.00]	0.00	-5.16	< .001
Age × Sleep regularity ²	0.00	[0.00, 0.00]	0.00	-0.78	.433	0.00	[0.00, 0.00]	0.00	-1.66	.096

Note. Adjusted for SES, sex, and BMI. Outcomes variables are listed in the row headers.




Figure 1. Sleep metrics on physical activity volume. The panels on the left show the curvilinear relationship between PA volume and each of the sleep outcomes at the ages indicated in each column. The panels on the right show the same relationships but by age continuously. The white band indicates predictions which were not significantly different from zero. These plots demonstrate exact turning points where the effects of PA volume change by age.




Figure 2. Sleep metrics on physical activity intensity. The panels on the left show the curvilinear relationship between PA intensity and each of the sleep outcomes at the ages indicated in each column. The panels on the right show the same relationships but by age continuously. The white band indicates predictions which were not significantly different from zero.




Figure 3. Physical activity by sleep duration. The panels on the left show the curvilinear relationship between sleep duration and each of the physical activity outcomes at the ages indicated in each column. The panels on the right show the same relationships but by age continuously. The white band indicates predictions which were not significantly different from zero.




Figure 4. Physical activity by sleep efficiency. The panels on the left show the curvilinear relationship between sleep efficiency and each of the physical activity outcomes at the ages indicated in each column. The panels on the right show the same relationships but by age continuously. The white band indicates predictions which were not significantly different from zero.




Figure 5. Physical activity by sleep onset. The panels on the left show the curvilinear relationship between sleep onset and each of the physical activity outcomes at the ages indicated in each column. The panels on the right show the same relationships but by age continuously. The white band indicates predictions which were not significantly different from zero.




Figure 6. Physical activity by sleep regularity. The panels on the left show the curvilinear relationship between sleep regularity and each of the physical activity outcomes at the ages indicated in each column. The panels on the right show the same relationships but by age continuously. The white band indicates predictions which were not significantly different from zero.