Running head: TITLE 1

The title

First Author¹ & Ernst-August Doelle^{1,2}

- ¹ Wilhelm-Wundt-University
- ² Konstanz Business School

Author Note

- Add complete departmental affiliations for each author here. Each new line herein must be indented, like this line.
- Enter author note here.

5

- The authors made the following contributions. First Author: Conceptualization,
- Writing Original Draft Preparation, Writing Review & Editing; Ernst-August Doelle:
- Writing Review & Editing, Supervision.
- Correspondence concerning this article should be addressed to First Author, Postal address. E-mail: my@email.com

TITLE 2

14 Abstract

One or two sentences providing a **basic introduction** to the field, comprehensible to a

scientist in any discipline.

17 Keywords: keywords

Word count: X

TITLE 3

The title

20 Results

The aggregated data-set describes 316,835 observations of daily physical activity and sleep from 24,935 unique participants. Table ?? shows demographic information. A table of study characteristics can be found in supplementary materials.

The effects of physical activity volume on sleep

We explored the effects of physical activity on sleep (RQ1) using mixed-effects models. 25 We estimated the effect of physical activity volume on sleep by age, and the results are 26 presented in Table 1 and Figure 1. There was no meaningful relationship between physical 27 activity volume and sleep duration. However, a positive curvilinear relationship was observed 28 between physical activity volume and sleep efficiency, onset, and regularity, which interacted 29 with age. Sleep efficiency improved with greater physical activity volume, but improvements 30 tapered off for older individuals. Physical activity volume and sleep onset had a positive 31 association for younger individuals, but a negative association for older individuals, where 32 sleep onset was reduced among those with the highest physical activity. There was a strong 33 positive association between physical activity volume and sleep regularity, which was strongest among older participants. For participants aged 35 years and above, this link weakened among those with a physical activity volume greater than two standard deviations.

Table 1 $Sleep\ on\ physical\ activity\ volume\ controlling\ for\ SES,\ gender\ and\ BMI$

Term	β [95% CI]	SE	t	p
Sleep duration				
(Intercept)	-0.37 [-0.94, 0.20]	0.29	-1.28	.291
Scale pa volume	$0.03 \ [0.00, \ 0.07]$	0.02	1.96	.050
Age	0.00 [-0.01, 0.00]	0.00	-0.29	.797
Scale pa volume ²	0.00 [-0.01, 0.01]	0.00	0.11	.912
Scale pa volume:age	$0.00 \ [0.00, \ 0.00]$	0.00	0.92	.359
Age:scale pa volume ²	$0.00 \ [0.00, \ 0.00]$	0.00	-0.94	.349
Sleep efficiency				
(Intercept)	-0.90 [-1.29, -0.50]	0.20	-4.46	.001
Scale pa volume	$0.23 \ [0.20, \ 0.27]$	0.02	12.97	< .001
Age	$0.01 \ [0.01, \ 0.01]$	0.00	9.99	.002
Scale pa volume ²	-0.02 [-0.03, -0.01]	0.00	-4.77	< .001
Scale pa volume:age	$0.00 \ [0.00, \ 0.00]$	0.00	-5.16	< .001
Age:scale pa volume ²	$0.00 \ [0.00, \ 0.00]$	0.00	-0.50	.617
Sleep onset				
(Intercept)	-0.98 [-1.50, -0.47]	0.26	-3.77	.060
Scale pa volume	$0.11 \ [0.09, \ 0.14]$	0.01	8.32	< .001
Age	$0.02 \ [0.02, \ 0.02]$	0.00	9.81	.007
$Scale pa volume^2$	-0.01 [-0.02, -0.01]	0.00	-4.73	< .001
Scale pa volume:age	$0.00 \ [0.00, \ 0.00]$	0.00	-8.70	< .001
Age:scale pa volume ²	$0.00 \ [0.00, \ 0.00]$	0.00	2.34	.034
Sleep regularity				
(Intercept)	-0.13 [-0.39, 0.13]	0.13	-0.96	.415
Scale pa volume	$0.33 \ [0.29, \ 0.36]$	0.02	19.10	< .001
Age	$0.01 \ [0.00, \ 0.01]$	0.00	5.07	.019
$Scale pa volume^2$	-0.03 [-0.03, -0.02]	0.00	-5.74	< .001
Scale pa volume:age	$0.00 \ [0.00, \ 0.00]$	0.00	0.64	.540
Age:scale pa volume ²	$0.00 \ [0.00, \ 0.00]$	0.00	-3.33	.006

Note. Adjusted for SES, BMI, and sex.

TITLE 5




Figure 1. Sleep metrics on Physical activity volume