

Authors:

Marca L. Sipski, MD
J. Scott Richards, PhD, ABPP

Affiliations:

From the Department of Physical
Medicine and Rehabilitation,
University of Alabama at
Birmingham, Birmingham, Alabama.

Correspondence:

All correspondence and requests for
reprints should be addressed to Marca
L. Sipski, MD, University of Alabama,
Spain Rehabilitation Center, 1717 6th
Ave. South, Room 157B,
Birmingham, AL 35233.

Editor's Note:

Tables are available at
www.amjphysmedrehab.com

Disclosures:

FIM™ is a trademark of Uniform
Data System for Medical
Rehabilitation, a division of UB
Foundation Activities, Inc.

0894-9115/06/8504-0310/0
*American Journal of Physical
Medicine & Rehabilitation*
Copyright © 2006 by Lippincott
Williams & Wilkins

DOI: 10.1097/01.phm.0000202105.87011.bf

**REVIEW & ANALYSIS**

Spinal Cord Injury Rehabilitation State of the Science

ABSTRACT

Sipski ML, Richards JS: Spinal cord injury rehabilitation: State of the science.
Am J Phys Med Rehabil 2006;85:310–342.

Key Words: Spinal Cord Injury, Rehabilitation Research, Outcomes

Spinal cord injury (SCI) is a low-incidence phenomenon (15–40 cases per million), resulting in about 11,000 new cases in the United States per year, and a prevalence near 250,000. Men sustain this injury about four times as often as women; age of onset is typically in the young 30s. As measured by the Model Systems database, over time, proportionately fewer whites have sustained SCIs, and more cervical injuries have occurred. Motor vehicle accidents, falls, violence, and sports have been the leading causes of SCI, respectively, over time, based on the Model Systems data, with a period of time in the 1990s in which violence superseded falls as the second-leading cause.¹ In very young children who sustain SCI, gender and injury characteristics are different from adults, becoming much more similar when onset is in adolescence.²

Research related to the epidemiology of SCI has largely been descriptive, with a focus on primary prevention.³ This information is important for estimating the impact of SCI as measured by incidence, prevalence, survival, and characteristics of those who receive such injuries. Methodologies for capturing this information have varied widely from retrospective attempts to capture all cases of a particular etiology over a given time period^{4,5} to prospective attempts to capture all cases in a geographic area.^{6,7,8} Countries with centralized health care and record keeping have come the closest to population-based estimates of incidence and prevalence,^{5,6} and several state-based registries in the United States have attempted to do the same.⁴ The National Institute on Disability and Rehabilitation Research–funded SCI Model Systems of Care database, though not population-based, has been collecting epidemiologic data on a large sample of persons with SCI back to 1973, with enough of the funded systems being the same throughout that period that important trends have been identified.⁹

The neurologic injury with its resultant paralysis and loss of sensation is only one of a myriad of physical consequences of SCI. Thus, the care of persons with SCI is not limited to the treatment of the neurologic injury. Medical consequences as a result of altered neurologic input such as the development of respiratory failure or neurogenic bladder and resultant complications such as pneumonia and urinary tract infections affect most organ systems. In addition, the psychological impact of the injury is vast, and complications such as depression and anxiety are frequent. Rehabilitation has traditionally focused on

maximizing the person's recovery from the injury and returning him or her to home as independently as possible. It has included such techniques as performing range of motion, strengthening weakened muscles, and retraining bladder and bowel functioning. However, once the milestone of returning home with a new SCI is crossed, the individual must conquer the barriers of societal reintegration with his or her altered body habitus.

The care and treatment of persons with SCIs has drastically improved, and we are now concerned with improving quality of life rather than merely extending survival. Despite this, many shortfalls are noted in the research supporting our past and current practices in the field of SCI. The purpose of this article is to develop an understanding of the level of evidence about practice patterns in the field of SCI and to make recommendations for where and how future research initiatives should be targeted.

METHODS

We realized that performing a review of all topics related to the treatment and needs of persons with SCIs would be impossible; therefore, we limited our search to a series of medical and psychosocial topics deemed most crucial to promote the rehabilitation of persons with SCI. We chose to study pharmacologic and surgical treatment of the acute injury, followed by neurologic recovery and functional potential. Medical topics that were searched included respiratory, pneumonia, neurogenic bowel dysfunction, acute abdomen, gastric ulcer, gallstone, management of neurogenic bladder, urinary tract infection, bladder stone, bladder carcinoma, neurogenic sexual dysfunction, pressure sore, deep vein thrombosis, musculoskeletal complications, heterotopic ossification and osteoporosis, metabolic dysfunction, body composition, spasticity, and pain. Outcome measures were reviewed. In addition, we searched the terms rehabilitation therapies, exercise, body-weight-supported treadmill training, and electrical stimulation. Psychosocial topics included community integration, depression, substance abuse, anxiety, family and caregiver issues, coping effectiveness, and life satisfaction. We also searched vocational issues, assistive technology, and aging.

We performed searches using MEDLINE, PubMed, and Psych Lit in addition to performing secondary reviews and conferring with known experts in the fields. We also surveyed ongoing research. After group discussions with other Model Systems representatives, we chose to include in the evidence tables the level I and level II studies published in the English language (as per the American Academy of Neurology Guidelines) from Fall 1999–2004 with a sample size of ≥ 20 for both

subjects and controls. For those areas in which limited recent research was available, we sought the most recent and meaningful articles published before 1999. In addition, we determined whether the article's focus was diagnostic, prognostic, or therapeutic. Published clinical practice guidelines were also reviewed. Because of the paucity of research in certain topics when level I or II evidence was not available, other level III and IV studies deemed important were reviewed and evaluated. If not specified in the text or tables as a level I, II, or III study, the reader can assume that the study reviewed was a level IV study. Based on this information, recommendations have been made for future directions in the field of SCI. Throughout the sections of this article, to the extent that various types of evidence were available for different topics, we have attempted to follow the pattern of discussion of diagnostic studies first followed by prognostic and therapeutic studies.

CLINICAL PRACTICE AND INTERVENTIONS RELATED TO THE NEUROLOGIC INJURY

Acute Management of Spinal Cord Injury Pharmacologic Management

Much attention has been paid to the acute management of SCI and whether pharmacologic therapy is justified. The National Acute Spinal Cord Injury Study (NASCIS) trials,^{10,11,12,13} all published before 1999, were the first to document a potential beneficial treatment for SCI and quickly led to methylprednisolone (MPSS) being known as a standard of care for acute SCI management. However, these results have been questioned due to the use of post hoc analyses,¹⁴ inappropriate use of statistics,¹⁵ and the potential harmful side effects of MPSS. Based on reviews of the literature,¹⁶ the use of MPSS post-SCI has been deemed a recommendation for acute, nonpenetrating SCI within 8 hrs of trauma. It is not recommended for acute nonpenetrating SCI over 8 hrs postinjury or for penetrating SCI. It is recommended that acute patients with SCI who present within 3 to 8 hrs postinjury may reasonably be given 48 hrs as opposed to 24 hrs of therapy.

Other drugs that were evaluated as part of the NASCIS studies included naloxone, a nonspecific opioid receptor agonist, studied as part of NASCIS 2, and tirilazad, a member of the 21-aminosteroid family of antioxidants called lazaroids developed to prevent lipid peroxidation without activation of glucocorticoid receptors, studied in NASCIS 3. Although a subsequent analysis¹⁷ of data showed that naloxone promoted significant neurologic recovery in subjects with incomplete injuries, it was originally touted as lacking efficacy. Furthermore, neu-

rologic recovery with tirilazad was found to be comparable with that of MPSS for 24 hrs.

Two level II therapeutic trials were noted between 1999 and 2004. Monosialotetrahexosylganglioside GM1 sodium salt (Sygen)¹⁸ is a naturally occurring compound that has previously been shown to have neuroprotective and regenerative effects. Subjects who participated in the multicenter clinical trial of Sygen all received MPSS as per the NASCIS 2 protocol, followed by either a 300-mg loading dose of Sygen followed by 100 mg/day for 56 days, a 600-mg loading dose of Sygen followed by 300 mg/day for 56 days, or placebo for a loading dose and subsequently for 56 days. A number of analyses were conducted. Significance was not achieved with the predefined primary efficacy analyses; however, various subgroups of subjects such as those with American Spinal Injury Association (ASIA) B injuries and those who did not have spinal surgery seemed to have earlier and greater recovery with Sygen. In a smaller, placebo-controlled study,¹⁹ the use of nimodipine, a calcium channel blocker, was compared with MPSS alone, MPSS combined with nimodipine, and placebo alone. Although this study may not have been sufficiently powered to achieve significance, no differences in neurologic recovery were noted between the groups at 1 yr postinjury.

Conclusion

Few clinical trials have targeted improved neurologic recovery post-SCI. It is clear that further research should and will continue to be performed to evaluate neuroprotective strategies post-SCI.

Spinal Surgery

Surgical stabilization of acute spinal trauma is common; however, there is little agreement with respect to the usefulness of surgery in promoting neurologic recovery post-SCI. The exact timing of "early" *vs.* "late" surgery has not been defined, and the logic behind the choice of different times in various studies varies. Some authors define early surgery as up to 8 hrs postinjury, whereas others have used a 72-hr window. Although a number of meta-analyses have been performed evaluating various issues related to surgery, the studies that have been performed are primarily level III, with only one prospective, randomized level II study performed before our time frame and eight prospective, nonrandomized case series. All other studies were retrospective case series. Thus, the number of recommendations that can be made are small. There are level II data to show that surgery for acute SCIs can safely be performed. There are also level III data that support urgent decompression with bilateral facet dislocation or in incomplete SCI with a neurologic deterioration. Aside from

these subcategories, however, the data only support the use of surgical decompression and stabilization as a practice option.

Conclusion

In light of the large number of persons with SCIs who undergo surgery, this is an area in which future prospective, randomized studies are definitely warranted. These should determine whether there is a benefit of early or late surgery, when those time frames are, and whether surgery does promote neurologic recovery.

Neurologic Recovery

It is important to be able to prognosticate the recovery of motor and sensory function in persons with SCIs so as to provide accurate information to patients and their families, to guide the patient's rehabilitation, and to determine if new treatment methods are effective. In addition, it is important to be able to prognosticate the recovery of voluntary bladder and bowel function and other autonomic functions. This area has been considered so important that clinical practice guidelines for Outcomes Following Traumatic Spinal Cord Injury were published in 1999.²⁰ However, as can be seen from the guidelines, the scientific evidence regarding prognosis for recovery of neurologic function post-SCI was primarily level V. Moreover, research in this area has primarily focused on the recovery of motor and sensory function and generally avoided the recovery of autonomic function. Multiple studies have shown that the prognosis for neurologic recovery after SCI is best predicted by the clinical neurologic examination using the International Standards for Neurological and Functional Classification of Spinal Cord Injury Patients.²¹ The examination at 72 hrs postinjury has been determined to be superior to that at 24 hrs postinjury for determining recovery post-SCI.^{22,23} Those subjects with incomplete injuries at the time of examination are well known to have a better prognosis for neurologic recovery than those with complete injuries.

Using the American Academy of Neurology criteria for rating of prognostic articles, we were able to find six level III studies and one level II study published between 1999 and 2004. Successive grades in both the Frankel and the ASIA impairment scale (AIS) have better prognosis for functional motor recovery.²⁴ A prospective multicenter study recently documented that subjects with incomplete injuries demonstrated a greater likelihood of improvement of neurologic function by one level than those subjects with complete injuries.²⁵ In another smaller prospective study, patients free from cervical fractures with central cord syndromes all demonstrated neurologic improve-

ment, and the absence of magnetic resonance imaging findings of abnormal signal intensity in the spinal cord was considered a positive predictor for neurologic recovery.²⁶ One issue that remains unresolved with regard to documenting the remaining motor control of subjects with motor complete paralysis post-SCI is that of the subclinical discomplete²⁷ syndrome used to describe persons who are clinically paralyzed but show subclinical evidence of translesional motor connections. In a recent study²⁸ of 67 subjects with chronic AISA A or B SCI, surface electromyographic evidence of this discomplete state was found in 68% of AISA A subjects and 58% of AISA B subjects. How this finding relates to the potential for recovery induced by new therapies in subjects with acute and chronic SCI needs to be determined.

Other recent studies that have looked at neurologic recovery have been limited in that they have retrospectively reviewed longitudinal data collected as part of the Model Systems or for other studies. These level III studies have begun to look at the late effects of SCI²⁹ and the impact of sex³⁰ and etiology²⁴ on neurologic recovery, whereas other studies have considered the potential for marked recovery at various levels of injury¹⁸ and the reasons for neurologic deterioration³¹ postinjury.

Conclusion

Based on the current literature, further prospective studies of the natural history of neurologic recovery for both motor and sensory function post-SCI and how these variables affect recovery are necessary. In addition, studies should address recovery of autonomic function. This information will be particularly relevant to effectively plan for clinical trials in SCI.

Functional Potential

Determining the potential functional outcome of an individual post-SCI is an important part of the rehabilitation process. Functional outcomes are traditionally determined based on the level of SCI in conjunction with a review of the age, medical status, motivation, and family support of the patient. A number of tables have been developed to document the optimal functional potential of persons with SCIs. These have recently been reviewed in conjunction with an evidenced-based review of the literature on functional potential after SCI performed by the Spinal Cord Medicine Consortium.²⁰ Unfortunately reviews of the literature conducted by the Consortium reveal that the majority of studies are retrospective rather than prospective. In addition, potential recovery is generally determined by the FIMTM motor subscale, which is not optimal to assess the progress of SCI patients.

Based on our criteria, we found eight level III studies addressing this topic. The issues of sex,³² age,^{33–36} and cause of injury^{37,38} and their impact on function have been studied. Magnetic resonance imaging³⁹ has also been used to prognosticate functional recovery post-SCI. Overall, these studies have not found a significant difference between groups with respect to function; however, there is some suggestion that with older age there is a negative impact on functional potential.

Conclusion

Further studies should be performed to document the optimal functional potential of persons with SCIs. Moreover, more sensitive outcome measures than the FIMTM instrument should be used to describe changes in functional status, and further research should be pursued to determine the impact of findings in using imaging on functional recovery.

TREATMENT OF MEDICAL CONSEQUENCES AND COMPLICATIONS

A large part of rehabilitation from acute SCI involves the management of the effects of the injury on various organ systems. These include the optimal management of the injury on bladder and bowel function, optimizing respiratory function, preventing pressure ulcers and deep vein thrombosis, among other effects. In addition to managing these problems, patients and their families are educated about the impact of their SCI on organ system functioning and how to manage potential problems.

Respiratory Function

Persons with SCIs at the level of T12 and above have a decrease in respiratory function. Impaired muscular function results in a decline in vital capacity and the ability to cough. Consequently, respiratory complications such as pneumonia are a major source of morbidity, and respiratory conditions are the number one cause of death post-SCI.⁴⁰ Despite this fact, only one level II and one level III therapeutic study and one level III prognostic study were found during our designated time period.

Traditional methods to improve pulmonary function after SCI include respiratory strengthening exercises and assistive coughing. These methods have not, however, been proven as efficacious in improving vital capacity above and beyond the natural recovery that occurs after SCI. Liaw et al.⁴¹ documented the benefits of resistive inspiratory muscle training in patients with complete SCI; however, their sample size was limited and it is

uncertain whether testing of pulmonary function was done in a blinded fashion.

Ventilator Dependency

Secondary complications associated with the decline in respiratory capacity post-SCI include the development of ventilator dependency, especially in patients at the level of C4 and above and in those persons with vital capacities of <1 liter. Weaning of persons with SCIs from ventilators in SCI units is generally done with progressive T-piece trials and with rest on the ventilator between trials using assist-control mode. However, the optimal ventilator tidal volume settings to facilitate weaning of patients from ventilatory support has not been prospectively studied. A retrospective evaluation⁴² demonstrated that persons with SCI ventilated with tidal volumes of ≥ 20 ml/kg body weight were able to wean from the ventilator substantially quicker than those who were ventilated with tidal volumes <20 ml/kg body weight. Furthermore, those patients who were weaned using large tidal volumes had substantially less atelectasis than those weaned using low tidal volumes. Nevertheless, in the acute hospital setting, weaning is generally performed using pressure-support ventilation, and when on assist-control mode, patients are ventilated using more traditional tidal volumes of 5–10 ml/kg.

Pneumonia

Along with atelectasis, pneumonia is also a frequent problem for persons with both acute and chronic SCI. Treatment of community-acquired pneumonia was recently reviewed in a level III prognostic study in the Veterans Administration Healthcare system,⁴⁰ with the finding that most SCI patients with pneumonia are admitted for treatment and that the short-term mortality is comparable with community-acquired pneumonia in the general population.

Conclusion

Although pulmonary consequences are the primary cause of death post-SCI, there is a paucity of current research. A multicenter, clinical trial to determine the optimal method to wean ventilator-dependent SCI patients is a priority for future research, as are studies addressing effective ways to improve overall pulmonary function and to prevent respiratory complications.

Neurogenic Bowel Dysfunction

General Care

Neurogenic bowel can result in incontinence, constipation, and a general decrease in quality of life post-SCI. A recent comparison of 467 SCI and

668 able-bodied age- and sex-matched control subjects revealed fecal incontinence was significantly higher in SCI persons than controls and that among SCI subjects was greater in those with complete *vs.* incomplete injuries.⁴³ Guidelines for neurogenic bowel management in adults with SCI⁴⁴ were developed in 1998. Unfortunately, there was an overall lack of high-level evidence in most topic areas; thus, the guidelines developed were primarily based on expert consensus. During our time frame, we found one level II prognostic study related to neurogenic bowel dysfunction, with three level III prognostic studies (of which two could also be considered diagnostic) and one level IV prognostic study related to gallstones.

It is generally accepted practice that SCI patients should be trained in a bowel program that takes into account the type of injury (upper motor neuron *vs.* lower motor neuron) affecting their bowel, their symptoms, and whether they have physical capability and support to perform a bowel program. Based on the patient's neurologic injury, a bowel management program is developed using an appropriate chemical or mechanical rectal stimulant for those with reflexive bowels and an appropriate manual evacuation technique with areflexive bowels. A sufficient fluid intake is recommended, along with proper diet and regular performance of the bowel program. Stool softeners and fiber are used when appropriate.

Gastrointestinal transit is delayed in subjects with SCI as compared with able-bodied control subjects.⁴⁵ Furthermore, the specific locations of segmental decreases in colonic transit time in respect to level of neurologic injury have recently been documented using radiopaque markers and radiographic examinations.⁴⁶ Cisapride, a prokinetic agent previously used to improve gastrointestinal transit time in SCI, has been discontinued.⁴⁷ Positive pilot data were also obtained that demonstrated potential efficacy and tolerability of prucalopride to improve transit time in patients with SCI.⁴⁸ Unfortunately, this drug was also discontinued. Thus, further testing of the newest prokinetic drug, tegaserod,⁴⁹ is warranted via a multicenter double-blind, placebo-controlled study.

Complications

Little is known about the incidence of acute abdomen in SCI and the optimal treatment of this unfortunate occurrence. No studies were found in the past 5 yrs on this topic or with regard to management of gastric ulcers after SCI. Little is also known about the effects of colostomy for long-term impact of neurogenic bowel and whether the impact on timing of bowel care and quality of life in persons with SCI justifies this procedure. Consortium Guidelines indicate a decision for colostomy

or ileostomy should be based on “the results of specialized screening procedures and the individual’s expectation” but that “a permanent stoma is the best option.” Although a recent retrospective study revealed a decrease in bowel care time from 10.3 hrs/wk to 1.9 hrs with a colostomy,⁵⁰ prospective controlled data are still necessary to determine the risk and benefits of this procedure.

Gallstones

Gallstones have been known to occur in almost one third of patients with SCIs,^{51–53} which is a 3- to 6-time greater risk than that seen in age- and sex-matched controls.⁵⁴ Whether the presentation of symptomatic gallstones is similar to that of the able-bodied person is uncertain, as is the recommended treatment of asymptomatic stones. Moonka et al.^{52,54,55} performed a number of level III studies in the veteran population using routine abdominal screening ultrasonography. They noted a prevalence of 31% for gallstones, with an annual rate of cholecystectomy of 6.3%. SCI subjects with stones were noted to have typical right upper quadrant and epigastric pain similar to able-bodied controls, and vague abdominal symptoms were not indicative of gallstones.⁵⁶ Regardless of level of injury, the presentation of gallbladder disease was not more advanced than that in neurologically intact patients. The weakness of these studies is that they were retrospective and performed in a veteran population; however, the findings regarding symptoms and treatment for gallbladder disease are probably relevant to a nonveteran SCI population.

Conclusion

Few prospective studies are available to support practice patterns for management of neurogenic bowel dysfunction. Further research should include double-blind, placebo-controlled studies of methods to optimize bowel function. In addition, optimal methods for diagnosing and treating complications such as gastric ulcers and the acute abdomen should be researched.

Neurogenic Bladder

General Care

A neurogenic bladder is an anticipated consequence of most SCIs. Over the past few years, the treatment of neurogenic bladder has improved such that the mortality from renal disease post-SCI has been exceeded by that of pulmonary disease. The management of a neurogenic bladder involves the use of physical examination and urodynamic testing to determine the type of injury affecting the bladder and the pressure profiles present in the bladder. Then patients are “bladder trained,” a pro-

cess which describes the physical and medical manipulation of the pressure profiles in the bladder to facilitate storage of urine or increased voiding.

In our time frame, we located eight level II, three level III, and four level IV notable studies related to neurogenic bladder dysfunction. Avoidance of high detrusor pressures is one of the caveats in maintenance of bladder function post-SCI. Level II evidence supports the use of tamsulosin⁵⁷ as a means to decrease maximal urethral pressure and the use of intravesical capsaicin and resiniferatoxin⁵⁸ for the treatment of detrusor hyperreflexia. Level IV evidence supports the efficacy of botulinum toxin A⁵⁹ for the treatment of detrusor hyperreflexia and the use of an endoprosthesis for detrusor sphincter dyssynergia.⁶⁰ It is notable that these level IV studies used pre-urodynamic and posturodynamic outcome measures rather than control groups due to the invasive nature of these procedures, and this underscores the difficulties of performing clinical research.

Regardless of the prescribed method of bladder emptying that individuals with SCI use at discharge from rehabilitation, they often change their bladder-emptying method over time. In a retrospective study of 235 SCI individuals at a mean of 24.1 yrs postinjury,⁶¹ the use of clean intermittent catheterization increased from 22% at initial discharge to 36%, suprapubic tapping decreased from 57% to 31%, and the use of Crede increased from 5% to 19%. A higher portion of patients continued with their bladder management method when they were using clean intermittent catheterization.

Urinary Tract Infections

Urinary tract infections (UTIs) are a common complication post-SCI, and much research has targeted their prevention. Level II evidence⁶² supports the use of an educational program to decrease the number of bacterial colony counts in persons with SCI. Two level II studies^{63,64} showed no benefit of the use of cranberry extract in reducing UTIs. Level II evidence supports the use of ofloxacin over trimethoprim sulfamethoxazole⁶⁵ in hospitalized persons with SCI and symptomatic UTI. Interestingly, level III evidence supports the use of bacterial interference in preventing UTI^{66,67} and noted this could develop into a safe, effective means of preventing symptomatic UTIs.

Other Complications

Other complications of neurogenic bladder include the development of renal stones and a higher incidence of bladder cancer post-SCI. The incidence of renal stones was recently noted to be unchanged over the past 25 yrs⁶⁸ and found to be approximately 3% in a new cohort of subjects with SCI, with an incidence going up to approximately

7% within 10 yrs of injury. Furthermore, once a person has a kidney stone, level II evidence documents that 34% develop a second stone within 5 yrs.⁶⁹ Bladder cancer was recently shown to have a standardized morbidity ratio in the SCI population of 25.4 as compared with the general population.⁷⁰ Level III⁷⁰ and level IV⁷¹ evidence supports the sole use of indwelling catheters as a risk factor for bladder cancer. The 5-yr mortality from bladder cancer in SCI was 38%.⁷⁰ To more effectively screen for bladder cancer, a new method to detect bladder cancer through analysis of urine for BLCA-4 was tested. A prospective study⁷² established a normal cutoff point for BLCA in the urine and showed that 19% of SCI patients had a higher level than normal, despite the lack of bladder cancer. Thus, it will be important to continue to follow these patients to see if this could be an early marker for the development of bladder cancer.

Conclusion

A number of studies have recently addressed prophylaxis against symptomatic UTI after SCI. Prospective controlled studies evaluating the efficacy of new therapies to manage neurogenic bladder are warranted. Moreover, determination of the natural history of neurogenic bladder function as it relates to remaining neurologic function postinjury is needed.

Neurogenic Sexual Function Natural History

Spinal cord injuries result in predictable effects on sexual responses. This is an area in which recent research has been relatively plentiful, and overall we noted two level I studies, eight level II, and one level III. There is level II evidence that a loss of psychogenic erection correlates with sympathetic denervation, and loss of somatic and parasympathetic reflex activity correlate with the loss of reflex erection.⁷³ In women, there is level I evidence that the ability to achieve psychogenic lubrication is related to the degree of preservation of sensation in the T11-L2 dermatomes and that reflex lubrication is maintained in women with upper motor neuron injury affecting the sacral segments.⁷⁴ There is also level II evidence that the ability to achieve orgasm is maintained in women post-SCI, provided they have an intact sacral reflex arc.

Treatment of Sexual Dysfunction

Treatment of sexual dysfunction has been revolutionized since the development of oral therapies for erectile dysfunction. Previous therapies included penile prostheses, intracavernosal injection of vasoactive agents,⁷⁵ the use of intraurethral al-

prostadil⁷⁶ and the use of vacuum erection devices. These have largely been replaced by pharmacologic therapies as first-line agents to treat erectile dysfunction after SCI. Level I⁷⁷ and level II⁷⁸ evidence exist to document the benefits of the use of sildenafil to treat erectile dysfunction post-SCI. Level II evidence has also noted slightly lowered blood pressure in men with SCIs above T6 after taking sildenafil.^{79,80} The use of new phosphodiesterase inhibitor tadalafil may have some advantages over sildenafil,⁸¹ primarily due to a longer half-life. Treatment of female sexual dysfunction after SCI has also recently been explored. Three level II studies were recently published. One⁸² revealed beneficial effects of sildenafil on subjective sexual arousal along with an acceptable safety profile, and another⁸³ documented potentially beneficial effects of anxiety-provoking stimulation. Finally, potentially beneficial effects of false-feedback on sexual arousal were noted in the subset of women with significantly impaired sympathetic input to their genitals.⁸⁴

Male Fertility

The study of reproductive health after SCI has focused on male infertility and the elicitation of ejaculation, primarily through the use of electroejaculation⁸⁵ or penile vibratory stimulation.⁸⁶ Predictors of successful ejaculation through vibratory stimulation have been studied,⁸⁷ and 80% of subjects who had both a bulbocavernosus reflex and hip flexor response ejaculated with penile vibratory stimulation, whereas only 8% of patients negative for both reflexes ejaculated. Although optimal parameters for vibration have been suggested,⁸⁸ it has been noted that repeated ejaculation in men with SCI does not change typically poor semen quality.⁸⁹

Female Reproductive Health

Women's reproductive health has received little attention as compared with that of men. A multicenter outpatient questionnaire study was recently performed⁹⁰ to document the unique reproductive health conditions, complications, and behaviors of women with SCIs. In 472 women, similar gynecologic problems were found preinjury and postinjury, with the exception of increased UTIs and vaginal yeast infections postinjury. Women with SCI were less likely to have routine mammograms, and although there were similar numbers of hysterectomies both preinjury and postinjury, the reasons were significantly different. Almost 14% of women with SCI became pregnant postinjury. Menopause after injury was reported by 14.6% of women.

Conclusion

Further research pertaining to neurogenic sexual function should address better methods to treat other components of male sexual dysfunction such as orgasmic dysfunction as opposed to the previous isolated focus on return of erectile function. Moreover, research should continue to target optimizing the sexual and reproductive issues of women with SCIs. Dissolution of barriers to mammography and routine gynecologic care should be pursued. Finally, any therapeutic effects associated with routine sexual activity should be explored.

Pressure Ulcers

Pressure ulcers are a serious and common consequence of SCI. The prevalence of pressure sores has been described as 33% in a community-based sample,⁹¹ and the 1998 incidence was noted as 34% in patients admitted to model SCI systems within 1 day of injury when one considered the time period through their discharge from rehabilitation.⁹² Having a complete cervical injury has been noted to be associated with a higher prevalence of pressure sores,⁹³ and age, injury duration, sex, marital status, education, activity, mobility, and comorbidities have been associated with increasing risk from pressure sores; however, findings are inconsistent. Pressure ulcers were noted to least likely occur in persons with normal weight who have returned to their work and family roles and who do not have a history of substance or tobacco use, suicide, or incarceration.⁹⁴

Pressure ulcer prevention and treatment⁹⁵ has recently been reviewed via the Consortium for Spinal Cord Medicine Clinical Practice Guidelines. Unfortunately, despite the fact that the cost of pressure ulcers has been estimated as \$1.335 billion per year,⁹⁶ there were no level I, II, or III studies related to pressure sores that met our review criteria; thus, this discussion is primarily based on the clinical practice guidelines. Pressure sores are graded from I to IV based on their severity. A standard in treating pressure ulcers is to determine their location and appropriately relieve pressure from the area. Other standards of care include instituting pressure relief as soon as emergency medical care and spinal stabilization allow, turning every 2 hrs in bed during the beginning of rehabilitation, avoiding direct positioning on the trochanters while in bed, and performing weight shifts while in the wheelchair. Shearing forces should be minimized, and appropriate seating and cushioning should be provided. When manual pressure relief is not possible, a power weight-shift system is appropriate. Patient education is generally considered paramount in emphasizing the need to prevent pressure sore development. There

is level I evidence⁹⁷ for the use of air-suspension beds to prevent pressure ulcers; however, the use of these beds is not practical for chronic SCI.

Treatment of pressure ulcers should be instituted as soon as possible. To determine appropriate therapies and document their efficacy, the existing pressure ulcer must be described in detail, including the anatomical location, size, stage, whether there is necrosis, exudates/odor, undermining, infection, the appearance of the wound margins, and surrounding tissue.⁹¹ The wound should be removed from the offending pressure source, and appropriate debridement and cleansing should be ordered. Dressings should be used that keep the ulcer bed moist and the surrounding tissue dry. Wound cavities should be packed enough to be effective but not too tightly so as to prevent granulation at the base of the wound.⁹³ A variety of dressings are available for treatment of wounds and should be selected based on the clinical appearance of the wound. Electrical stimulation is the sole therapy for which level I evidence is present to support its efficacy in promoting closure of stage III or IV wounds in SCI in conjunction with standard wound care,^{98–100} and surgical closure is an appropriate option in patients with nonhealing pressure ulcers.

Conclusion

As pressure ulcers are common, costly, but often easy-to-treat, the lack of overall level I evidence to support treatment paradigms is surprising. Multicenter clinical trials to determine the most effective means to treat pressure sores after SCI are clearly overdue.

Thrombophlebitis

Thrombophlebitis is a common complication of SCI that occurs in conjunction with the paralysis of the injury. Thrombophlebitis is the cause of pulmonary embolism after SCI and can also be a cause of untimely death. Due to this potentially lethal complication, the second edition of Clinical Practice Guidelines for thromboembolism prevention was published in 1999.¹⁰¹ The clinical practice guidelines noted nine studies with a total of 419 subjects that documented a 40% rate of objectively diagnosed deep vein thrombosis (DVT) in patients with acute SCI. Based on consensus opinion and the literature at the time, the guidelines recommended first determining the level of risk for DVT based on whether SCI patients are motor complete or incomplete or motor complete with other risks such as lower limb fracture, previous thrombosis, cancer, heart failure, obesity, or age of >70 yrs. The guidelines recommended the use of compression hose or pneumatic devices for the first 2 wks

after SCI, with anticoagulant prophylaxis with either low molecular weight heparin or adjusted-dose unfractionated heparin within 72 hrs after SCI, provided there are no contraindications. Recommendations are that prophylaxis be continued for 8 wks in patients with uncomplicated motor complete injury and, in patients with complications, that prophylaxis continue for 12 wks or until rehabilitation discharge.

A recent retrospective review of 243 persons with acute SCI,¹⁰² all of whom had objective testing, revealed 51 subjects had DVT and eight died. The highest likelihood for DVT was in persons >35 yrs of age who had cancer. Obesity was also a risk factor. Women between 36 and 58 without cancer and men with flaccid paralysis were also noted to be at high risk. In our time period, there were two level II and one level three therapeutic studies addressing prophylaxis against DVT. Two studies^{103,104} of prophylaxis compared the use of unfractionated heparin or enoxaparin combined with pneumatic compression during the first 2 wks postinjury and alone for the next 6 wks. No significant differences were found in the efficacy or the complications associated with either treatment. Another study looked at the incidence of thromboembolism in acute SCI patients *vs.* other trauma patients and found that the subgroup of patients with SCI and long bone fractures were significantly more likely to incur a DVT. They noted that routine use of prophylactic Greenfield filters does not seem warranted as a preventative measure against pulmonary thromboembolism post-SCI; however, those patients with long bone fractures and SCI, those with DVT formation despite prophylactic anticoagulation, and patients with contraindications to anticoagulation may be appropriate candidates for prophylactic caval filters.¹⁰⁵

Conclusion

Future prospective studies should be designed to evaluate the adherence to and efficacy of the clinical practice guidelines in preventing DVT. Optimal methods for treatment of DVT should be determined.

Musculoskeletal Effects of SCI Heterotopic Ossification

Heterotopic ossification is a frequent complication post-SCI that is generally diagnosed between 1 and 6 mos postinjury but most commonly at 2 mos postinjury.¹⁰⁶ Two level II therapeutic studies showed successful prevention of heterotopic ossification through the use of indomethacin¹⁰⁷ and rofecoxib.¹⁰⁸ Both studies showed a significant decrease in the development of heterotopic ossification post-acute SCI; however, although sta-

tistical significance was achieved, the studies were limited by a small *n* and the use of the clinical presence of heterotopic ossification as a primary screening measure. Moreover, rofecoxib has recently been taken off of the market.

Once heterotopic ossification forms, the usual treatment involves the use of etidronate sodium and/or indomethacin and can also include radiotherapy and surgical resection.¹⁰⁹ However, surgical resection involves significant risks and thus is generally only undertaken in the presence of complications such as joint immobility or pressure sores as a result of the ossification.

Osteoporosis

Osteoporosis is also a known consequence of SCI. We were able to find two level I, three level II, and three level III prognostic studies related to osteoporosis but no intervention studies. A level II study¹¹⁰ recently documented a significant decrease in radial bone density at 6 mos postinjury and both radial and ulnar bone density by 12 mos post-SCI in tetraplegics but not paraplegics. Furthermore, tibial trabecular bone mineral density was significantly decreased in all SCI subjects at 6 mos and 1 yr post-SCI. A recent level I analysis¹¹¹ showed that completeness of injury followed by body mass index were the most likely variables to predict bone loss at the knee leading to pathologic fractures. Moreover, the patterns of regional bone loss in women¹¹² include the greatest loss of density at the knee, followed by the hip, and complete preservation of bone mineral density at the spine.

Conclusion

In conclusion, most research pertaining to the musculoskeletal consequences of SCI has focused on the treatment of heterotopic ossification and determining prognostic factors related to the occurrence of osteoporosis. Although significant information is available relating to prognosis of osteoporosis, double-blind, placebo-controlled trials are warranted to study prevention and treatment of osteoporosis after SCI. Furthermore, further studies are needed in relation to the occurrence of musculoskeletal consequences of SCI such as shoulder pain, carpal tunnel syndrome, and other overuse syndromes.

Metabolic Consequences and Body Composition Metabolic Consequences

A number of metabolic consequences have been identified after SCI. As compared with the general population, persons with SCI have been shown to be more likely to have oral carbohydrate intolerance,¹¹³ insulin resistance,¹¹⁴ elevated low-

density lipoprotein cholesterol,¹¹⁵ and reduced high-density lipoprotein cholesterol.¹¹⁶ We were able to find three level I and two level II prognostic studies related to metabolic consequences along with one level III intervention study. Level I evidence documents that SCI individuals with the greatest neurologic deficits have an increased risk of developing disorders of carbohydrate metabolism, men with SCI are more insulin resistant than women, and glucose tolerance seems to be independent of the effects of ethnicity.¹¹⁷ Level I evidence indicates that in white and Latino men, but not in women or African Americans, immobilization subsequent to SCI seems associated with lower high-density lipoprotein cholesterol levels than in controls.¹¹⁸ Level III evidence supports the use of a dietary intervention to reduce total cholesterol and low-density lipoprotein cholesterol in persons with SCI and elevated cholesterol.¹¹⁹

Body Composition

Body composition in individuals with SCIs differs from that of the able-bodied. There is level I¹²⁰ evidence that men with SCI are $13\% \pm 1\%$ fatter per unit of body mass index (kg/m^2) than an age-, height-, and ethnicity-matched able-bodied control population and level II¹²¹ evidence documenting that as compared with able-bodied men with similar body mass indexes, body fat percentage was 9.4% greater in the SCI group.

Conclusion

In light of the triad of altered body composition, glucose metabolism, and lipoprotein profiles, it is not surprising that there is a significant risk of cardiovascular disease in persons with SCIs. Over the time period 1973–1998, out of 1,543 deaths occurring in persons followed through the SCI Model Systems program, 18.8% of deaths were attributable to cardiac causes.¹²² Further research is warranted to determine effective interventions to treat the negative metabolic and body composition effects of SCI. In addition, the efficacy of wellness programs in improving metabolic profiles and decreasing cardiovascular morbidity and mortality should be studied.

Spasticity

Spasticity is a well-known consequence of upper motor neuron injury and can be as disabling to an individual with SCI as the injury itself. The first-line treatment of spasticity has traditionally been with oral therapies including baclofen, tizanidine, Dantrium, and Valium. However, each of these medications has known side effects and limited efficacy; thus, more invasive treatments have begun to be used, including botulinum toxin injections

and neurosurgical procedures. Peripheral neurosurgical ablative procedures include rhizotomy and peripheral neurectomy. Central ablative procedures include cordectomy, myelotomy, and stereotactic procedures. Nonablative procedures include peripheral nerve or motor point blocks, the implantation of cerebellar or spinal stimulators, and the implantation of subdural catheters for relief of spasticity.¹²³ Despite the significant functional problems associated with spasticity and the expense and difficulty with treatments, we were unable to find any level I or II intervention studies during our time frame.

Evaluation of Spasticity

Part of the difficulty in evaluating treatments for spasticity is the reality that spasticity changes with the time of day, whether a person has received range-of-motion exercise, whether the person has other acute medical problems or other conditions, and the weather. The Ashworth scale,¹²⁴ originally intended for use in multiple sclerosis, is part of a recommended triad for evaluations including spasm frequency and self-report. The Ashworth scale is reported to be used most commonly clinically post-SCI¹²⁵; however, more exact outcome measures are needed. A recent study evaluating spasticity post-SCI included visual analog scales¹²⁶ and the Modified Ashworth Scale¹²⁷ as methods of documentation of spasticity. The latter study also documented significant fluctuation of the spasticity of cervical SCI subjects during the day as compared with thoracic-injured subjects. Findings confirmed that passive movement immediately decreased spasticity ratings in thoracic patients. Visual analog scale ratings correlated significantly with Modified Ashworth Scale ratings.

Other methods to document spasticity have included more quantitative analysis such as the use of isokinetic dynamometry¹²⁸ using maximum torque values and the sum of torque amplitudes. A positive correlation was noted between these values and Ashworth scores. However, even with such quantitative analysis, variability with different positions has been noted,¹²⁹ underscoring the importance of exact descriptions of test positions and procedures in research reports. Brain motor control assessment¹²⁵ using surface electromyography recordings has also been used to document degree of spasticity. In a study of 97 subjects with SCI¹²⁵ and varying levels of motor dysfunction, the analyses of averages of surface electromyography for a limb were able to document a significant difference for scores between those with Ashworth of 0 *vs.* 2 and 3 and 1 *vs.* 2 and 3 but not between 0 and 1.

Conclusion

Spasticity is an area in which our ability to assess treatment in a laboratory may be excellent, but our clinical outcome measures are still in their infancy. In light of this fact, new therapies should have their efficacy documented in a laboratory-based setting before advancement into clinical trials. The development of a validated outcome measure to assess spasticity via self-report and clinical examination is necessary not only for the development of more effective therapies to treat spasticity but also to assess the effects on spasticity of therapies for promoting neurologic recovery.

Pain

There is a substantial literature that has accumulated over the years documenting a significant prevalence of pain in persons with SCI.¹³⁰ Prevalence figures range from 65% to 79% for community-based samples,^{131,132,133} with one third of those reporting severe pain. Age, sex, completeness of lesion, pathogenesis, and duration of injury have been examined cross-sectionally as risk factors for the development of chronic pain with inconsistent results across studies.^{134,135} Chronic pain has been identified as having a negative influence on quality of life.^{136–138} Although considerable attention has been given to neuropathic pain at and below the level of injury, increasing attention is being paid to upper limb pain and to ways to minimize it.^{139,140}

One of the shortcomings of pain research is the failure to distinguish subtypes of SCI pain. Recently there have been a number of attempts to remedy this problem by the development of classification schemes¹³⁰; however, there is no consensus. Reports have just recently started appearing in the literature demonstrating different time courses, levels of severity, implications for quality of life, and underlying pathology for different subtypes of SCI pain.^{141–143} The International Association for the Study of Pain has recently proposed a classification scheme in the hope that it will be adopted world-wide,¹⁴⁴ but this has not occurred to date. Performance of reliability and validity studies for pain classification schemes, a necessary precursor for widespread acceptance, is imperative so that the subtypes of pain can be evaluated and treated appropriately. Our search, however, revealed no level I studies and only one level II study¹⁴⁵ supporting the validity of proposed SCI pain diagnostic schemes.

Several studies of prognostic outcomes derived from longitudinal sampling were recently published. Persons with recent-onset SCI were recently studied for 5 yrs in a level III study.¹³⁸ Neuropathic pain tended to persist over time, whereas musculoskeletal and visceral pain did not

predict future pain. In a level II study using case-control methodology,¹⁴⁶ persons developing pain interference from one year to the next reported decreased life satisfaction and decreased physical and mental health, whereas those reporting decreased pain interference over that same period reported improvements in those measures. In another level II¹⁴² study, demographic, injury-related, and radiographic variables were predictive of increased shoulder pain and range-of-motion problems over a 3-yr period. Furthermore, these problems were related to functional limitations, disability, and perceived health.

Much of the literature concerning treatment of SCI pain has been characterized by clinical reports and uncontrolled case series.¹⁴⁷ However, level I and II controlled trials for particular subtypes of pain have begun to emerge. Gabapentin has been proven efficacious for the treatment of neuropathic pain in two small, randomized, controlled trials (level I and II, respectively).^{148,149} Intravenous ketamine was shown to significantly reduce neuropathic pain in a level II trial,¹⁵⁰ as did the intrathecal combination of morphine and clonidine (level II),¹⁵² and oral lamotrigine (level II).¹⁵³ Amitriptyline did not prove efficacious for pain across all of the subtypes of SCI pain combined nor did the authors find a differential effect for the specific subtypes of pain for which their samples were large enough to examine (level I).¹⁵⁴ Exercise has been shown to reduce pain in a small level II, randomized, controlled trial in which subtypes of pain were not reported¹⁵⁵ and in a level II trial of stretching and exercise for shoulder pain in long-term wheelchair users.¹⁵¹ Controlled trials of implanted spinal cord stimulation are lacking, and reported results from clinical series are not encouraging.¹⁵⁷ Dorsal root entry zone microsurgical ablative approaches have been reported in clinical reports as highly successful, primarily for selected cases of neuropathic pain.^{158,159} Cognitive-behavioral interventions have not been subjected to controlled trials for their effectiveness in this population.¹⁶⁰ A large number of other modalities are in use,¹⁶¹ and alternative medicine approaches such as acupuncture have been described in clinical series as helpful.¹⁶² A level II study of the treatment of shoulder pain demonstrated the effectiveness of both acupuncture and Trager treatments.¹⁵⁶

Conclusion

Given the significant negative impact of pain on quality of life, continued efforts to demonstrate efficacious treatments are needed. Specifically, research is needed to validate and support the adoption of a standardized classification scheme with accompanying diagnostic procedures and criteria. Level I multicenter, randomized trials of interven-

tions are also needed to reduce the severity and impact of the different subtypes of SCI pain. Such trials could include both primary prevention and treatment of acute and chronic pain.

Outcome Measures

Outcome measures to address function post-SCI are in a state of dynamic change. Despite the fact that they do not fit into evidence guidelines, we thought it important to review the recent literature and document ongoing initiatives. The International Standards for the Classification of SCI remain the gold standard for the documentation of neurologic function post-SCI.¹⁶³ However, newer techniques such as quantitative sensory testing¹⁶⁴ have been assessed to see whether they could be used as an improvement over the testing of sensory function via the International Standards. The voluntary response index was also recently recommended¹⁶⁵ as a means for objective interpretation of surface electromyographic responses obtained after SCI. The voluntary response index consists of one numeric value derived from the magnitude of muscle activity recorded during a specific voluntary task and another numeric value (called the similarity index) from the surface electromyographic distribution across recorded areas during the same task. Although this instrument needs further validation, having a more sensitive method to quantify recovery of neurologic function would be of great benefit in order to develop methodology to study SCI.

The FIM™ motor score remains the most commonly used and studied scale to address the functional recovery of persons with SCIs. However, use of the FIM™ instrument in isolation is inadequate to describe function post-SCI. Therefore, the Walking Index for Spinal Cord Injury (WISCI)¹⁶⁶ and revision (WISCI II)¹⁶⁷ were recently developed as measures to quantify ambulatory function. To address other functional capacities, the Spinal Cord Independence Measure¹⁶⁸ was recently validated, and a short form of the Quadriplegia Index of Function¹⁶⁹ was developed. In addition to monitoring the recovery of motor and sensory function and functional recovery, documentation of the recovery of bladder, bowel, and sexual function is also important. However, there is a notable absence of a single scale to address autonomic function. A group involving the international collaboration of the International Spinal Cord Society and the American Spinal Injury Association is currently in the process of developing such a measure.

With regard to outcome measures used in psychosocial, epidemiologic, and technology-related SCI research, there is such a diversity of measures used that comparing outcomes across studies is problematic. Two unresolved issues characterize

this literature: whether SCI-specific as opposed to generic measures are needed and the extent to which generic measures are “biased” or lack validity when used with persons who have SCI. With well-validated generic measures, comparisons of outcomes can occur across different samples; however, some generic measures do not capture all relevant domains for persons with SCI. Moreover, specific item content can be irrelevant or misleading (e.g., sleep disturbance can reflect many medical phenomena other than depression). Ongoing efforts are directed toward developing SCI-specific measures of perceived needs,^{170,171} quality of life,¹⁷² and activity.¹⁷³ On the other hand, there are also ongoing efforts to validate the use of generic measures of various constructs with the SCI population in areas such as vocational rehabilitation intervention effectiveness,¹⁷⁴ environmental factors,¹⁷⁵ participation,¹⁷⁶ depression,¹⁷⁷ quality of life,^{178,179} and receptivity to assistive technology.^{180,181}

Conclusion

Research is needed to determine the most sensitive yet practical outcome measures to be used to document both remaining functional and neurological status post-SCI. Research on the psychosocial aspects of SCI would be enhanced by funded efforts to identify, develop, or validate a set of instruments across the most important domains that could be proposed as the standard set of instruments for researchers to adopt in this area.

REHABILITATION Exercise Therapies

Traditionally, management of the person with SCI involves the determination of the person's level of injury and functional capacity and then the generation of a problem list and prescription for therapies, with an overall goal of having the patient achieve his or her maximum functional potential. Based on this evaluation, various therapies including muscle strengthening, range of motion, transfer training, gait training, and activities of daily living training are pursued, generally through a team approach. The person is also prescribed and trained in the use of various pieces of equipment, including wheelchairs, transfer boards, and splints. Although the benefits of inpatient interdisciplinary rehabilitation after SCI are intuitively obvious, there have been few evidence-based analyses of the efficacies of specific exercise therapies. We were only able to find one level II study meeting our criteria; thus, we did not construct a separate table. This study compared the central cardiovascular responses of 20 complete thoracic SCI subjects to those of 20 able-bodied subjects during arm-crank exercise.¹⁸² Despite significantly greater values of

heart rate at rest and during matched absolute workloads compared with able-bodied subjects, cardiac output was lower in SCI subjects at both periods in time.

One area that has recently received significant attention is the use of body-weight-supported treadmill training. It is notable though that during our time period, none of the studies met levels I–III evidence criteria. This therapy has shown the ability to induce a locomotor pattern even in patients with complete paraplegia in conjunction with increases in leg extensor electromyographic responses.¹⁸³ Level IV evidence supports the use of body-weight-supported treadmill training combined with electrical stimulation in patients with acute¹⁸⁴ SCI and its use in subjects with incomplete SCI for ≥ 1 yr postinjury.¹⁸⁵ However, neither of these studies had control groups, and the final results of more recent multicenter studies were not yet published at the time of this review. More recently, a robotic orthosis has been used for performance of treadmill training¹⁸⁶ and has been made commercially available as the Lokomat¹⁸⁷ in the United States (Hokoma, Chicago, IL). Whether this device will prove cost-effective and beneficial as compared with conventional therapies targeting the improvement of gait post-SCI needs to be tested.

Conclusion

As the foundation of rehabilitation involves exercise, it is surprising that so little evidence is available to support its use. Substantial research is needed to document the benefits of exercise interventions post-SCI. Issues that also need to be determined are the optimal methods for strengthening muscles, range of motion, gait, activities of daily living, and transfer training. In addition, the relative merits of endurance *vs.* strength training should be determined. Due to the impact of body composition, age, concomitant medical problems, and our limited knowledge of the natural history of recovery post-SCI, this research will need to be performed through well-designed multicenter trials. Moreover, because these therapies are costly and have traditionally been billed for despite lack of proven efficacy, consensus should be sought on an ethical way to finance this research.

Electrical Stimulation

The use of electrical stimulation in persons with SCIs has been developed for both therapeutic and functional purposes. Although a significant amount of work has been published in this area, most of it has been in demonstration studies. Thus, traditional evidence levels often do not apply, and we have not constructed a table on this topic, as there was not any level I or II evidence. Therapeutic

uses for electrical stimulation after SCI have included its use for prevention of DVT,¹⁸⁸ prevention of pressure ulcers,¹⁸⁹ and for the exercise of muscles post-SCI.¹⁹⁰ Documented benefits of the use of electrical stimulation for exercise post-SCI include increases in strength in paralyzed muscles,¹⁹¹ increased muscle mass,¹⁹² and improved muscle endurance for lower limb muscles.¹⁹³ Increased high-density lipoprotein levels¹⁹⁴ and improved insulin sensitivity¹⁹⁵ have been documented through the use of a lower limb electrical stimulation exercise program. With chronic use, the use of electrical stimulation has also been shown to potentially have a positive effect on bone density.¹⁹⁶ A recent controlled study tested 14 SCI subjects and controls who underwent isokinetic stimulation of the left quadriceps 1 hr daily, 5 days/wk for 24 wks. Although the initial bone density of the SCI subjects was lower than that of the controls, after training, 30% of the bone lost in the distal femur and proximal tibia was regained as compared with controls. These findings were associated with increased strength.¹⁹⁷ Another recent study examined 90 male paraplegics who performed isokinetic exercise through electrical stimulation to the quadriceps¹⁹⁸ to determine the optimal training protocol for exercise. Training for 3 days/wk for 30 mins/day with 6 secs of rest and 6 secs of exercise proved the optimal protocol.

Functional Electrical Stimulation for Walking

Functional electrical stimulation (FES) has been used in a very limited fashion to allow individuals with SCI to stand and walk. In a recent postal survey in individuals with SCI in the UK, 4,840 questionnaires were mailed and 1,122 completed surveys were received.¹⁹⁹ Two percent of subjects had used FES for walking, whereas 13% had used some kind of orthosis. Of those who had used FES for walking, >50% had no functional walking ability, whereas most orthotic users had some independent walking ability. Another study compared three groups: 1) FES to the common peroneal nerve through a single channel stimulator, 2) this same stimulator combined with a hinged ankle-foot orthosis, and 3) the use of a hinged ankle-foot orthosis alone with baseline readings with respect to walking.²⁰⁰ In 19 persons with incomplete SCI, gait speed was shown to increase with FES and with the use of a hinged ankle-foot orthosis as compared with baseline; however, the greatest increase in gait speed and endurance occurred with the combined ankle-foot orthosis and FES condition. Foot clearance was shown to improve with FES but not the orthosis. Those subjects with increases in gait speed from FES had weaker hip flexors, knee flexors, and ankle

dorsiflexors than those who did not benefit from FES.

Surface, percutaneous, and implanted electrodes have been used as means of multichannel stimulation to produce standing and walking. With the availability of a commercial FES gait system, various groups have reported on its effects. Active standing through the use of the Parastep-1²⁰¹ (Altimed, Fresno, CA) was shown to result in significantly greater values of $\dot{V}O_2$ than performance of passive standing. In another report,²⁰² 15 patients were able to complete training and ambulate independently with a walker. The mean consecutive walking distance was 52.8 m, and the mean speed was 0.15 m/sec. Follow-up was 40 mos, and only five patients continued to use the system primarily for exercise. The authors concluded that this approach has limited practical usefulness for mobility in daily life but should be considered a resource for persons with SCIs. The use of the Case Western Reserve University/Veterans Affairs implanted standing neuroprosthesis was evaluated in 11 subjects with SCIs²⁰³ who had used the device for >1 yr. Nine subjects continued to use the device regularly for standing or exercising. The system was considered safe, reliable, and easy to use, and all subjects expressed a willingness to repeat the surgery and rehabilitation to achieve the same outcome.

Upper Limb FES

Another use of FES has been to promote improved function in the upper limbs. An implanted neuroprosthesis that allowed C5 or C6 tetraplegic individuals to restore the ability to grasp, hold, and release objects was developed²⁰⁴ and eventually became commercially available as the FREEHAND System (NeuroControl, Valley View, OH). In an international multicenter cohort trial of this system,²⁰⁵ 51 tetraplegic adults underwent concurrent tendon transfer surgery and implantation of the system in which electric stimulation of the grasping muscles of one arm were controlled by using contralateral shoulder movements. With >3 yrs of follow-up, pinch force was significantly greater with the prosthesis in all subjects and grasp-release abilities were improved in 49. Activities of daily living improved in 49 of 49 tested patients, and regular home use was reported in >90% of subjects. In a postal survey²⁰⁶ of 34 individuals with C5 or C6 tetraplegia who received the prosthesis, 87% reported they were very satisfied with the device, 87% reported improvements in activities of daily living, 81% reported improved independence, and 88% reported a positive impact on their lives. A total of 15 participants used the device daily, whereas five individuals did not use the device at all. Another method of use of FES for

the upper limbs was reported through the use of the "Bionic Glove,"²⁰⁷ an external device that uses three channels of electrical stimulation to stimulate finger flexors, extensors, and thumb flexors, with a control signal coming from a wrist-position transducer. In 12 persons with SCIs at C5-C7 who had used the device for ≥ 6 mos,²⁰⁸ daily use of the glove was shown to increase the subject's power grasp and range of movements. The authors recommended that the best candidates for use of the device are complete C6-7 tetraplegics with a FIMTM score between 25 and 50 and a Quadriplegia Index of Function between 0 and 13 and who are motivated to become independent. Another use of electrical stimulation for improved upper limb function is restoration of elbow extension through triceps stimulation. In 11 arms of persons with SCIs who received a triceps electrode as an addition to a hand-grasp neuroprosthesis,²⁰⁹ triceps stimulation provided a significantly stronger elbow extension moment than a posterior deltoid to triceps tendon transfer. The elbow extension moment generated through simultaneous activation of the tendon transfer and triceps electrode was always greater than either method used alone. Addition of elbow extension to the neuroprosthesis resulted in a decreased amount of time required to acquire an object while reaching and significantly increased the ability to successfully reach and move an object.

Other Uses

Other uses for FES include inspiratory muscle pacing and implantable stimulation to control bladder function. In individuals with high tetraplegia but preserved phrenic nerve function, electrical stimulation of these nerves can be used to restore respiratory function.²¹⁰ This procedure has generally been performed via thoracotomy; however, most recently laparoscopic placement of electrodes into the intramuscular portion of the diaphragm has been piloted,²¹¹ and it is hoped that intramuscular pacing can provide similar results to conventional phrenic nerve pacing. An externally controlled implantable neuroprosthesis is available (VOCARE Bladder System, NeuroControl) for management of the neurogenic bladder and bowel.²¹² A multicenter trial evaluated the safety and efficacy of this device 3, 6, and 12 mos postimplantation in 23 patients with SCI. At 1-yr follow-up, 18 of 21 patients were able to urinate >200 ml and 15 had postvoid volumes <50 ml. UTI, reflex incontinence, catheter use, anticholinergic drug use, and autonomic dysreflexia were substantially reduced, and 15 of 17 patients also reported reduction in the time spent in bowel management. One problem with this technique is that device placement is usually performed in conjunction with a sacral

rhizotomy that generally leads to the loss of reflex ejaculation and erection in men. In addition, it has been hypothesized that this procedure could abolish the ability to achieve orgasm post-SCI.⁷⁴

Conclusion

As can be seen from the above discussion, the use of electrical stimulation continues to show great potential for both therapeutic and functional purposes post-SCI. Performance of research in this area is, however, costly and difficult because of its potentially invasive nature and the difficulties associated with blinding of subjects to this type of therapy. Whereas great progress has been made in this field, reports are generally uncontrolled case series. In conclusion, further research targeting randomized controlled trials of noninvasive means of stimulation is needed as a means of assessing the utility of electrical stimulation over standard orthoses. In light of the push toward curative therapies, the development of less-destructive techniques for internal stimulation and the development of consensus on the means to best test them in randomized controlled trials is also warranted. Moreover, if this research is truly to be performed in a randomized fashion, the ethics of subjects paying for the surgical procedures should be discussed.

COMMUNITY INTEGRATION

Successful community integration is an end goal of the rehabilitation process. It is a multifactorial construct that reflects the extent to which a person's preferred level of social and physical involvement in the community is achieved. Given its inherent subjective core, some authors have used qualitative research methodology to explore correlates of successful community integration or participation.²¹³ Other quantitative efforts have been largely descriptive or cross-sectional, with the exception of a level II prognostic study we were able to identify, which revealed that only 9% and 29% of the variance in social integration and the ability to get out of the house and into the community, respectively, are explained by demographic and neurologic variables.²¹⁴ Other work suggests that community integration is better explained by such factors as education level, employment, income, functional independence, and health status than by demographics or neurologic variables.²¹⁵ A level II prognostic study suggested that for a population averaging 20 yrs postinjury, community integration declines over time, with a corresponding decline in satisfaction with life.²¹⁶ Recently, researchers have examined environmental factors as they facilitate or inhibit community integration.²¹⁷⁻²¹⁹ Qualitative investigations²¹⁶ have supported the

importance of the physical environment and attitudinal barriers for community integration, but quantitative approaches have found positive but less robust relationships between environmental factors and societal participation.^{217,218} Availability of independent living services was not associated with better community integration outcomes, but this study was not a true randomized controlled trial.²²⁰

Conclusion

The lack of level I or II intervention attempts in this area is striking but not surprising; interventions would likely have to be community based; such "remote" intervention studies are difficult logistically. However, with the advent of relevant technology (e.g., tele-health) and the availability of existing statewide networks with whom partnerships could be established (e.g., vocational rehabilitation, public health nursing, independent living centers), creative interventions could be formulated (see Forchheimer and Tate²²⁰ for an example). With shortened rehabilitation lengths of stay, such attempts to link and enhance continuity between acute and longer-term care resources could be demonstrably helpful with regard to community integration.

PSYCHOSOCIAL ISSUES

In a report for the American Association of Spinal Cord Injury Psychologists and Social Workers in 1997 reviewing the status of psychosocial research for persons with SCI,²²¹ the authors concluded that there was evidence for psychosocial interventions being "somewhat effective." Effect sizes were found to be small to medium but with clinical relevance. The authors also noted the preponderance of correlational research in this area and the lack of true intervention trials. In the following sections, the reader will note a maturing of this promising body of work in terms of research rigor, but with considerable room for further development.

Depression

Recent published estimates of the prevalence of depression in persons with SCI vary as a function of measures used, how depression was defined, and years postinjury, among other factors. Studies suggest significant rates of depressive syndromes in this group of individuals, with estimates ranging from 16% to 30%.²²²⁻²²⁴ Of note is a Canadian population-based level I prognostic study²²⁵ in which all persons from Alberta who sustained an SCI for the 2-yr period between April 1992 and March 1994 were followed over 6 yrs postinjury. Cases were paired with able-bodied controls matched for age, sex, and region of residence. In-

ternational Classification of Diseases-9 definitions for depression were used. The authors found that 27.5% of persons with SCI were treated for depression during the follow-up interval compared with 10.8% of controls. An attempt in the United States to assess depressive symptoms in persons with SCI using a community-based sampling framework yielded an estimate of 31%, scoring above the cut-off on a widely used measure of depressive symptoms.²²⁶ Based on the more limiting category of probable major depression, 11.4% of 849 persons from the Model Systems²²⁷ met criteria for that diagnosis 1 yr postinjury.

A number of studies have identified risk and protective factors for depression, but other than the level I prognostic study described above,²²⁵ these have been based on cross-sectional designs; hence, they do not meet criteria for prognostic studies reviewed in detail in this article. High levels of wheelchair sports,²²⁸ and higher quality social support,²²⁹ have been associated with lower depression scores. Diagnosed posttraumatic stress syndrome has also been associated with a higher risk for depressive reactions.²³⁰ Finally, a number of authors have demonstrated that coping style can affect outcomes such as depression.²³¹ All of these risk and protective factors could be or have been developed as interventions.

There have been few controlled trials for the treatment of depression for persons with SCI. No controlled trials of antidepressants for this group of individuals were noted, although several noncontrolled trials before the 1999 to present time period did suggest efficacy.²³² In a recent randomized controlled trial of Elavil for pain in SCI,²³³ depression was assessed before and after administration in the study arm. No impact on depression was noted; however, doses were typically lower than would be used for depression. Thus, although this study was classified as a level I trial for pain (see earlier discussion), it was not considered an adequate trial of the efficacy of Elavil for depression. Consistent with the risk factor research reviewed above, two recent level III studies^{234,235} have been reported suggesting that cognitive-behavioral coping-focused psychotherapy either in groups or individually may help ameliorate depressive symptoms. Although both used control groups, they were not true randomized treatment designs. In a small randomized controlled level III study, a group of persons who received massage demonstrated lower posttreatment depression scores than controls who were randomized to an exercise group.²³⁶ Finally, in a small randomized controlled level III trial of the benefits of physical exercise, participants randomized to the exercise group showed improved depression scores after 9 mos of

twice-weekly exercise compared with controls who were provided bimonthly education classes.²³⁷

Conclusion

Given the high prevalence of major depression in community-dwelling individuals with SCI, funding is needed to pursue level I intervention trials through medical, psychological, environmental, or physical (e.g., exercise) modalities. As is the case for most other topics, however, multisite studies are likely needed to ensure adequate sample sizes.

Substance Abuse

Substance abuse has been well documented as both a preinjury and postinjury problem for persons with SCI. In one university-based SCI treatment center, 96% of a consecutive admission sample of 26 newly injured individuals reported preinjury alcohol use, with 57% assessed as heavy drinkers.²³⁸ In a mixed consecutive sample of persons with traumatic brain injury, SCI, or both admitted to a level I trauma center, 17.9% of the sample were characterized as persons who would meet criteria for current alcohol dependence, 21.1% were characterized by heavy alcohol use but with less evidence of physical dependence and fewer negative consequences, and 14.7% characterized by high levels of lifetime alcohol-related problems, but little current alcohol consumption.²³⁹ Alcohol has also been demonstrated through statewide injury surveillance systems to be a risk factor in the accidents that cause SCI.²⁴⁰ The converse, however, may not be true; SCI was not a risk factor in alcohol use patterns for identical twins who sustained an SCI compared with their non-SCI twin.²⁴¹ It has also been well demonstrated in both current²⁴² and previous work that there is a high recidivism rate in this population, although not always to the extent of preinjury use.

Substance use history has been examined as a risk factor in three level II prognostic studies. In a consecutive sample of 87 newly injured persons with SCI admitted to a university medical center, 53% presented with positive screens: 44% alcohol only, 30% drug only, 26% both. Rehabilitation outcomes were the same for those with positive and negative screens, however.²⁴³ Alcohol use postinjury has also been associated with a higher risk for post-SCI injuries,²⁴⁴ pressure sores (level II),²⁴⁵ pain, and diminished quality of life.²⁴⁶ Problem drinking before injury has also been demonstrated in a level II study to be a risk factor for longer initial rehabilitation and lower FIM™ efficiency scores pertaining to SCI and substance abuse.²⁴⁷

There were no treatment studies identified in the recent literature. There are recent treatment

approaches such as motivational interviewing that are being evaluated for efficacy in related populations. A recent randomized, controlled, prospective trial of a brief intervention with 366 patients at a level I trauma center who screened positive for alcohol at admission, suggested a marked positive impact on drinking and injuries requiring hospitalization over 12 mos compared with controls.²⁴⁸

Conclusion

In summary, given the well-documented recidivism rate in this population and the relationship between substance abuse and secondary complications, research is needed to determine the most efficacious methods, both during acute and follow-up care, to prevent a return to substance abuse after discharge from initial hospitalization.

Anxiety Disorders

There has been less research emphasis on anxiety disorders in the SCI population, with relatively few intervention trials. The clinical literature²⁴⁹ has highlighted the need to screen for anxiety disorders based on the cause of SCI (recollection of the trauma) and the experiences that can occur postinjury, ranging from the physical sequelae of SCI to the uncertainties of community, family, vocational, and avocational re-integration.

Base rates for anxiety disorders in non-SCI community samples are relatively low: 1–2% for panic disorder (1-yr prevalence), 3% for generalized anxiety (1-yr prevalence), and 1–14% for post-traumatic disorder.²⁵⁰ Recent estimates of the prevalence of anxiety disorders in the SCI population are typically higher than that, although there is considerable variability.²⁵¹ Virtually all of these estimates are not community- or population-based estimates, however, but are hospital or clinic convenience samples. Fourteen percent of 85 SCI subjects were noted to have high levels of anxiety 6–24 wks postinjury.²⁵²

In a longitudinal study of a sample of persons with SCI from acute rehabilitation through discharge into the community out to 2 yrs postinjury, scores were above a cutoff on a state anxiety measure ranging from 10% to 60%, with scores increasing up to the point of initial discharge from rehabilitation.²²³ In contrast, in a study of 33 persons with SCI and 33 able-bodied controls,²⁵³ no differences were found on a self-report anxiety measure, but there were indications of higher anxiety using the Rorschach, a projective test.

In a presumably more at risk population of veterans many years post-SCI, a study using a screening instrument found that 40% of subjects scored above a cutoff for any anxiety disorder, with 9% at risk for generalized anxiety, 24% for panic

attack, and 21% for posttraumatic stress disorder (PTSD).²²² In another sample of veterans with SCI,²⁵⁴ 11% were noted to have a current diagnosis of PTSD and a lifetime rate of 28%. In a pediatric sample of children with SCI averaging 5.9 yrs postinjury, 25.4% met DSM-IV criteria for a current diagnosis of PTSD.²⁵⁵

More focus has occurred in the recent literature regarding risk and protective factors for PTSD than other anxiety disorders for persons with SCI, but these are typically cross-sectional studies in design; hence, they do not meet criteria for inclusion in our prognostic tables. War theater exposure before onset of SCI increased the risk of a current PTSD diagnosis in veterans compared with their peers who had not been exposed to combat.²⁵⁶ The availability of social support from friends and a higher level of education have been correlated with a lower risk of a current PTSD diagnosis,²⁵⁶ whereas another study noted a relationship between anxiety about death and the presence of PTSD.²⁵⁷ High levels of participation in sports activity were found to be related to lower levels of anxiety after SCI.²²⁸

Few controlled trials exist for the treatment of anxiety for persons with SCI. A recent level III study²³⁵ reported that cognitive-behavioral coping-focused psychotherapy provided in groups, compared with outcomes from historical controls, may help ameliorate anxiety symptoms. In a small, randomized, level III, controlled trial of the benefits of massage, participants assigned to the massage condition showed greater reductions in anxiety scores after 5 wks of treatment than did controls who were assigned to range of motion and strengthening exercises.²³⁶

Conclusion

In conclusion, effective pharmacologic and behavioral treatments for anxiety disorders exist, but there is a virtual absence of evidence for their application to persons with SCI. Future research should include controlled trials of these interventions.

Family Issues: Caregiving, Parenting, Marital Stability

Increasingly, in recent years, the requirements for providing physical care to persons with SCI have fallen on family members. We are beginning to explore and understand this important but complex area in terms of its positive and negative impact on the person who is providing care and in terms of the particular mix of individual, cultural, social, and environmental characteristics that help explain short- and long-term physical and emotional outcomes for those individuals. The impor-

tance of social support for the person with SCI, often provided principally by family members, is also increasingly well recognized.

Marriage

With the profound effects of SCIs, impacts are expected on family issues with regard to marriage and divorce in the SCI population. Authors have generally found marginally higher divorce rates for persons who are married at the time of injury compared with national norms, with the greatest risk for divorce the first year postinjury.²⁵⁸ Postinjury marriages have generally been found to be more stable than preinjury ones.²⁵⁸ Being married may lead to improved quality-of-life outcomes. Putzke et al.²⁵⁹ matched 53 single and married individuals with SCI on education, race, sex, lesion level, pathogenesis, and hospitalizations during the previous year. Married individuals reported higher quality of life and greater levels of participation than their single counterparts.

Caregivers

Caregivers of persons with SCI have been shown to have declines in self-reported quality of life over time²⁶⁰ and cross-sectionally when compared with matched, able-bodied controls.²⁶¹ Individual differences within caregivers in terms of how they cope with distress are important. A negative problem orientation among caregivers has been associated with poorer health, greater depression, lower social functioning, and lower vitality²⁶² and, in a level II prognostic study, a greater rate of developing psychological and health problems over the first year postinjury.²⁶³ In a level II prognostic study, caregivers with a tendency to problem-solve carelessly and impulsively were associated with lower acceptance of disability in the person for whom they provide care and with the development of pressure sores 1 yr postinjury in that person.²⁶⁴

Parenting

Parenting a child with SCI can be associated with elevated distress levels, even several years postinjury. In contrast, there is no evidence that SCI itself results in poorer outcomes for children raised by those individuals. In a cross-sectional study, 31 children raised by a parent with SCI were compared across a variety of dimensions with a matched group of children raised by an able-bodied parent. No differences were noted on any of the several adjustment measures utilized.²⁶⁵

Conclusion

Family caregivers, whether providing care for a child, spouse, parent, or relative, may not experience caregiving as stressful; indeed, for some it

may provide a sense of purpose. However, controlled research is needed for ways to identify caregivers in the community who are significantly distressed and to demonstrate ways of ameliorating that distress. The physical and emotional well-being of the person with SCI who is the recipient of that care should be secondary outcomes of those intervention studies.

Mediating Factors in Individual Coping/Adjusting Spirituality

Experience suggests that the sudden onset of SCI can be either a test of spirituality, potentially weakening personal belief systems, or a source of comfort and confirmation of those belief systems. However, we are not far along in the development of psychometrically sound ways of assessing spirituality,^{266,267} and empirical attempts to study the relationship between spirituality and coping/adjusting have been generally cross-sectional and not prognostic.²⁶⁸ The importance of social support in the coping/adjusting process is also widely accepted. There has been some progress in developing psychometrically sound ways of assessing different types of social support such as emotion-oriented, problem-oriented,²⁶⁹ esteem, informational, and tangible support,^{270,271} and also negatively experienced support.²⁶⁹ However, we were unable to find any therapeutic or prognostic studies in these areas.

Recreation

Much of the recent literature on the effectiveness of recreation therapy on participation for persons with SCI is descriptive in nature²⁷²⁻²⁷⁷ or poorly controlled,²⁷⁸ so cause and effect cannot be inferred. Recent attempts to develop and evaluate a theoretic, predictive model of the impact of leisure on adjusting,²⁷⁹ and to enhance motivation in therapy tasks by modifying traditional exercise activities to recreational ones,^{280,281} provide encouragement that more empirically sound evaluations of the impact of recreation and recreation services can be forthcoming.

Environment

The importance of understanding the limiting or facilitating impact of the environment in terms of adjusting and participation has received appropriate recent attention, with significant theoretic models addressing not only main effects but the importance of the interaction between individual characteristics and the environment.^{282,283} Most environmental facilitators have been shown to be social, whereas the main obstacles are physical. Moreover, the extent to which environment influ-

ences individuals varies, among other factors, as a function of age, sex, and level of injury.²⁸² Other studies have identified level of education²⁸⁴ and transportation^{285,286} as barriers to employment and participation. Use of a wheelchair seems to improve participation in some instances, but in a complex interaction between personal and environmental characteristics.²⁸⁷ Others have struggled with ways to measure and assess the impact of community structure or cohesion in terms of facilitating or limiting participation.²⁸⁸ All of these studies have used cross-sectional methodology and, hence, did not meet the criteria used here for systematic review.

Conclusion

Prognostic studies are needed that utilize well-validated measures of the environment. Therapeutic studies in this area are likely to be complex and ambitious undertakings, focused on the impact of policy change or measured changes in the environment on participation. Conceivably, individuals could be taught advocacy/political activism skills to cause environmental change, with concomitant assessment of the impact of that achievement on participation. Policy changes aimed at improving participation have likely done so, but not often within a scientific context so that the impact can be measured.

Individual Differences

A recent study exploring the prevalence of personality disorders in the SCI population did not find a higher prevalence in this group compared with matched medical controls.²⁸⁹ However, personality characteristics have been associated with subsequent additional injuries post-SCI.²⁹⁰ In two level II prognostic studies,^{295,299} social problem solving orientation and coping abilities were predictive of a variety of psychological and physical outcomes at one and 2 yrs postinjury. Coexisting cognitive dysfunction, be that from concomitant traumatic brain injury, previous head injury, prolonged substance abuse, or others, has been frequently overlooked²⁹¹ or has been associated with negative outcomes.^{292,293} However, prognostic as opposed to cross-sectional descriptive studies are lacking, and no treatment studies (e.g., cognitive remediation) were identified.

One of the more fruitful lines of investigation in recent years has been the adoption of cognitive-behavioral formulations of coping, which have well-developed models, accompanying measures, and importantly, lend themselves to intervention.²⁹⁴ Two level III studies, both of which used historic controls, and both of which adopted a stress and coping training intervention,²⁹⁵ re-

ported fewer hospital readmissions, higher self-reported adjustment after 2 yrs,²⁹⁶ and lower levels of anxiety and depression after 6 wks.²⁹⁷ Other positive constructs of recent interest in the SCI adjustment literature include hope^{300–302} and purpose in life,³⁰³ although prognostic studies and attempts to instill hope or purpose in life, and to examine subsequent outcomes, are lacking. Still others have attempted to delineate positive outcomes as a function of SCI³⁰⁴ rather than measuring negative ones like depression and anxiety. One level III randomized controlled trial³⁰⁵ of the effectiveness of a “wellness” intervention demonstrated some positive benefits in terms of short-term health outcomes.

Conclusion

Prognostic research is needed here rather than the cross-sectional descriptive work that has characterized psychosocial research in SCI over many years. The thrust of this body of work should be in identifying and measuring enduring individual differences that significantly predict outcomes at a future time. Tied to this should be an emphasis on individual differences that are capable of modification, such as coping style and education, rather than more immutable characteristics of a person. In addition, the interaction between individual differences and environmental barriers/facilitators needs further exploration. Controlled interventions should be applied to subgroups with identified deficits or needs, rather than across the board to a convenience sample; not everyone, for example, needs coping effectiveness training. Multicenter trials will be needed to attain adequate sample sizes for such targeted interventions.

Quality of Life

Quality of life has emerged as an increasingly important research endpoint in recent years. However, it is a subjective construct that has proven elusive in terms of the development of any standardized system, accepted across research groups, for its assessment.^{306,307} Some have argued that given its inherent subjectivity, qualitative research methods are warranted.³⁰⁸ Others have argued that although qualitative methods might be appropriate for early exploratory work, they are difficult to replicate, and more objective methods are needed. Accordingly, a variety of standardized quality-of-life scales have been used across studies.

There is a substantial literature in which quality of life (or similar terms) is an outcome. The bulk of this literature, however, is descriptive in methodology, chiefly using interview or survey methods cross-sectionally to identify correlates of quality of life.^{308–325} Because these studies are not

predictive, they do not meet criteria for detailed review here, but they remain important contributions in pointing out potential areas to develop interventions to improve quality of life such as increasing community mobility,³²⁰ decreasing pain,³¹² improving bowel and bladder control,³¹⁴ improving marital stability,³¹⁰ and increasing access to physical activity.³²¹

Quality of life has been used as a secondary outcome in intervention trials in which the primary outcomes were more proximal to the intervention. It has been demonstrated in a level I double-blind randomized trial, for example, that aspects of quality of life were improved via treatment of erectile dysfunction, where the primary outcomes were measures of sexual performance and satisfaction.³²⁶ Similarly, in a level III therapeutic study, exercise training as administered via a randomized controlled trial resulted in improvements in quality of life along with the primary endpoints of strength and stamina.³²⁷ When quality of life is the primary endpoint, studies have more frequently been prognostic in nature rather than interventions designed to improve quality of life.^{328–334} Several of these studies did meet criteria for inclusion in the evidence tables as level I or II prognostic studies.^{329,332–334}

Conclusion

Future research should concentrate on developing or identifying a well-validated, psychometrically sound set of methods for assessing quality of life post-SCI, with their advantages and disadvantages clearly spelled out. Until/unless there is some uniformity of measurement of this important construct across studies, this particular aspect of the SCI literature will be troubled by ambiguity.

EMPLOYMENT

Most recent studies of employment post-SCI indicate that fewer people are employed after injury than was the case preinjury, with some studies showing that gap to be relatively narrow³³⁵ and others indicating a much wider disparity.^{336,337} It has been well demonstrated that increasing time postinjury is associated with increasing rates of return to employment,³³⁵ with one estimate of 6.3 yrs to the first full-time employment.³³⁵ A number of factors have been associated with return to employment, including whether the individual held a professional occupation before injury or was able to return to his or her previous job.³³⁵ Other factors associated with return to employment have included sex, the level of physical exertion required, Barthel Index, education level, age postinjury,³³⁶ and race.³³⁷ Psychological variables including locus of control, and work attitudes have also been

associated with return to work.³³⁸ Individuals injured as children who grow into employment age are less likely to be engaged in paid employment than comparable able-bodied peers,³³⁹ and education, community mobility, functional independence, and decreased medical complications have been associated with their success in employment.³⁴⁰ Factors external to the individual have also been associated with return to employment, including availability of transportation, access difficulties, and perceived workplace discrimination.³⁴¹ However, for those able to overcome these barriers and return to employment, even part-time, benefits have been noted, including increased satisfaction with life and with social integration and decreased levels of depressive symptoms.³⁴² Similar benefits have been noted for volunteer activities.³⁴³

Most research in this area is cross-sectional. Two longitudinal, prognostic studies were identified. In a level I study,³⁴⁴ variables derived at injury were used to predict employment at follow-up years 1, 2, 5, 10, 15, or 20 yrs postinjury. Being white, younger at injury, having lived more years post-SCI, having a less severe injury, and being better educated were predictive of being employed. In another level III prognostic study, positive expectations about returning to work assessed during initial rehabilitation were associated with later return to work.³⁴⁵ Our searches did not turn up any controlled intervention trials and located only one description of early outcomes of an uncontrolled intervention to increase employment outcomes.³⁴⁶

Conclusion

Low return-to-work rates postinjury have been adequately demonstrated over the years, along with descriptive studies identifying correlates of those who do and do not return to work. What is needed are controlled studies of attempts to encourage a return to employment with adequate follow-up to discern long-term work maintenance rates and to demonstrate the positive personal impact (or lack thereof) of a return to work.

ASSISTIVE TECHNOLOGY

Most studies reviewed in this area did not fit well conceptually into the classification rubric used for this review. Much of the literature over the past 5 yrs related to assistive technology documents technological advancements in equipment design, particularly wheelchair design and interfaces. Subjective³⁴⁷ and objective³⁴⁸ evaluations of new equipment are provided by persons with SCI.³⁴⁹ The most rigorous evaluations use both objective and subjective data from both users and experienced staff in comparisons of new technology *vs.* more typical technology.^{350,351} The use of indepen-

dent, experienced observers to evaluate the effectiveness of technology helps somewhat with the obvious methodological problem of the impossibility of “blinding” the assessor. Cost-effectiveness evaluations are not prominent.

Wheelchairs

New wheelchair designs have improved mobility across a variety of terrains,^{350,351} and new, innovative interfaces, including head³⁴⁸ and brainwave interfaces,^{352,353} provide promise for increased mobility, community access, and environmental control³⁵⁴ for persons with very high tetraplegia. Power-assist designs, filling a gap between fully manual and fully powered designs, have been developed, and their proper place in the wheelchair prescription process is being evaluated.³⁵⁵ Considerable recent attention has been given to the role of improvements in the biomechanics of wheelchair designs in the preservation of upper limb function after SCI (see Consortium for Spinal Cord Medicine³⁵⁶ for a thorough evidence-based review). Laboratory-based investigations of the impact of different wheelchair designs^{357–359} and their interaction with individual differences and neurological level of SCI provide promise for optimizing efficiency of wheelchair propulsion,^{360,361} at the same time minimizing the development of upper limb pain and dysfunction over time. In a level II prognostic study³⁶² using a combination of subjective and objective measures and subjects as their own controls, a pushrim-activated power assisted wheelchair was shown to have considerable energy demand advantages over the participants' own wheelchairs. Others have been exploring the biomechanics of push-up maneuvers^{363,364,365}; these studies are the necessary precursor to intervention trials. Another important area of investigation is the assessment of equipment reliability and durability. Objective data in this arena could have important policy implications—for example, providing guidelines to funding agencies as to what normal lifetime expectancy would be for certain types of equipment. In a level II prognostic study of this type, the durability of three different types of manual wheelchairs was objectively demonstrated.³⁶⁶ More work is needed in wheelchair design and evaluation because persons with SCI have listed the wheelchair itself as the most important factor limiting their social participation, above physical impairment and the environment.³⁶⁷

Other Assistive Devices

Work has also been reported on applications of telemedicine for the remote management of complications.³⁶⁸ Access to the Internet is generally reported to be lower for persons with a disability compared with the able-bodied population, but for those who do have access, positive benefits may accrue, particularly with regard to information

gain and knowledge about access to services.³⁶⁹ Finally, access to technology remains problematic based on subjective reports from persons with SCI³⁷⁰ and may be particularly problematic for socioeconomically disadvantaged individuals.³⁷¹

Conclusion

Ongoing attention to objective evaluation of evolving technology is needed, along with cost-benefit analyses of such technology and longer-term evaluations of its utility and resilience in day-to-day community use.

AGING

Most of the work identified in this area was cross-sectional or descriptive, with one prognostic and no therapeutic studies identified. Aging has become an important research focus in recent years, in part because acute and postacute medical care has improved, and people who sustain SCIs are living much longer. Aging is not an outcome or an independent variable but a marker for the passage of time and the cumulative impact of physical, psychosocial, and environmental events and experiences. Research has been focused on discovering trends that might help alert the clinician as to high-risk outcomes that are likely to occur as the result both of chronologic age and years postinjury. Many studies in this area are cross-sectional, with a few notable attempts to gather long-term longitudinal information.^{372–374}

Secondary Conditions

Secondary conditions such as kidney stones and musculoskeletal conditions like fractures³⁷² have been associated with increasing years postinjury. Depressive symptoms, including probable major depression, may also vary with years postinjury, although in a curvilinear fashion: those with the most recent (<5 yrs) and the greatest number of years (≥26 yrs) postinjury demonstrate the highest depression symptomatology.³⁷⁵ There is some evidence based on qualitative research,³⁷⁶ and as summarized in reviews,³⁷⁷ that women have unique concerns and needs as a function of aging with SCI, including concerns about osteoporosis, bladder incontinence and UTIs, dependency, and financial security. Increasingly, physical medicine and rehabilitation physicians are assuming both primary care and geriatric care roles in managing these aging related concerns across the lifespan.³⁷⁸ Interestingly, chronologic age in contrast to years postinjury has not been highly predictive of a variety of outcome domains across measures of functional independence, activity, and participation.^{379,380}

There have been some attempts to examine prognostic factors related to aging in non-cross-sectional designs, but the predictive power of such studies in some cases is limited by the retrospective

nature of the data gathered^{381,382} or cross-sectional analyses of longitudinal data bases.^{383–385} Such studies have been valuable nonetheless in demonstrating concerns about fatigue,³⁸³ decreasing expectations for life satisfaction and independence with aging,³⁸² and marked sex differences in complications and concerns.³⁸¹ In one level II prognostic study³⁸⁶ examining a large cohort of persons with SCI injured ≥ 20 yrs at study entry, measures of stress and depression at entry did not predict stress and depression 3, 6, or 9 yrs later, but perceived quality of life at entry did predict later stress and depression.

Conclusion

Longitudinal, prospective risk factor studies, though difficult, time consuming, and costly to complete, are warranted to provide the best evidence for early identification of markers for later problems, thereby making possible the design and execution of prevention studies.

CURRENT STATUS AND FUTURE RECOMMENDATIONS

Systems of Care

Persons with SCIs have been treated in specialized centers since World War II in England through the National Health Service and in the United States through the Veterans Administration. The goal of establishing these centers was to improve the medical and rehabilitative care of persons with this costly disability. The SCI Model Systems program was started in 1970 with the funding of the Good Samaritan Hospital in Phoenix by the then Rehabilitation Services Administration. Twenty-seven centers in various institutions in the United States have been designated Model Systems since that time. These systems were conceptualized as places where superior clinical care could be given, education provided regarding SCI and its prevention, and lifetime management and research could be conducted, primarily through the development of a longitudinal database and the performance of collaborative studies among the centers. These systems continue to be funded through the National Institute of Disability and Rehabilitation Research.

Rehabilitation Services

Over the years, a number of forces have acted to affect how and where SCI rehabilitation services are provided. Reimbursement patterns have led to dramatically shortened inpatient lengths of stay and a shift to greater outpatient service delivery. Reimbursement policies also encouraged the proliferation of rehabilitation units, with the result that SCI care began being

provided in many more locations than primarily large, tertiary care centers.

Research

Much of the early research emphasis for the field has been on the care and prevention of secondary conditions, both medical and psychosocial, with the goal of improving community participation and quality of life. Such efforts have led to improvements in survival, reductions in morbidity, a greater understanding of the process of adjusting, and an appreciation of the factors involved in living many years with such an injury. Recent theoretical and applied work has highlighted the importance of measuring and ultimately trying to affect the barriers imposed by the natural and built environment in terms of community participation.

Over the past 10 yrs, largely due to the SCI and resultant efforts of the late Christopher Reeve, there has been a burgeoning international interest in research aimed at diminishing the neurologic injury after SCI. New partnerships have formed between basic scientists and clinical researchers, and the schism between these two disciplines has lessened. There are a number of ongoing studies evaluating the efficacy of various pharmacologic and cellular therapies for the treatment of the neurologic consequences associated with acute and chronic SCI. These include pharmacologic studies with minocycline, HP184, and fampridine. In addition, cellular therapies such as the use of autologous incubated macrophages are in phase II studies for the treatment of acute SCI in the United States, and other therapies, such as the transplantation of olfactory ensheathing glia in patients as little as 6 mos post-SCI, are being tested in other parts of the world. At the same time, there are international attempts to bring together basic science and clinical researchers to develop standard protocols for clinical trials post-SCI. Unfortunately, efforts to overcome the challenge of translational research post-SCI still require more coordination. Within these groups of international investigators, there is a need to add more representation from the field of Physical Medicine and Rehabilitation, whose clinicians and researchers have the most experience pertaining to the functioning of persons with chronic SCIs.

Recommendations for Research Development

Within each review section above, we have delineated proposed areas for needed research in diagnostic, prognostic, and/or therapeutic studies. Rather than repeat that information here, we add these comments intended to address broader issues that cut across specific topics. Many of the topics we reviewed were not amenable for classification

according to the evidence-based criteria. Less than 20% of the literature reviewed here met criteria for class I or II studies. Inadequate sample sizes and attrition limit the impact of many studies. Many SCI centers do not have enough persons with new-onset SCI moving through their systems alone to mount adequately powered trials. Adequate funding is needed, therefore, to generate much needed multicenter trials.

Given the current and probable future absence in the United States of mechanisms for determining true population-based estimates of incidence and prevalence, longitudinal epidemiologic work like that of the Model Systems program and of other researchers will continue to be important to inform policy makers about the impact of SCI, its causes, its basic outcomes, and changes in those parameters over time. Effective SCI registries with demonstrated ability to capture all cases of SCI occurring in a region over a specific time frame would need to be funded and in place to assess changes in policy that might affect either the incidence of SCI (such as changes in highway safety regulations, improvements in sports equipment/rules) or the prevalence of SCI (such as improvements in healthcare procedures).

Despite the excitement with respect to cure and the optimism regarding the development of therapies to improve neurologic function after SCI, a number of areas continue to require significant research to optimize the lives of persons with SCI. A better understanding of the natural history of SCI is imperative, as is the development of outcome measures, to effectively describe its physiologic, psychosocial, and functional consequences. Improved management of secondary complications of SCI continues to remain a priority. Moreover, an evidence-based understanding of the benefits of rehabilitation is long overdue. The optimal use of electrical stimulation, robotics, and other assistive technologies in improving the quality of lives of persons with SCI needs to be defined, as does determining the best methods for fully integrating persons with SCI into society. At present, there are few multicenter, large clinical trials related to treatment and prevention of secondary complications of SCI, with the notable exception of a Veterans Administration cooperative study that will be evaluating the efficacy of oxandrolone as an augmentative therapy in the treatment of pressure sores.

Because of the inherent difficulties, small potential sample size, and expenses associated with performance of research in persons with SCIs, it is paramount that funded research topics target the highest priority areas. Requests for proposals for major funding programs (not pilot studies) should require designs capable of attaining level I evidence

in chosen topic areas. This may require expansion of the traditional Model Systems or Veterans Administration study population to include select community-based partners as has been required to complete industry-sponsored studies in a timely fashion. In addition, collaboration with other federal agencies, industry, professional organizations, and consumer groups is imperative. With its rich history of system development, maintenance of a longitudinal database, documented accomplishments, and superior quality of its clinicians and investigators, the SCI Model Systems are well poised to serve as a catalyst to facilitate and focus future clinical research in the field of SCI. Areas to be addressed should be those that are most meaningful in terms of outcomes: both in terms of decreasing the morbidity that occurs in association with SCI and in terms of the psychosocial and physical needs of individuals affected by SCIs and their families.

ACKNOWLEDGMENTS

We thank Michelle Zamora, Neyton M. Balto-dano, MD, and Carol Ramos, MD, for all their help and assistance.

REFERENCES

1. Jackson A, Dijkers M, Devivo M, et al: A demographic profile of new traumatic spinal cord injuries: Change and stability over 30 years. *Arch Phys Med Rehabil* 2004;85: 1740–48
2. DeVivo MJ, Vogel LC: Epidemiology of spinal cord injury in children and adolescents. *J Spinal Cord Med* 2004;27: S4–10
3. Ackery A, Tator C, Krassioukov A: A global perspective on spinal cord injury epidemiology. *J Neurotrauma* 2004;21: 1355–70
4. Burke D, Linden R, Zhang Y, et al: Incidence rates and populations at risk for spinal cord injury: A regional study. *Spinal Cord* 2001;39:274–78
5. Cantu R, Mueller F: Catastrophic spinal injuries in American football, 1977–2001. *Neurosurgery* 2003;53:358–62
6. Surkin J, Gilbert B, Harkey H, et al: Spinal cord injuries in Mississippi: Findings and evaluations, 1992–1994. *Spine* 2000;25:716–21
7. Dryden D, Saunders L, Rowe B, et al: The epidemiology of traumatic spinal cord injury in Alberta, Canada. *Can J Neurol Sci* 2003;30:113–21
8. Pickett W, Simpson K, Walker J, et al: Traumatic spinal cord injury in Ontario Canada. *J Trauma* 2003;55:1070–6
9. DeVivo MJ: The history of the National Spinal Cord Injury Statistical Center Database, 1973–2000 with epidemiological findings from the United States and other countries. *Europa Medicophysica* 2000;36:109–14
10. Bracken MB, Shepard MJ, Collins WF, et al: A randomized, controlled trial of methylprednisolone or naloxone in the treatment of acute spinal-cord injury: Results of the Second National Acute Spinal Cord Injury Study. *N Engl J Med* 1990;322:1405–11
11. Bracken MB, Shepard MJ, Collins WF, et al: Methylprednisolone or naloxone treatment after acute spinal cord injury: 1-year follow-up data. Results of the second National Acute Spinal Cord Injury Study. *J Neurosurg* 1992; 76 23–31

12. Bracken MB, Shepard MJ, Holford TR, et al: Administration of methylprednisolone for 24 or 48 hours or tirilazad mesylate for 48 hours in the treatment of acute spinal cord injury: Results of the Third National Acute Spinal Cord Injury Randomized Controlled Trial. National Acute Spinal Cord Injury Study. *JAMA* 1997;277:1597-604
13. Bracken MB, Shepard MJ, Holford TR, et al: Methylprednisolone or tirilazad mesylate administration after acute spinal cord injury: 1-year follow up. Results of the third National Acute Spinal Cord Injury randomized controlled trial. *J Neurosurg* 1998;89:699-706
14. Hurlbert RJ: The role of steroids in acute spinal cord injury: An evidence-based analysis. *Spine* 2001;26:S39-46
15. Apuzzo : Pharmacological therapy after acute cervical spinal cord injury. *Neurosurg Online* 2002;50:63
16. Fehlings MG: Recommendations regarding the use of methylprednisolone in acute spinal cord injury: Making sense out of the controversy. *Spine* 2001;26:S56-7
17. Bracken MB, Holford TR: Effects of timing of methylprednisolone or naloxone administration on recovery of segmental and long-tract neurological function in NASCIS 2. *J Neurosurg* 1993;79:500-7
18. Geisler FH, Coleman WP, Grieco G, et al: The Sygen Multicenter Acute Spinal Cord Injury Study. *Spine* 2001;26:S87-98
19. Pointillart V, Petitjean ME, Wiart L, et al: Pharmacological therapy of spinal cord injury during the acute phase. *Spinal Cord* 2000;38:71-6
20. Consortium for Spinal Cord Medicine Clinical Practice Guidelines: *Outcomes Following Traumatic SCI: Clinical Practice Guidelines for Health-Care Professionals*. Paralyzed Veterans of America, 1999
21. American Spinal Injury Association/International Medical Society of Paraplegia International Standards for Neurological and Functional Classification of Spinal Cord Injury Patients. Revised 2000, reprinted 2002. Chicago, ASIA/IMSP, 2002
22. Burns AS, Ditunno JF: Establishing prognosis and maximizing functional outcomes after spinal cord injury. *Spine* 2001;26:137-45
23. Brown PJ, Marino RJ, Herbison GH, et al: The 72-hour examination as a predictor of recovery in motor complete quadriplegia. *Arch Phys Med Rehabil* 1991;72:546-8
24. Marino RJ, Ditunno JF, Donovan WH, et al: Neurologic recovery after traumatic SCI: Data from the model SCI systems. *Arch Phys Med Rehabil* 1999;80:1391-6
25. Ditunno JF, Cohen ME, Hauck WW, et al: Recovery of upper-extremity strength in complete and incomplete tetraplegia: A multicenter study. *Arch Phys Med Rehabil* 2000;81:389-93
26. Ishida Y, Tominaga T: Predictors of neurologic recovery in acute central cervical cord injury with only upper extremity impairment. *Spine* 2002;27:1652-7
27. Sherwood AM, Dimitrijevic MR, McKay WB: Evidence of subclinical brain influence in clinically complete spinal cord injury: Discomplete SCI. *J Neurol Sci* 1992;110:90-8
28. McKay WB, Lim HK, Priebe MM, et al: Clinical neurophysiological assessment of residual motor control in post-spinal cord injury paralysis. *Neurorehabil Neural Repair* 2004;18:144-53
29. Kirshblum S, Millis S, McKinley W, et al: Late neurologic recovery after traumatic SCI. *Arch Phys Med Rehabil* 2004;85:1811-7
30. Sipski ML, Jackson A, Gomez-Marin O, et al: Effects of gender on neurological and functional recovery following spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1826-36
31. Harrop JS, Sharan AD, Vaccaro AR, et al: The cause of neurologic deterioration after acute cervical spinal cord injury. *Spine* 2001;26:340-46
32. Greenwald BD, Seel RT, Cifu DX, et al: Gender-related differences in acute rehabilitation lengths of stay, charges, and functional outcomes for a matched sample with spinal cord injury: A multicenter investigation. *Arch Phys Med Rehabil* 2001;82:1181-7
33. Cifu DX, Seel RT, Kreutzer JS, et al: A multicenter investigation of age-related differences in lengths of stay, hospitalization charges, and outcomes for a matched tetraplegia sample. *Arch Phys Med Rehabil* 1999;80:733-40
34. Cifu DX, Huang ME, Kolakovsky-Hayner SA, et al: Age, outcome, and rehabilitation costs after paraplegia caused by traumatic injury of the thoracic spinal cord, conus medullaris, and cauda equine. *J Neurotrauma* 1999;16:805-15
35. Seel RT, Huang ME, Cifu DX, et al: Age-related differences in length of stays, hospitalization costs, and outcomes for an injury-matched sample of adults with paraplegia. *J Spinal Cord Med* 2001;24:241-50
36. Garcia KA, Gaebler-Spira D, Sisom C, et al: Functional improvement after pediatric spinal cord injury. *Am J Phys Med Rehabil* 2002;81:458-63
37. McKinley WO, Johns JS, Musgrove JJ: Clinical presentations, medical complications, and functional outcomes of individuals with gunshot wound-induced spinal cord injury. *Am J Phys Med Rehabil* 1999;78:102-7
38. Putzke JD, Richards JS, Devivo MJ: Gunshot versus non-gunshot spinal cord injury: Acute care and rehabilitation outcomes. *Am J Phys Med Rehabil* 2001;80:366-70
39. Flanders AE, Spettell CM, Friedman DP, et al: The relationship between the functional abilities of patients with cervical spinal cord injury and the severity of damage revealed by MR Imaging. *Am J Neuroradiol* 1999;20:925-34
40. Burns SP, Weaver FM, Parada JP: Management of community-acquired pneumonia in persons with spinal cord injury. *Spinal Cord* 2004;42:450-458
41. Liaw M, Lin M, Cheng P, et al: Resistive Inspiratory muscle training: Its effectiveness in patients with acute complete cervical cord injury. *Arch Phys Med Rehabil* 2000;81:725-6
42. Peterson WP, Barbalata L, Brooks CA, et al: The effect of tidal volumes on the time to wean persons with high tetraplegia from ventilators. *Spinal Cord* 1999;37:284-8
43. Lynch AC, Wong C, Anthony A, et al: Bowel dysfunction following spinal cord injury: A description of bowel function in a spinal cord-injury population and comparison with age and gender matched controls. *Spinal Cord* 2000;38:717-23
44. *Neurogenic Bowel Management in Adults with Spinal Cord Injury: Clinical Practice Guidelines*. Consortium for Spinal Cord Medicine, Paralyzed Veterans of America, 1998
45. Chen C, Chuang T, Tsai Y, et al: Loss of sympathetic coordination appears to delay gastrointestinal transit in patients with spinal cord injury. *Digest Dis Sci* 2004;49:738-43
46. Krogh K, Mosdal C, Laurverg S: Gastrointestinal and segmental colonic transit times in patients with acute and chronic spinal cord lesions. *Spinal Cord* 2000;38:615-21
47. Rajendran SK, Reiser JR, Bauman, et al: Gastrointestinal transit after spinal cord injury: Effects of cisapride. *Am J Gastroenterol* 1992;87:1614-7
48. Krogh K, Jensen M, Gandrup P, et al: Efficacy and tolerability of prucalopride in patients with constipation due to spinal cord injury. *Scand J Gastroenterol* 2002;37:431-6
49. *Physicians Desk Reference*, ed 54. New Jersey, Medical Economics Company, 2000
50. Branagan G, Tromans A, Finnis D: Effect of stoma formation on bowel care and quality of life in patients with spinal cord injury. *Spinal Cord* 2003;41:680-3

51. Apstein MD, Dalecki-Chipperfield K: Spinal cord injury is a risk factor for gallstone disease. *Gastroenterology* 1987;92:966-8
52. Moonka R, Steins SA, Resnick WJ, et al: The prevalence and natural history of gallstones in the spinal cord injured population. *J Am Coll Surg* 1999;189:274-81
53. Rotter KP, Larrain CG: Gallstones in spinal cord injury (SCI): A late medical complication? *Spinal Cord* 2003;41:105-8
54. Moonka R, Stiens SA, Stelzner M: Atypical gastrointestinal symptoms are not associated with gallstones in patients with spinal cord injury. *Arch Phys Med Rehabil* 2000;81:1085-9
55. Moonka R, Stiens SA, Eubank WB, et al: The presentation of gallstones and results of biliary surgery in a spinal cord injured population. *Am J Surg* 1999;178:246-50
56. Tola VB, Chamberlain S, Kostyk SK, et al: Symptomatic gallstones with spinal cord injury. *J Gastrointest Surg* 2000;4:642-7
57. Abrams P, Amarenco G, Bakke A, et al: Tamsulosin: Efficacy and safety in patients with neurogenic lower urinary tract dysfunction due to suprasacral spinal cord injury. *J Urol* 2003;170 (4 pt 1):1242-51
58. De Seze M, Wiart L, de Seze MP, et al: Intravesical capsaicin versus resiniferatoxin for the treatment of detrusor hyperreflexia in spinal cord injured patients: A double-blind, randomized, controlled study. *J Urol* 2004;171:251-5
59. Schurch B, Stohrer M, Kramer G, et al: Botulinum-A toxin for treating detrusor hyperreflexia in spinal cord injured patients: a new alternative to anticholinergic drugs? Preliminary results. *J Urol* 2000;164 (3 pt 1):692-7
60. Chancellor MB, Gajewski J, Ackman CF, et al: Long term followup of the North American multicenter UroLume trial for the treatment of external detrusor-sphincter dys-synergia. *J Urol* 1999;161:1545-50
61. Hansen RB, Biering-Sorensen F, Kristensen JK: Bladder emptying over a period of 10-45 years after a traumatic spinal cord injury. *Spinal Cord* 2004;42:631-7
62. Cardenas DD, Hoffman JM, Kelly E, et al: Impact of a urinary tract infection educational program in persons with spinal cord injury. *J Spinal Cord Med* 2004;27:47-54
63. Waites KB, Canupp KC, Armstrong S, et al: Effect of cranberry extract on bacteruria and pyuria in persons with neurogenic bladder secondary to spinal cord injury. *J Spinal Cord Med* 2004;27:35-40
64. Linsenmeyer TA, Harrison B, Oakley A, et al: Evaluation of cranberry supplement for reduction of urinary tract infections in individuals with neurogenic bladders secondary to spinal cord injury: A prospective, double-blinded, placebo-controlled, crossover study. *J Spinal Cord Med* 2004;27:29-34
65. Reid G, Potter P, Delaney G, et al: Ofloxacin for the treatment of urinary tract infections and biofilms in spinal cord injury. *Intl J Antimicrobial Agents* 2000;13:305-7
66. Darouiche RO, Donovan WH, Del Terzo M, et al: Pilot trial of bacterial interference for preventing urinary tract infection. *Urology* 2001;58:339-44
67. Hull R, Rudy D, Donovan W, et al: Urinary tract infection prophylaxis using *Escherichia coli* 83972 in spinal cord injured patients. *J Urol* 2000;163:872-7
68. Chen Y, DeVivo MJ, Roseman JM: Current trend and risk factors for kidney stones in persons with spinal cord injury: A longitudinal study. *Spinal Cord* 2000;38:346-53
69. Chen Y, DeVivo MJ, Stover SL, et al: Recurrent kidney stone: A 25 year follow-up study in persons with spinal cord injury. *Urology* 2002;60:228-32
70. Groah SL, Weitzenkamp DA, Lammerste DP, et al: Excess risk of bladder cancer in spinal cord injury: Evidence for an association between indwelling catheter use and bladder cancer. *Arch Phys Med Rehabil* 2002;83:346-51
71. West DA, Cummings JM, Longo WE, et al: Role of chronic catheterization in the development of bladder cancer in patients with spinal cord injury. *Urology* 1999;53:292-7
72. Konety BR, Nguyen TT, Brenes G, et al: Clinical usefulness of the novel marker BLCA-4 for the detection of bladder cancer. *J Urol* 2000;164:634-9
73. Schmid DM, Curt A, Hauri D, et al: Clinical value of combined electrophysiological and urodynamic records to assess sexual disorders in spinal cord injured men. *Neuro-urol Urodyn* 2003;22:314-21
74. Sipski ML, Alexander CJ, Rosen RC: The neurologic basis of sexual arousal and orgasm in women: Effects of spinal cord injury. *Ann Neurol* 2001;49:35-44
75. Zaslaw S, Nicoli C, Galea G, et al: A simplified pharmacologic erection program for patients with spinal cord injury. *J Spinal Cord Med* 1999;22:303-7
76. Bodner DR, Haas CA, Krueger B, et al: Intraurethral al-prostadil for treatment of erectile dysfunction in patients with spinal cord injury. *Urology* 1999;53:199-202
77. Giuliano F, Hultling C, El Masry WE, et al: Randomized trial of sildenafil for the treatment of erectile dysfunction in spinal cord injury. *Ann Neurol* 1999;46:15-21
78. Maytom MC, Derry FA, Dinsmore WW: A two-part pilot study of sildenafil (VIAGRA) in men with erectile dysfunction caused by spinal cord injury. *Spinal Cord* 1999;37:110-6
79. Ethans KD, Casey AR, Schryvers OI, et al: The effects of sildenafil on the cardiovascular response in men with spinal cord injury at or above the sixth thoracic level. *J Spinal Cord Med* 2003;27:222-6
80. Sipski ML, Alexander CJ, Guo X, et al: Cardiovascular effects of sildenafil in men with SCIs at and above T6. *Top Spinal Cord Inj Rehabil* 2003;8:26-34
81. Del Popolo G, Li arzi V, Mondaini N, et al: Time/duration effectiveness of sildenafil versus tadalafil in the treatment of erectile dysfunction in spinal cord-injured patients. *Spinal Cord* 2004;42:643-8
82. Sipski ML, Rosen RC, Alexander CJ, et al: Sildenafil effects on sexual and cardiovascular responses in women with spinal cord injury. *Urology* 2000;55:812-5
83. Sipski ML, Rosen RC, Alexander CJ, et al: Sexual responsiveness in women with spinal cord injuries: Differential effects of anxiety-eliciting stimulation. *Arch Sex Behav* 2004;33:295-302
84. Sipski ML, Rosen RC, Alexander CJ, et al: Effects of positive feedback on sexual arousal in women with spinal cord injury. *Neurorehabil* 2000;15:145-3
85. Heruti RJ, Katz H, Menashe Y, et al: Treatment of male infertility due to spinal cord injury using rectal probe electroejaculation: The Israeli experience. *Spinal Cord* 2001;39:168-75
86. Brackett NL, Ferrell SM, Aballa TC, et al: An analysis of 653 trials of penile vibratory stimulation in men with spinal cord injury. *J Urol* 1998;159:1931-4
87. Bird VG, Brackett NL, Lynne CM, et al: Reflexes and somatic responses as predictors of ejaculation by penile vibratory stimulation in men with spinal cord injury. *Spinal Cord* 2001;39:514-9
88. Sonksen J, Biering-Sorensen F, Kristensen JK: Ejaculation induced by penile vibratory stimulation in men with spinal cord lesion. The importance of the vibratory amplitude. *Paraplegia* 1994;32:651-60
89. Sonken J, Ohl DA, Giwerzman A, et al: Effect of repeat ejaculation on semen quality in spinal cord injured men. *J Urol* 1999;161:1163-5
90. Jackson AB, Wadley V: A multicenter study of women's self-reported reproductive health after spinal cord injury. *Arch Phys Med Rehabil* 1999;80:1420-8

91. Fuhrer MJ, Garber SL, Rintala DH, et al: Pressure ulcers in community-resident persons with spinal cord injury: Prevalence and risk factors. *Arch Phys Med Rehabil* 1993;74:1172-7
92. Model SCI Centers Annual Report. National SCI Statistical Center, University of Alabama at Birmingham, 1998
93. Richardson RR, Meyer PR: Prevalence and incidence of pressure sores in acute spinal cord injuries. *Paraplegia* 1981;19:235-47
94. Krause JS, Vines CL, Farley TL, et al: An exploratory study of pressure ulcers after spinal cord injury: Relationship to protective behaviors and risk factors. *Arch Phys Med Rehabil* 2001;82:107-13
95. Pressure ulcer prevention and treatment following spinal cord injury: A clinical practice guideline for health-care professionals. Consortium for Spinal Cord Medicine, Paralyzed Veterans of America, 2000
96. Miller H, Delozier J: Cost implications, in Bergstrom N, Cuddigan J (eds): *Treating Pressure Ulcers: Guideline Technical Report, Vol II, No. 14*. Rockville, MD, US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research, 1994. Publication 96-N015
97. Inman KJ, Sibbald WJ, Rutledge FS, et al: Clinical utility and cost-effectiveness of an air-suspension bed in the prevention of pressure ulcers. *JAMA* 1993;269:1139-43
98. Baker L, Rubayi S, Villar F, et al: Effect of electrical stimulation waveform on healing of ulcers in human beings with spinal cord injury. *Wound Rep Regen* 1996;4:21-8
99. Griffin J, Tooms R, Mendius RA, et al: Efficacy of high-voltage pulsed current for healing of pressure ulcers in patients with spinal cord injury. *Phys Ther* 1991;72:433-42
100. Stefanovska A, Vodovnik L, Benko H, et al: Treatment of chronic wounds by means of electric and electromagnetic field: Part 2. Value of FES parameters for pressure sore treatment. *Med Biol Eng Comput* 1993;31:213-20
101. Consortium for Spinal Cord Medicine Clinical Practice Guidelines: *Prevention of Thromboembolism in Spinal Cord Injury*, ed 2. Paralyzed Veterans of America, 1999
102. Green D, Hartwig D, Chen D, et al: Spinal cord injury risk assessment for thromboembolism. *Am J Phys Med Rehabil* 2003;82:950-6
103. Prevention of venous thromboembolism in the acute treatment phase after SCI: A randomized, multicenter trial comparing low-dose heparin plus intermittent pneumatic compression with enoxaparin. Spinal Cord Injury Thromboprophylaxis Investigators. *J Trauma* 2003;54:1116-26
104. Prevention of venous thromboembolism in the rehabilitation phase after SCI: Prophylaxis with low-dose heparin or enoxaparin. Spinal Cord Injury Thromboprophylaxis Investigators. *J Trauma* 2003;54:1111-5
105. Maxwell RA, Chavarria-Aguilar M, Cockerham WT, et al: Routine prophylactic vena cava filtration is not indicated after acute spinal cord injury. *J Trauma* 2002;52:902-6
106. Kuijk AA, Geurts ACH, van Kuppevelt HJM: Neurogenic heterotopic ossification in spinal cord injury. *Spinal Cord* 2002;40:313-26
107. Banovac K, Williams JM, Patrick LD, et al: Prevention of heterotopic ossification after spinal cord injury with indomethacin. *Spinal Cord* 2001;39:370-4
108. Banovac K, Williams JM, Patrick LD, et al: Prevention of heterotopic ossification after spinal cord injury with COX-2 selective inhibitor (rofecoxib). *Spinal Cord* 2004;42:707-10
109. Sautter-Bihl ML, Liebermeister E, Nanassy A: Radiotherapy as a local treatment option for heterotopic ossification in patients with spinal cord injury. *Spinal Cord* 2000;38:33-6
110. Frey-Rindova P, deBruin ED, Stussi E, et al: Bone mineral density in upper and lower extremities during 12 months after spinal cord injury measured by peripheral quantitative computed tomography. *Spinal Cord* 2000;38:26-32
111. Garland DE, Adkins RH, Kushwaha Vand Stewart C: Risk factors for osteoporosis at the knee in the spinal cord injury population. *J Spinal Cord Med* 2004;27:202-6
112. Garland DE, Adkins RH, Stewart CA, et al: Regional osteoporosis in women who have a complete spinal cord injury. *J Bone Joint Surg Am* 2001;83:1195-200
113. Bauman WA, Spungen AM: Disorders of carbohydrate and lipid metabolism in veterans with paraplegia or quadriplegia: A model of premature aging. *Metabolism* 1994;43:749-56
114. Bauman WA, Spungen AM: Carbohydrate and lipid metabolism in chronic spinal cord injury. *J Spinal Cord Med* 2001;24:266-77
115. Moussavi RM, Ribas-Cardus F, Rintala DH, et al: Dietary and serum lipids in individuals with spinal cord injury living in the community. *J Rehabil Res Dev* 2001;38:225-33
116. Dallmeijer AJ, van der Woude LH, van Kamp GJ, et al: Change in lipid, lipoprotein and apolipoprotein profiles in persons with spinal cord injuries during the first 2 years post-injury. *Spinal Cord* 1999;37:96-102
117. Bauman WA, Adkins RH, Spungen AM, et al: The effect of residual neurological deficit on oral glucose tolerance in persons with chronic spinal cord injury. *Spinal Cord* 1999;37:765-71
118. Bauman WA, Adkins RH, Spungen AM, et al: Is immobilization associated with an abnormal lipoprotein profile? Observations from a diverse cohort. *Spinal Cord* 1999;37:485-93
119. Szlachcic Y, Adkins RH, Adal T, et al: The effect of dietary intervention on lipid profiles in individuals with spinal cord injury. *J Spinal Cord Med* 2001;24:26-9
120. Spungen AM, Adkins RH, Stewart CA, et al: Factors influencing body composition in persons with spinal cord injury: A cross-sectional study. *J Appl Physiol* 2003;95:2398-407
121. Jones LM, Legge M, Goulding A: Health body mass index values often underestimate body fat in men with spinal cord injury. *Arch Phys Med Rehabil* 2003;84:1068-71
122. DeVivo MJ, Krause JS, Lammertse DP: Recent trends in mortality and causes of death among persons with spinal cord injury. *Arch Phys Med Rehabil* 1999;80:1411-9
123. Smyth MD, Peacock WJ: The surgical treatment of spasticity. *Muscle Nerve* 2000;23:153-63
124. Ashworth B: Preliminary trial of carisoprodol in multiple sclerosis. *Practitioner* 1964;192:540-2
125. Sherwood AM, Graves DE, Priebe MM: Altered motor control and spasticity after spinal cord injury: Subjective and objective assessment. *J Rehabil Res Dev* 2000;37:41-52
126. Skold C: Spasticity in spinal cord injury: self- and clinically rated intrinsic fluctuations and intervention-induced changes. *Arch Phys Med Rehabil* 2000;81:144-9
127. Bohannon RW, Smith MB: Interrater reliability of a modified Ashworth scale of muscle spasticity. *Phys Ther* 1987;67:206-7
128. Akkman MN, Bengi R, Karatas, et al: Assessment of spasticity using isokinetic dynamometry in patients with spinal cord injury. *Spinal Cord* 1999;37:638-43
129. Kakebeeke TH, Lechner H, Baumberger M, et al: The importance of posture on the isokinetic assessment of spasticity. *Spinal Cord* 2002;40:236-43
130. Bryce T, Ragnarsson K: Epidemiology and classification of pain after spinal cord injury. *Top Spinal Cord Inj Rehabil* 2001;7:1-17
131. Finnerup N, Johannesen I, Sindrup S, et al: Pain and

- dysesthesia in patients with spinal cord injury: A postal survey. *Spinal Cord* 2001;39:256–62
132. Siddall P, Loeser J: Pain following spinal cord injury. *Spinal Cord* 2001;39:63–73
 133. Ravenscroft A, Ahmed Y, Burnside I: Chronic pain after SCI: A patient survey. *Spinal Cord* 2000;38:611–4
 134. Putzke J, Richards J: Pain after spinal cord injury: Important predictors, outcome, future directions in cost-effectiveness analysis. *Top Spinal Cord Inj Rehabil* 2001;7:84–92
 135. Werhagen L, Budh C, Hutling C, et al: Neuropathic pain after traumatic spinal cord injury: Relations to gender, spinal level, completeness, and age at the time of injury. *Spinal Cord* 2004;18 (2):164–71
 136. Price G, Kendall M, Amsters D, Perhouse K: Perceived causes of change in function and quality of life for people with long duration spinal cord injury. *Clin Rehabil* 2004;18:164–71
 137. Widerstrom-Noga E, Felipe-Cuervo E, Yezierski R: Chronic pain after spinal cord injury: Interference with sleep and daily activities. *Arch Phys Med Rehabil* 2001;82:1571–7
 138. Siddall P, McClelland J, Rutkowski S, et al: A longitudinal study of the prevalence and characteristics of pain in the first 5 years following spinal cord injury. *Pain* 2003;103:249–57
 139. Nyland J, Quigley P, Huang C, et al: Preserving transfer independence among individuals with spinal cord injury. *Spinal cord* 2000;38:649–57
 140. Dalyan M, Cardenas D, Gerard B: Upper extremity pain after spinal cord injury. *Spinal Cord* 1999;37:191–5
 141. Pattany P, Yezierski R, Widerstrom-Noga E, et al: Proton magnetic resonance spectroscopy of the thalamus in patients with chronic neuropathic pain after spinal cord injury. *Am J Neuroradiol* 2002;23:901–5
 142. Ballinger D, Rintala D, Hart K: The relation of shoulder pain and range of motion problems to functional limitations, disability and perceived health of men with spinal cord injury: A multifaceted longitudinal study. *Arch Phys Med Rehabil* 2000;81:1575–81
 143. Putzke JD, Richards JS, Hicken BL, et al: Interference due to pain following spinal cord injury: Important predictors and impact on quality of life. *Pain* 2002;231–42
 144. Siddall P, Yezierski R, Loeser J: Pain following spinal cord injury: Clinical features, prevalence, and taxonomy. *IASP Newslett* 2000;3–7
 145. Boninger M, Towers J, Cooper R, et al: Shoulder imaging abnormalities in individuals with paraplegia. *J Rehabil Res Dev* 2001;38:401–8
 146. Putzke J, Richards J, Dowler R: The impact of pain in spinal cord injury: A case-control study. *Rehabil Psychol* 2000;45:386–401
 147. Widerstrom-Noga E, Turk D: Types and effectiveness of treatments used by people with chronic pain associated with spinal cord injuries: Influence of pain and psychosocial characteristics. *Spinal Cord* 2003;41:600–9
 148. Levenhdoglu F, Ogun C, Ozerbil O, et al: Gabapentin is a first line drug for the treatment of neuropathic pain in spinal cord injury. *Spine* 2004;29:743–51
 149. Tai Q, Kirshblum S, Chen B, et al: Gabapentin in the treatment of neuropathic pain after spinal cord injury: A prospective, randomized, double-blind, crossover trial. *J Spinal Cord Med* 2002;25:100–5
 150. Kvarnstrom A, Karlsten R, Quiding H, et al: The analgesic effect of intravenous ketamine and lidocaine on pain after spinal cord injury. *Acta Anaesth Scand* 2004;48:498–506
 151. Curtis K, Tyner T, Zachary L, et al: Effect on a standard exercise protocol on shoulder pain in long-term wheelchair users. *Spinal cord* 1999;37:421–9
 152. Siddall P, Molloy A, Walker S, et al: The efficacy of intrathecal morphine and clonidine in the treatment of pain after spinal cord injury. *Anesth Analg* 2000;91:1493–8
 153. Finnerup N, Sindrup S, Bach F, et al: Lamotrigine in spinal cord injury pain: A randomized controlled trial. *Pain* 2002;96:375–83
 154. Cardenas D, Warms C, Turner J, et al: Efficiency of amitriptyline for the relief of pain in spinal cord injury: Results of a randomized controlled trial. *Pain* 2002;96:365–73
 155. Hicks A, Martin K, Ditor D, et al: Long-term exercise training in persons with spinal cord injury: Effects on strength, arm ergometry performance, and psychological well-being. *Spinal Cord* 2003;41:34–43
 156. Dyson-Hudson T, Shifflet S, Kirshblum S, et al: Acupuncture and Trager psychophysical integration in the treatment of wheelchair user's shoulder pain in individuals with spinal cord injury. *Arch Phys Med Rehabil* 2001;82:1038–46
 157. Penn R: Treatment of pain in persons with spinal cord injury: Spinal cord stimulation and intrathecal drugs. *Top Spinal Cord Inj Rehabil* 2001;7:51–6
 158. Lammertse DP, Falci SP: Surgical management of central pain in persons with spinal cord injury: The dorsal root entry zone procedure. *Top Spinal Cord Inj Rehabil* 2001;7:41–50
 159. Falci S, Best L, Bayles R, et al: Dorsal root entry zone microcoagulation for spinal cord injury-related central pain: Operative intramedullary electrophysiological guidance and clinical outcome. *J Neurosurg* 2002;97:193–200
 160. Wegener S, Haythornthwaite JA: Psychological and behavioral issues in the treatment of pain after spinal cord injury. *Top Spinal Cord Inj Rehabil* 2001;7:73–83
 161. Apple D, Richards J, Ragnarsson, K: Pain management. *Top Spinal Cord Inj Rehabil* 2001;7:1–105
 162. Nayak S, Shifflet S, Schoenberger N, et al: Is acupuncture effective in treating chronic pain after spinal cord injury? *Arch Phys Med Rehabil* 2001;82:1578–86
 163. Jonsson M, Tollback A, Gonzalez H, et al: Inter-rater reliability of the 1992 international standards for neurological and functional classification of incomplete spinal cord injury. *Spinal Cord* 2000;38:675–9
 164. Krassioukov A, Wolfe DL, Hsieh JTC, et al: Quantitative sensory testing in patients with incomplete spinal cord injury. *Arch Phys Med Rehabil* 1999;80:1258–63
 165. Lee DC, Lim HK, McKay WB, et al: Toward an objective interpretation of surface EMG patterns: A voluntary response index (VRI). *J Electromyogr Kinesiol* 2004;14:379–88
 166. Ditunno JF, Ditunno PL, Graziani V, et al: Walking index for spinal cord injury (WISCI): An international multi-center validity and reliability study. *Spinal Cord* 2000;38:234–43
 167. Ditunno PL, Ditunno, JF: Walking index for spinal cord injury (WISCI II): Scale revision. *Spinal Cord* 2001;39:654–6
 168. Itzkovich M, Tripolski M, Zeilig G, et al: Rasch analysis of the Catz-Itzkovich spinal cord independence measure. *Spinal Cord* 2002;40:396–407
 169. Gresham GE, Labi MLC, Dittmar SS, et al: The quadriplegia index of function (QIF): Sensitivity and reliability demonstrated in a study of thirty quadriplegic patients. *Paraplegia* 1986;24:38–44
 170. Kennedy P, Evan M, Berry C, et al: Comparative analysis of goal achievement during rehabilitation for older and younger adults with spinal cord injury. *Spinal Cord* 2003;41:44–52
 171. Berry C, Kennedy P: A psychometric analysis of the Needs Assessment Checklist (NAC). *Spinal Cord* 2003;41:490–501
 172. May L, Warren S: Measuring quality of life of persons with

- spinal cord injury: Substantive and structural validation. *Qual Life Res* 2002;40:341–50
173. Itzkovich M, Tamir A, Philo O, et al: Reliability of the Catz-Itzkovich spinal cord independence measure assessment by interview and comparison observation. *Am J Phys Med Rehabil* 2003;81:267–72
174. Young A, Murphy G: A social psychology approach to measuring vocational rehabilitation intervention effectiveness. *J Occup Rehabil* 2002;12:175–89
175. Whiteneck G, Harrison-Felix C, Mellick D, et al: Quantifying environmental factors: A measure of physical, attitudinal, service, productivity, and policy barriers. *Arch Phys Med Rehabil* 2004;1324–35
176. Walker N, Mellick D, Brooks et al: Measuring participation across impairment groups using Craig Handicap Assessment Reporting Technique. *Am J Phys Med Rehabil* 2003; 82:936–41
177. Campagnolo D, Filart R, Millis S, et al: Appropriateness of the Ifield Psychiatric Symptom Index as a screening tool for depressive symptomatology on persons with spinal cord injury. *J Spinal Cord Med* 2002;129–32
178. Scherer M, Cushman L: Measuring subjective quality of life following spinal cord injury: A validation study of the assistive technology device predisposition assessment. *Disabil Rehabil* 2001;23:387–93
179. Jang Y, Hsieh C, Wang Y, et al: A validation study of the WHOQOL-BREF assessment in persons with traumatic spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1890–5
180. Scherer M, Cushman L: Predicting satisfaction with assistive technology for a sample of adults with new spinal cord injuries. *Psychol Rep* 2000;87:981–7
181. Cushman L, Scherer M: A pilot study of perceived needs of persons with new spinal cord injury. *Psychol Rep* 2002; 90:1153–60
182. Jacobs PL, Mahoney ET, Robbins A, et al: Hypokinetic circulation in persons with paraplegia. *Med Sci Sports Exerc* 2002;34:1401–7
183. Dietz V, Colombo G, Jensen L, et al: Locomotor capacity of spinal cord in paraplegic patients. *Ann Neurol* 1995;37: 574–82
184. Postans NJ, Hasler JP, Granat MH, et al: Functional electric stimulation to augment partial weight-bearing supported treadmill training for patients with acute incomplete spinal cord injury: A pilot study. *Arch Phys Med Rehabil* 2004;85:604–10
185. Field-Fote EC: Combined use of body weight support, functional electric stimulation, and treadmill training to improve walking ability in individuals with chronic incomplete spinal cord injury. *Arch Phys Med Rehabil* 2001;82: 818–24
186. Colombo G, Joerg M, Schreiber R, et al: Treadmill training of paraplegic patients using a robotic orthosis. *J Rehabil Res Dev* 2000;37:693–700
187. Jezernik S, Colombo G, Keller T, et al: Robotic Orthosis Lokomat: A rehabilitation and research tool. *Neuromodulation* 2003;6:108–15
188. Merli CJ, Crabbe S, Paluzzi RG, et al: Etiology, incidence and prevention of deep vein thrombosis in acute spinal cord injury. *Arch Phys Med Rehabil* 1993;74:1199–205
189. Creasey FH, Ho CH, Triolo RJ, et al: Clinical applications of electrical stimulation after spinal cord injury. *J Spinal Cord Med* 2004;27:365–75
190. Ragnarsson KR: Physiologic effects of functional electrical stimulation-induced exercises in spinal cord-injured individuals. *Clin Orthop Rel Res* 1988;233:53–63
191. Rodgers MM, Glaser RM, Figoni SF, et al: Musculoskeletal responses of spinal cord injured individuals to functional neuromuscular stimulation-induced knee extension exercise training. *J Rehabil Res Dev* 1991;28:19–26
192. Scremin AM, Kurta L, Gentili A, et al: Increasing muscle mass in spinal cord injured persons with a functional electrical stimulation exercise program. *Arch Phys Med Rehabil* 1999;80:1531–6
193. Hooker SP, Figoni SF, Rodgers MM, et al: Physiologic effects of electrical stimulation leg cycle exercise training in spinal cord injured persons. *Arch Phys Med Rehabil* 1992;73:470–6
194. Bauman WA, Alexander LR, Zhong YG, et al: Stimulated leg ergometry training improves body composition and HDL-cholesterol values. *J Am Paraplegia Soc* 1994;17:201
195. Morh T, Dela F, Handberg A, et al: Insulin action and long-term electrically induced training in individuals with spinal cord injuries. *Med Sci Sports Exerc* 2001;33: 1247–52
196. Mohr T, Podenphant J, Biering-Sorensen F, et al: Increased bone mineral density after prolonged electrically induced cycle training of paralyzed limbs in spinal cord injured man. *Calcif Tissue Int* 1997;61:22–5
197. Belanger M, Stein RB, Wheeler BD, et al: Electrical stimulation: Can it increase muscle strength and reverse osteopenia in spinal cord injured individuals? *Arch Phys Med Rehabil* 2000;81:1090–8
198. Petrofsky JS, Stacy R, Laymon M: The relationship between exercise work intervals and duration of exercise on lower extremity training induced by electrical stimulation in humans with spinal cord injuries. *Eur J Appl Physiol* 2000;82:504–9
199. Maxwell DJ, Granat MH, Baardman G, et al: Demand for and use of functional electrical stimulation systems and conventional orthoses in the spinal lesioned community of the UK. *Artif Organs* 1999;23:410–2
200. Kim CM, Eng JJ, Whittaker MW: Effects of a simple functional electric system and/or hinged ankle-foot orthosis on walking in persons with incomplete spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1718–23
201. Jacobs PL, Johnson B, Mahoney ET: Physiologic responses to electrically assisted and frame-supported standing in persons with paraplegia. *J Spinal Cord Med* 2003;6:384–9
202. Brissot R, Gallien P, LeBot MP, et al: Clinical experience with functional electrical stimulation-assisted gait with Parastep in spinal cord-injured patients. *Spine* 2000;25: 501–8
203. Agarwal S, Triolo RJ, Kobetic R, et al: Long-term user perceptions of an implanted neuroprosthesis for exercise, standing and transfers after spinal cord injury. *J Rehabil Res Dev* 2003;40:241–52
204. Keith MS, Peckham PH, Thrope GB, et al: Implantable functional FNS in the tetraplegic hand. *J Hand Surg Am* 1989;14:524–30
205. Peckham PH, Keith MW, Kilgore KL, et al: Efficacy of an implanted neuroprosthesis for restoring hand grasp in tetraplegia: A multicenter study. *Arch Phys Med Rehabil* 2001;82:1380–8
206. Wuolle KS, Van Doren CL, Bryden AM, et al: Satisfaction with and usage of a hand neuroprosthesis. *Arch Phys Med Rehabil* 1999;80:206–13
207. Prochazka A, Gauthier M, Wieler M, et al: The bionic glove: An electrical stimulator garment that provides controlled grasp and hand opening in quadriplegia. *Arch Phys Med Rehabil* 1997;78:608–14
208. Popovic D, Stojanovic A, Pjanovic A, et al: Clinical evaluation of the bionic glove. *Arch Phys Med Rehabil* 1999;80: 299–304
209. Member WD, Crago PE, Keith MW: Restoration of elbow extension via functional electrical stimulation in individuals with tetraplegia. *J Rehabil Res Dev* 2003;40:477–86
210. DiMarco AG: Diaphragm pacing in patients with spinal cord injury. *Top Spinal Cord Inj Rehabil* 1999;5:6–20
211. DiMarco AF, Onders RP, Kowalski KE, et al: Phrenic nerve pacing in a tetraplegic patient via intramuscular dia-

- phragm electrodes. *Am J Respir Crit Care Med* 2002;166:1604-6
212. Creasey GH, Grill JH, Korsten M, et al: An implantable neuroprosthesis for restoring bladder and bowel control to patients with spinal cord injuries: A multicenter trial. *Arch Phys Med Rehabil* 2001;82:1512-9
 213. Boschen K, Tonack M, Gargaro J: Long-term adjustments and community reintegration following spinal cord injury. *Int J Rehabil Res* 2003;26:157-64
 214. Whiteneck G, Tate D, Charlifue, S: Predicting community reintegration after spinal cord injury from demographic and injury characteristics. *Arch Phys Med Rehabil* 1999;80:1485-91
 215. Anderson C, Krajci K, Vogel L: Community integration among adults with spinal cord injuries sustained as children or adolescents. *Dev Med Child Neurol* 2003;45:129-34
 216. Charlifue S, Gerhart, K: Community integration in spinal cord injury of long duration. *NeuroRehabil* 2004;19:91-101
 217. Whiteneck G, Meade M, Dijkers M, et al: Environmental factors and their role in participation and life satisfaction after spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1793-803
 218. Dijkers M, Yavuzer G, Ergin S, et al: A tale of two countries: Environmental impacts on social participation after spinal cord injury. *Spinal Cord* 2002;40:351-62
 219. Levins S, Redenbach D, Dyck I: Individual and societal influences on participation in physical activity following spinal cord injury: A qualitative study. *Phys Ther* 2004;84:496-509
 220. Forchheimer JM, Tate D: Enhancing community re-integration following spinal cord injury. *NeuroRehabil* 2004;19:103-13
 221. McAweeney JM, Tate DG: The status of psychosocial research in the rehabilitation of persons with spinal cord injury: Implications for practitioners, researchers, and health care reform. American Association of SCI Psychologists and Social Workers, 1997
 222. Weingardt K, Hsu J, Dunn M: Brief screening for psychosocial and substance abuse disorders in veterans with long-term spinal cord injury. *Rehabil Psychol* 2001;46:271-8
 223. Kennedy P, Rogers B: Anxiety and depression after spinal cord injury: A longitudinal analysis. *Arch Phys Med Rehabil* 2000;81:932-7
 224. Kemp B, Krause J: Depression and life satisfaction among people aging with post-polio and spinal cord injury. *Disabil Rehabil* 1999;21:241-9
 225. Dryden D, Saunders L, Rowe B, et al: Utilization of health services following spinal cord injury: A 6-year follow up study. *Spinal Cord* 2004;42:513-25
 226. Fuhrer M, Rintala D, Hart K, et al: Depressive symptomatology in persons with spinal cord injury who reside in the community. *Arch Phys Med Rehabil* 1993;74:255-60
 227. Bombardier C, Richards J, Krause J, et al: Symptoms of major depression in people with spinal cord injury: Implications for screening. *Arch Phys Med Rehabil* 2004;85:1749-56
 228. Muraki S, Tsunawake N, Hiramatsu S, et al: The effect of frequency and mode of sports activity on the psychosocial status in tetraplegics and paraplegics. *Spinal Cord* 2001;38:309-14
 229. Beedie A, Kennedy P: Quality of social support predicts homelessness and depression post spinal cord injury. *J Clin Psychol Med Settings* 2002;9:227-34
 230. Nielsen M: Post-traumatic stress disorder and emotional distress in persons with spinal cord lesions. *Spinal Cord* 2003;41:296-302
 231. Galvin L, Godfrey H: The impact of coping on emotional adjustment to spinal cord injury (SCI): Review of the literature and applications of a stress appraisal and coping formulation. *Spinal Cord* 2001;39:615-27
 232. Elliot T, Kennedy P: Treatment of depression following spinal cord injury: An evidence-based review. *Rehabil Psychol* 2004;49:134-9
 233. Cardenas D, Warms C, Turner J, et al: Efficiency of amitriptyline for the relief of pain in spinal cord injury: Results of a randomized controlled trial. *Pain* 2002;96:365-73
 234. Kemp B, Kahan J, Krause J, et al: Treatment of major depression in individuals with spinal cord injury. *J Spinal Cord Med* 2004;27:22-8
 235. Kennedy P, Duff J, Evans M, et al: Coping effectiveness training reduces depression and anxiety following traumatic spinal cord injuries. *Br J Clin Psychol* 2003;42:41-52
 236. Diego M, Field T, Hernandez-Reif M, et al: Spinal cord patients benefit from massage therapy. *Int J Neurosci* 2002;112:133-42
 237. Hicks A, Martin K, Ditor D, et al: Long-term exercise training in persons with spinal cord injury: Effects on strength, arm ergometry performance, and psychological well-being. *Spinal Cord* 2003;41:34-43
 238. Kolakowsky-Hayner S, Gourley E, Kreutzer J, et al: Pre-injury substance abuse among person with brain injury and persons with spinal cord injury. *Brain Inj* 1999;13:571-81
 239. Turner A, Bombardier C, Rimmele C: A typology of alcohol use patterns among persons with recent traumatic brain injury or spinal cord injury: Implications for treatment matching. *Arch Phys Med Rehabil* 2003;84:358-64
 240. Levy D, Mallonee S, Miller J, et al: Alcohol involvement in burn, submersion, spinal cord, and brain injuries. *Med Sci Monitor* 2004;10:CR17-24
 241. Seltz M, Radnite C, Bauman W: The relationship of spinal cord injury trauma to alcohol misuse: A study of monozygotic twins. *J Spinal Cord Med* 2004;27:18-21
 242. Kolakowsky-Hayner S, Gourley E, Kreutzer J, et al: Post-injury substance abuse among persons with brain injury and persons with spinal cord injury. *Brain Inj* 2002;16:583-92
 243. Mc Kinley W, Kolakowsky S, Kreutzer J, et al: Substance abuse, violence, and outcome after traumatic spinal cord injury. *Am J Phys Med Rehabil* 1999;78:306-12
 244. Krause J, Coker J, Charlifue S, et al: Health outcomes among American Indians with spinal cord injuries. *Arch Phys Med Rehabil* 2000;81:924-31
 245. Elliot T, Kurylo M, Chen Y, et al: Alcohol abuse history and adjustment following spinal cord injury. *Rehabil Psychol* 2002;47:278-90
 246. Tate D, Forchheimer M, Krause J, et al: Patterns of alcohol and substance use and abuse in persons with spinal cord injury: Risk factors and correlates. *Arch Phys Med Rehabil* 2004;85:1837-49
 247. Bombardier C, Stroud M, Esselman P, et al: Do pre-injury alcohol problems predict poorer rehabilitation progress in person with spinal cord injury? *Arch Phys Med Rehabil* 2004;85:1488-92
 248. Gentilello L, Rivara F, Donovan D, et al: Alcohol interventions in a trauma center as a means of reducing the risk of injury recurrence. *Ann Surg* 1999;230:473-80
 249. Richards J, Kewman D, Pierce C: Spinal cord injury, in Frank R, Elliot T (eds): *Handbook of Rehabilitation Psychology*. American Psychological Association, pp 1127
 250. *Diagnostic and Statistical Manual of Mental Disorders*, ed 4. Washington, DC, American Psychiatric Association, 1994
 251. Kennedy P, Duff J: Post traumatic stress disorder and spinal cord injuries. *Spinal Cord* 2001;39:1-10

252. Kennedy P, Evans M: Evaluation of post traumatic distress in the first 6 months following SCI. *Spinal Cord* 2001;39:381-6
253. Benony H, Daloz L, Bungener C, et al: Emotional factors and subjective quality of life in subjects with spinal cord injuries. *Am J Phys Med Rehabil* 2002;81:437-45
254. Radnitz C, Schleini I, Hsu L: The effect of prior trauma exposure on the development of PTSD following spinal cord injury. *J Anxiety Disord* 2000;14:313-24
255. Boyer B, Knolls M, Kafkalas C, et al: Prevalence and relationships of posttraumatic stress in families experiencing pediatric spinal cord injury. *Rehabil Psychol* 2002;45:339-55
256. Danner G, Radnitz C: Predictive factors and posttraumatic stress disorder in veterans with spinal cord injury. *Int J Rehabil Health* 2000;5:195-203
257. Martz E: Death anxiety as a predictor of posttraumatic stress levels among individuals with spinal cord injuries. *Death Stud* 2004;28:1-17
258. Kreuter M: Spinal cord injury and partner relationships. *Spinal Cord* 2000;38:2-6
259. Putzke J, Elliot T, Richards J: Marital status and adjustment year post-spinal-cord-injury. *J Clin Psychol Med Settings* 2001;8:101-7
260. Lucke K, Coccia H, Goode J, et al: Quality of life in spinal cord injured individuals and their caregivers during the initial 6 months following rehabilitation. *Qual Life Res* 2004;13:97-110
261. Unalan H, Gencosmanoglu B, Akgun K, et al: Quality of life of primary caregivers of spinal cord injury survivors living in the community: Controlled study with Short Form 36 questionnaire. *Spinal Cord* 2001;39:318-22
262. Elliot T, Shewchuk R: Social problem-solving abilities and distress among family members assuming a caregiver role. *Br J Health Psychol* 2003;8:149-63
263. Elliot T, Shewchuk R, Richards J: Family caregiver social problem-solving abilities and adjustment during the initial year of the caregiving role. *J Counsel Psychol* 2001;48:223-32
264. Elliott T, Schewchuk R, Richards J: Caregiver social problem-solving abilities and family member adjustment to recent-onset physical disability. *Rehabil Psychol* 1999;44:104-23
265. Alexander C, Hwang K, Sipski M: Mothers with spinal cord injuries: Impact on marital, family and children's adjustment. *Arch Phys Med Rehabil* 2002;83:24-30
266. McColl M, Bickenbach J, Johnston J, et al: Spiritual issues associated with traumatic-onset disability. *Disabil Rehabil* 2000;22:655-64
267. McColl M, Bickenbach J, Johnston J, et al: Changes in spiritual beliefs after traumatic disability. *Arch Phys Med Rehabil* 2000;81:817-23
268. Tate D, Forchheimer M: Quality of life, life satisfaction, and spirituality: Comparing outcomes between rehabilitation and cancer patients. *Am J Phys Med Rehabil* 2002;81:400-10
269. Post M, Ros W, Schrijvers A: Impact of social support on health status and life satisfaction in people with a spinal cord injury. *Psychol Health* 1999;14:679-95
270. Rees T, Smith B, Sparkes A: The influence of social support on the lived experiences of spinal cord injured sportsmen. *Sport Psychol* 2003;17:135-56
271. Sherman J, DeVinney D, Sperling K: Social support and adjustment after spinal cord injury: Influence of past peer mentoring experiences. *Rehabil Psychol* 2004;49:140-9
272. Beringer A: Spinal cord injury and outdoor experiences. *Int J Rehabil Res* 2004;27:7-15
273. Dattilo J, Caldwell L, Lee Y, et al: Returning to the community with a spinal cord injury: Implications for therapeutic recreation specialists. *Ther Recreation J* 1998;32:13-27
274. Wise J, Hale S: Strength and generalizing self-efficacy in male with spinal cord injury. *Ther Recreation J* 1999;33:333-41
275. Sable J, Craig P, Lee D: Promoting health and wellness: A research-based case report. *Ther Recreation J* 2000;34:348-61
276. Batavia A, Batavia M: Karaoke for quads: A new application of an old recreation with potential therapeutic benefits for people with disabilities. *Disabil Rehabil* 2003;25:297-300
277. Slater D, Meade M: Participation in recreation and sports for persons with spinal cord injury: Review and recommendations. *NeuroRehabil* 2004;19:121-9
278. Hanson C, Nabavi D, Yuen H: The effect of sports on level of community integration as reported by persons with spinal cord injury. *Am J Occup Ther* 2001;55:332-8
279. Loy D, Dattilo J, Kleiber D: Exploring the influence of leisure on adjustment: Development of the leisure and spinal cord injury adjustment model. *Leisure Sci* 2003;25:231-55
280. O'Connor T, Cooper R, Fitzgerald S, et al: Evaluation of a manual wheelchair interface to computer games. *Neuro-rehabil Neural Repair* 2000;14:21-31
281. O'Connor T, Fitzgerald S, Cooper R, et al: Does computer game play aid in motivation of exercise and increase metabolic activity during wheelchair ergometry? *Med Eng Phys* 2001;23:267-73
282. Noreau L, Fougereyrollas P, Boschen K: Perceived influence of the environment on social participation among individuals with spinal cord injury. *Top Spinal Cord Inj Rehabil* 2002;7:56-72
283. Fougereyrollas P, Noreau L, Boschen K: Interaction of environment with individual characteristics and social participation: Theoretical perspectives and application in persons with spinal cord injury. *Top Spinal Cord Inj Rehabil* 2002;7:1-16
284. Noreau L, Dion S, Vacho J, et al: Productivity outcomes of individuals with spinal cord injury. *Spinal Cord* 1999;37:730-6
285. Brown M, Gordon W, Spielman L, et al: Participation by individuals with spinal cord injury in social and recreational activity outside the home. *Top Spinal Cord Inj Rehabil* 2002;7:83-100
286. Fiedler I, Indermuehler D, Drobac W, et al: Perceived barriers to employment in individuals with spinal cord injury. *Top Spinal Cord Inj Rehabil* 2002;7:73-82
287. Carlson D, Myklebust J: Wheelchair use and social integration. *Top Spinal Cord Inj Rehabil* 2002;7:28-46
288. Roach M: Community social structure as an indicator of social integration and its effect on quality of life for persons with spinal cord injury. *Top Spinal Cord Inj Rehabil* 2002;7:101-11
289. Bockian N, Lee A, Fidanque C: Personality disorders and spinal cord injury: A pilot study. *J Clin Psychol in Med Settings* 2003;10:307-13
290. Krause J: Factors associated with risk for subsequent injuries after traumatic spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1503-8
291. Buchanan K, Elias L: Psychological distress and family burden following spinal cord injury: Concurrent traumatic brain injury cannot be overlooked. *Axone* 2001;22:16-7
292. Macciocchi S, Bowman B, Coker J, et al: Effect of comorbid brain injury on functional outcome of persons with spinal cord injuries. *Am J Phys Med Rehabil* 2004;83:22-6
293. Hanks RA, Rapport LJ, Millis SSR, et al: Measures of executive functioning as predictors of functional ability and social integration in the rehabilitation sample. *Arch Phys Med Rehabil* 1999;80:1030-7

294. Galvin LR, Godfrey HP: The impact on going on emotional adjustment to spinal cord injury (SCI): Review of the literature and application of a stress appraisal and coping formulation. *Spinal Cord* 2001;39:615-27
295. Kennedy P, Marsh N, Lowe R, et al: A longitudinal analysis of psychological impact and coping strategies following spinal cord injury. *Br J Health Psychol* 2000;5:157-72
296. Craig A, Hancock K, Dichson H: Improving the long-term adjustment of spinal cord injured persons. *Spinal Cord* 1999;37:345-50
297. Kennedy P, Duff J, Evans M, et al: Coping effectiveness training reduces depression and anxiety following traumatic spinal cord injuries. *Br J Clin Psychol* 2003;42:41-52
298. Dreer L, Elliot T, Tucker E: Social problem solving abilities and health behaviors among persons with recent-onset spinal cord injury. *J Clin Psychol Med Settings* 2004;11:7-13
299. Elliot T: Social problem solving abilities and adjustment to recent-onset spinal cord injury. *Rehabil Psychol* 1999;44:315-32
300. Lohne Vand Severinsson E: Hope during the first months after acute spinal cord injury. *J Adv Nurs* 2004;47:279-86
301. Lohne V: Hope in patients with spinal cord injury: A literature review related to nursing. *J Neurosci Nurs* 2001;33:317-25
302. Lohne Vand Severinsson E: Hope and despair: The awakening of hope following acute spinal cord injury-an interpretative study. *Int J Nurs Stud* 2004;41:881-90
303. Thompson N, Coker J, Krause J, et al: Purpose in life as a mediator of adjustment after spinal cord injury. *Rehabil Psychol* 2003;48:100-8
304. McMille J, Cook C: The positive by-products of spinal cord injury and their correlates. *Rehabil Psychol* 2003;48:77-85
305. Zemper E, Tate D, Roller S, et al: Assessment of a holistic wellness program for persons with spinal cord injury. *Am J Phys Med Rehabil* 2003;12:957-68
306. Dijkers M: Quality of life of individuals with spinal cord injury: Conceptualization, measurement, and research findings, in Third International Congress Restorative of (Wheeled) Mobility in SCI Rehabilitation: State of the Art III, April 19, 2004
307. Hammell K: Exploring quality of life following high spinal cord injury: A review and critique. *Spinal Cord* 2004;42:491-502
308. Manns P, Chad K: Components of quality of life for persons with a quadriplegic and paraplegic spinal cord injury. *Qual Health Res* 2001;11:795-811
309. Leduc B, Lepage Y: Health-related quality of life after spinal cord injury. *Disabil Rehabil* 2002;24:196-202
310. Holicky R, Charlifue S: Ageing with spinal cord injury: The impact of spousal support. *Disabil Rehabil* 1999;21:250-7
311. Anderson C, Krajci K, Vogel L: Life satisfaction in adults with pediatric-onset spinal cord injuries. *J Spinal Cord Med* 2002;25:184-90
312. Price G, Kendall M, Amsters D, et al: Perceived causes of change in function and quality of life for people with long duration spinal cord injury. *Clin Rehabil* 2004;18:164-71
313. Benony H, Daloz L, Bungerner C, et al: Emotional factors and subjective quality of life in subjects with spinal cord injuries. *Am J Phys Med Rehabil* 2002;81:437-45
314. Hicken B, Putzke J, Richards J: Bladder management and quality of life after spinal cord injury. *Am J Phys Med Rehabil* 2001;80:916-22
315. McColl M: Expectations of health, independence, and quality of life among aging spinal-injured adults. *Assist Technol* 1999;11:130-6
316. Putzke J, Elliott T, Richards J: Marital status and adjustment year post-spinal-cord-injury. *J Clin Psychol Med Settings* 2001;8:101-7
317. Hicken B, Putzke J, Novack T, et al: Life satisfaction following spinal cord and traumatic brain injury: A comparative study. *J Rehabil Res Dev* 2002;39:359-65
318. Krause J, Broderick L, Broyles J: Subjective well-being among African-Americans with spinal cord injury: An exploratory study between men and women. *NeuroRehabil* 2004;19:81-9
319. Dijkers M: Correlates of life satisfaction among persons with spinal cord injury. *Arch Phys Med Rehabil* 1999;80:867-76
320. Richards J, Bombardier C, Tate D, et al: Access to the environment and life satisfaction after spinal cord injury. *Arch Phys Med Rehabil* 1999;80:1501-71
321. Manns P, Chad K: Determining the relation between quality of life, handicap, fitness, and physical activity for persons with spinal cord injury. *Arch Phys Med Rehabil* 1999;80:1566-71
322. McColl M, Stirling P, Walker J, et al: Expectations of independence and life satisfaction among ageing spinal cord injured adults. *Disabil Rehabil* 1999;21:231-40
323. Randell N, Lynch A, Anthony A, et al: Does a colostomy alter quality of life in patients with spinal cord injury? A controlled study. *Spinal Cord* 2001;39:279-82
324. Anderson C, Vogel L: Domain-specific satisfaction in adults with pediatric-onset spinal cord injuries. *Spinal Cord* 2003;41:684-91
325. Dowler R, Richards J, Putzke J, et al: Impact of demographic and medical factors on satisfaction with life after spinal cord injury: A normative study. *J Spinal Cord Med* 2001;24:87-91
326. Hulting C, Giuliano F, Quirk F, et al: Quality of life in patients with spinal cord injury receiving Viagra (sildenafil citrate) for the treatment of erectile dysfunction. *Spinal Cord* 2000;38:363-70
327. Hicks A, Martin K, Ditor D, et al: Long-term exercise training in persons with spinal cord injury: Effects on strength, arm ergometry performance, and psychological well-being. *Spinal Cord* 2003;41:34-43
328. Johnston M, Nissim E, Wood K, et al: Objective and subjective handicap following spinal cord injury: Interrelationships and predictors. *J Spinal Cord Med* 2002;25:11-22
329. Gerhart K, Weitzenkamp D, Kennedy P, et al: Correlates of stress in long-term spinal cord injury. *Spinal Cord* 1999;37:183-90
330. Kennedy P, Rogers B: Reported quality of life of people with spinal cord injuries: A longitudinal analysis of the first 6 months post-discharge. *Spinal Cord* 2000;38:498-503
331. Lucke K, Coccia H, Goode J, et al: Quality of life in spinal cord injured individuals and their caregivers during the initial 6 months following rehabilitation. *Qual Life Res* 2004;13:97-110
332. Putzke J, Richards J, DeVivo M: Quality of life after spinal cord injury caused by gunshot. *Arch Phys Med Rehabil* 2001;82:949-54
333. Putzke J, Richards J, Hicken B, et al: Interference due to pain following spinal cord injury: Important predictors and impact on quality of life. *Pain* 2002;100:231-42
334. Putzke J, Richards J, Hicken B, et al: Predictors of life satisfaction: A spinal cord injury cohort study. *Arch Phys Med Rehabil* 2002;83:555-61
335. Krause J: Years to employment after spinal cord injury. *Arch Phys Med Rehabil* 2003;84:1282-9
336. Tomassen P, Post M, van Asbeck F: Return to work after spinal cord injury. *Spinal Cord* 2000;38:51-5
337. Meade M, Lewis A, Jackson M, et al: Race, employment,

- and spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1782–92
338. Murphy G, Young A, Brown D, et al: Explaining labor force status following spinal cord injury: The contribution of psychological variables. *J Rehabil Med* 2003;35:276–83
339. Anderson C, Vogel L: Work experience in adolescents with spinal cord injuries. *Dev Med Child Neurol* 2000;42:515–7
340. Anderson C, Vogel L: Employment outcomes of adults who sustained spinal cord injuries as children or adolescents. *Arch Phys Med Rehabil* 2002;83:791–801
341. Conroy L, McKenna K: Vocational outcomes following spinal cord injury. *Spinal Cord* 1999;37:624–33
342. Hess D, Meade M, Forcheimer M, et al: Psychological well-being and intensity of employment in individuals with a spinal cord injury. *Top Spinal Cord Inj Rehabil* 2004;9:1–10
343. Yuen H, Burik J, Krause J: Physical and psychological well-being among adults with spinal cord injury: The role of volunteer activities. *Top Spinal Cord Inj Rehabil* 2004;9:19–25
344. Krause J, Kewman D, DeVivo M, et al: Employment after spinal cord injury: An analysis of cases from the Model Spinal Cord Injury Systems. *Arch Phys Med Rehabil* 1999;80:1492–1500
345. Schonherr M, Groothoff J, Mulder G, et al: Vocational outcomes following spinal cord injury: Expectations, participation, and interventions. *Spinal Cord* 2004;42:177–84
346. King T, Emery R, Warren S, et al: A collaborative approach to returning clients to work during the first year after spinal cord injury. *Top SCI Rehabil* 2004;9:33–42
347. Malassigne P, Nelson A, Cors A, et al: Iterative design and evaluation of new prone carts for individuals with SCDs: A technical note. *J Rehabil Res Dev* 2002;39:127–39
348. Chen Y, Chen S, Chen W, et al: A head oriented wheelchair for people with disabilities. *Disabil Rehabil* 2003;25:249–53
349. Malassigne P, Nelson A, Cors M, et al: Design of the advanced commode-shower chair for spinal cord injured individuals. *J Rehabil Res Dev* 2000;37:373–82
350. Cooper RA, Boninger M, Cooper R, et al: Preliminary assessment of a prototype advanced mobility device in the work environment of veterans with spinal cord injury. *NeuroRehabil* 2004;19:161–70
351. Cooper RA, Boninger M, Cooper R, et al: Use of the independence 3000 IBOT transported at home and in the community. *J Spinal Cord Med* 2003;26:79–85
352. Birch G, Bozorgzadeh Z, Mason S: Initial on-line evaluations of the LF-ASD brain computer interface with able-bodied and spinal-cord subjects using imagined voluntary motor potentials. *IEEE Trans Neural Syst Rehabil Eng* 2002;10:219–24
353. Wolpaw J, Birbaumer N, McFarland D, et al: Brain-computer interfaces for communication and control. *Clin Neurophysiol* 2002;113:767–91
354. Craig A, Moses P, Tran Y, et al: The effectiveness of hands-free environmental control system for the profoundly disabled. *Arch Phys Med Rehabil* 2002;83:1455–8
355. Algood S, Cooper R, Fitzgerald et al: Impact of pushrim-activated power-assisted wheelchair on the metabolic demands, stroke frequency, and range of motion among subjects with tetraplegia. *Arch Phys Med Rehabil* 2004;85:1865–71
356. Consortium for Spinal Cord Medicine: Preservation of upper limb function following spinal cord injury: A clinical practice guideline for health-care professionals. Washington, DC, Paralyzed Veterans of America, 2005, pp 1–29
357. Newsam CJ, Rao SS, Mulroy S, et al: Three dimensional upper extremity motion during manual wheelchair propulsion in men with different levels of spinal cord injury. *Gait Posture* 1999;10:223–32
358. Boninger M, Baldwin M, Cooper R, et al: Manual wheelchair pushrim biomechanics and axle position. *Arch Phys Med Rehabil* 2000;81:608–13
359. Boninger M, Cooper R, Baldwin M, et al: Wheelchair pushrim kinetics: Body weight and median nerve function. *Arch Phys Med Rehabil* 1999;80:910–5
360. Boninger M, Souza A, Cooper R, et al: Propulsion patterns and pushrim biomechanics in manual wheelchair propulsion. *Arch Phys Med Rehabil* 2002;83:718–23
361. Beekman C, Miller-Porter L, Schoneberger M: Energy cost of propulsion in standard and ultralight wheelchairs in people with spinal cord injuries. *Phys Ther* 1999;79:146–58
362. Cooper R, Fitzgerald S, Boninger M, et al: Evaluation of a pushrim-activated, power assisted wheelchair. *Arch Phys Med Rehabil* 2001;82:702–10
363. Harvey L, Crosbie J: Weight bearing through flexed upper limbs in quadriplegics with paralyzed triceps brachii muscles. *Spinal Cord* 1999;37:780–5
364. Harvey L, Crosbie J: Biomechanical analysis of a weight-relief maneuver in C5 and C6 quadriplegia. *Arch Phys Med Rehabil* 2000;81:500–5
365. Gagnon D, Nadeau S, Gravel D, et al: Biomechanical analysis of a posterior transfer maneuver on a level surface in individuals with high and low level spinal cord injuries. *Clin Biomech* 2003;18:319–31
366. Fitzgerald S, Cooper R, Boninger M, et al: Comparison of fatigue life for 3 types of manual wheelchairs. *Arch Phys Med Rehabil* 2001;82:1484–8
367. Chaves E, Boninger M, Cooper R, et al: Assessing the influence of wheelchair technology on perception of participation in spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1854–79
368. Vesmarovich S, Walker T, Hauber RP, et al: Use of telerehabilitation to manage pressure ulcers in persons with spinal cord injury. *Adv Wound Care* 1999;12:164–9
369. Drainoni M, Houlihan B, Williams S, et al: Patterns of internet use by persons with spinal cord injuries and relationships to health-related quality of life. *Arch Phys Med Rehabil* 2004;85:1872–79
370. Bingham S, Beatty P: Rates of access to assistive equipment and medical rehabilitation services among people with disabilities. *Disabil Rehabil* 2003;25:487–90
371. Hunt P, Boninger M, Cooper R, et al: Demographic and socioeconomic factors associated with disparity in wheelchair customizability among people with traumatic spinal cord injury. *Arch Phys Med Rehabil* 2004;85:1859–64
372. Weitzenkamp D, Jones R, Whiteneck G, et al: Aging with spinal cord injury: Cross-sectional and longitudinal effects. *Spinal Cord* 2001;39:301–9
373. Krause J: Aging after spinal cord injury: An exploratory study. *Spinal Cord* 2000;38:77–83
374. Charlifue S, Weitzenkamp D, Whiteneck G: Longitudinal outcomes in spinal cord injury: Aging, secondary conditions, and well-being. *Arch Phys Med Rehabil* 1999;80:1429–34
375. Krause J, Kemp B, Coker J: Depression after spinal cord injury: Relations to gender, ethnicity, aging, and socioeconomic indicators. *Arch Phys Med Rehabil* 2000;81:1099–109
376. Pentland W, Walker J, Minnes P, et al: Women with spinal cord injury and the impact of aging. *Spinal Cord* 2000;40:374–87
377. McColl M: A house of cards: Women, aging, and spinal cord injury (editorial). *Spinal Cord* 2002;40:371–3

378. Groah S, Stiens S, Gittler M, et al: Spinal cord injury medicine: Preserving wellness and independence of the aging patient with spinal cord injury. A primary care approach for the rehabilitation medicine specialist. *Arch Phys Med Rehabil* 2004;83:S82-9
379. Putzke J, Barrett J, Richards J, et al: Age and spinal cord injury: An emphasis on outcomes among the elderly. *J Spinal Cord Med* 2003;26:37-44
380. Barrett J, Putzke J, Richards J: Early versus late onset of spinal cord injury among the elderly. *J Clin Psychol Med Settings* 2002;9:219-26
381. Thompson L: Functional changes in person aging with spinal cord injury. *Assist Technol* 1999;11:123-9
382. McColl M, Charlifue S, Glass C, et al: Aging, gender and spinal cord injury. *Arch Phys Med Rehabil* 2004;58:363-7
383. McColl M, Stirling P, Walker J, et al: Expectations of independence and life satisfaction among ageing spinal cord injured adults. *Disabil Rehabil* 1999;21:231-40
384. McColl M, Arnold R, Charlifue S, et al: Aging, spinal cord injury, and quality of life: Structural relationships. *Arch Phys Med Rehabil* 2003;84:1137-44
385. Thompson L: Functional changes in persons aging with spinal cord injury *Assist Technol* 1999;11:123-9
386. Charlifue S, Gerhart K: Changing psychosocial morbidity in people aging with spinal cord injury. *NeuroRehabil* 2004;19:15-23