

SIR models for the spread of COVID-19

Maksim Gordienko

Alexander Sysoev

January 9, 2021

1 Introduction

In this article we are going to show how Kermack and McKendrick SIR models can be used for epidemic process simulation. Then we add vital dynamics to SIR model. Next we solve simple models in analytical way. Finally, we introduce sophisticated SEIRS-like model especially for COVID-19 conditions, close to real life.

2 Mathematical model of epidemics

Let's consider a group of N people and classify them into these types:

- **S** – Susceptible
- **I** – Infected
- **R** – Recovered

This is called the SIR model for the spread of epidemic diseases that describes the changes in numbers of the three types of individuals. We denote the number of susceptible persons by $s(t)$, the number of infected individuals by $i(t)$ and the number of recovered people by $r(t)$. The time t is measured in days. Also we suppose that each person is in contact with m persons per day average. Hence the number of contacts between the susceptible and infected people becomes $\frac{m}{N}i(t)s(t)$. If we set the probability of infection for each contact as p then the number of newly infected individuals within Δt days becomes

$$\frac{m}{N}i(t)s(t)p\Delta t \quad (1)$$

in total. Let $\beta = mp$, the number of non-infected people (**S**) from the t -th day to $(t + \Delta t)$ -th day changes as

$$s(t + \Delta t) - s(t) = -\frac{\beta}{N}s(t)i(t)\Delta t \quad (2)$$

When $\Delta t \rightarrow 0$, we can rewrite it as differential equation

$$\frac{ds(t)}{dt} = -\frac{\beta}{N}s(t)i(t) \quad (3)$$

Meanwhile, the infected individuals recover at a removal rate γ per day. Subsequently, increase in the number of the recovered persons becomes

$$\frac{dr(t)}{dt} = \gamma i(t) \quad (4)$$

Respectively, $\frac{1}{\gamma}$ is the expected duration of infection. Also the number of recovered people includes the amount of deceased persons because they cannot possibly infect others.

As well as the total number of individuals is N , we can express $i(t)$

$$i(t) = N - s(t) - r(t) \quad (5)$$

The change in the number of infected people can be written as

$$\frac{di(t)}{dt} = -\frac{ds(t)}{dt} - \frac{dr(t)}{dt} = \frac{\beta}{N}s(t)i(t) - \gamma i(t) = \frac{\beta}{N}(s(t) - \gamma)i(t) \quad (6)$$

Gathering the equations above, we have

$$\begin{cases} \frac{ds(t)}{dt} = -\frac{\beta}{N}s(t)i(t) \end{cases} \quad (7)$$

$$\begin{cases} \frac{di(t)}{dt} = (\frac{\beta}{N}s(t) - \gamma)i(t) \end{cases} \quad (8)$$

$$\begin{cases} \frac{dr(t)}{dt} = \gamma i(t) \end{cases} \quad (9)$$

As initial conditions, we use

$$\begin{cases} s(0) = N_1 \\ i(0) = N_2 \\ r(0) = 0 \end{cases} \quad (10)$$

If we introduce the following transformations of variables

$$\tilde{s}(t) = \frac{s(t)}{N}, \tilde{i}(t) = \frac{i(t)}{N}, \tilde{r}(t) = \frac{r(t)}{N}, \tilde{t} = \beta t$$

Then system of the equations becomes

$$\begin{cases} \frac{d\tilde{s}(t)}{d\tilde{t}} = -\tilde{s}(t)\tilde{i}(t) \\ \frac{d\tilde{i}(t)}{d\tilde{t}} = (\tilde{s}(t) - \frac{1}{R_0})\tilde{i}(t) \\ \frac{d\tilde{r}(t)}{d\tilde{t}} = \frac{1}{R_0}\tilde{i}(t) \end{cases} \quad (11)$$

The number $R_0 = \frac{\beta}{\gamma}$ is known as the basic reproduction number. The number of infected people increases when $R_0 > 1$ and decreases when $R_0 < 1$.

3 Analytical solution of SIR model

When model is defined, we can solve the system of equations (7) – (9). For simplicity forget about function arguments. Firstly, rewrite the equation (7) as

$$i = -\frac{1}{\tilde{\beta}}\left(\frac{s'}{s}\right) \quad (12)$$

where $\tilde{\beta} = \frac{\beta}{N}$. Then differentiate both sides

$$i' = -\frac{1}{\tilde{\beta}}\left(-\frac{s'^2}{s^2} + \frac{s''}{s}\right) \quad (13)$$

Next insert the equation (12) into (8)

$$i' = -(\tilde{\beta}s - \gamma)\frac{1}{\tilde{\beta}}\left(\frac{s'}{s}\right) \quad (14)$$

Comparing equations (13) and (14) we have

$$s \frac{d^2 s}{dt^2} - \left(\frac{ds}{dt} \right)^2 + (\gamma - \tilde{\beta} s) s \frac{ds}{dt} = 0 \quad (15)$$

Now we introduce the following function

$$\phi = \frac{dt}{ds} \quad (16)$$

Afterward (15) becomes

$$\frac{d\phi}{ds} + \frac{\phi}{s} = (\gamma - \tilde{\beta} s) \phi^2 \quad (17)$$

This is a Bernoulli differential equation. Divide both parts by ϕ^2

$$\frac{\phi'}{\phi^2} + \frac{1}{\phi s} = \gamma - \tilde{\beta} s \quad (18)$$

Then make a substitution

$$z = \frac{1}{\phi}, z' = -\frac{\phi'}{\phi^2} \quad (19)$$

Now solve the first-order linear ordinary equation

$$-z' + \frac{z}{s} = \gamma - \tilde{\beta} s \quad (20)$$

with general solution where C is constant

$$z = -\gamma s \ln s + \tilde{\beta} s^2 + Cs \quad (21)$$

Returning back to ϕ

$$\phi = \frac{1}{s(C - \gamma \ln s + \tilde{\beta} s)} \quad (22)$$

From the relation of inverse function in equation (16), we have

$$\frac{1}{\phi} = \frac{ds}{dt} \quad (23)$$

Using equation (7), we obtain

$$i(t) = -\frac{1}{\tilde{\beta}} (C - \gamma \ln s(t) + \tilde{\beta} s(t)) \quad (24)$$

Moreover, from equations (7) and (9), we get

$$\frac{dr}{dt} = -\frac{\gamma}{\tilde{\beta}} \left(\frac{s'}{s} \right) \quad (25)$$

Subsequently, the relation between $s(t)$ and $r(t)$ becomes

$$r(t) = -\frac{\gamma}{\tilde{\beta}} \ln \frac{s(t)}{C_1} \quad (26)$$

where C_1 is a constant. According to our initial conditions (10)

$$C_1 = N_1 \quad (27)$$

From the relation $s(0) + i(0) + r(0) = N$, we have

$$C = -\tilde{\beta}N + \gamma \ln N_1 \quad (28)$$

If we substitute C into equation (22), we obtain

$$\frac{dt}{ds} = \frac{1}{s(-\tilde{\beta}N - \gamma \ln \frac{s}{N_1} + \tilde{\beta}s)} \quad (29)$$

Integrate this and express t as a function of s

$$t = \int_{s(0)}^{s(t)} \frac{d\varepsilon}{\varepsilon(-\tilde{\beta}N - \gamma \ln \frac{\varepsilon}{N_1} + \tilde{\beta}\varepsilon)} \quad (30)$$

For convenience, we change a variable

$$\xi = \frac{\varepsilon}{N_1} \quad (31)$$

Rewrite the equation (30)

$$t(s) = \int_1^{\frac{s(t)}{N_1}} \frac{d\xi}{\xi(-\beta - \gamma \ln \xi + \beta \xi \frac{N_1}{N})} \quad (32)$$

Now we can calculate $t(s)$ using numerical integration with a small step size and $s(t)$ as a parameter like that

$$\int_a^b f(\xi) d\xi \simeq \sum_{i=0}^{n-1} f(a + ih)h, \text{ where } h = \frac{b-a}{n} \quad (33)$$

If $t(s)$ is obtained, then $i(t)$ and $r(t)$ can be calculated from equations (24) and (26) respectively.

4 SIR Model with natural deaths and births

Let's rewrite our system of equations (7) – (9) with an addition of the death and birth processes. Consider these rates are equal to each other, we introduce new variable D :

$$\begin{cases} \frac{ds(t)}{dt} = -\frac{\beta}{N}s(t)i(t) + D(N - s(t)) \end{cases} \quad (34)$$

$$\begin{cases} \frac{di(t)}{dt} = (\frac{\beta}{N}s(t) - \gamma - D)i(t) \end{cases} \quad (35)$$

$$\begin{cases} \frac{dr(t)}{dt} = \gamma i(t) - Dr(t) \end{cases} \quad (36)$$

As we did before, we rewrite it into one equation:

$$i = \frac{s' - D(N - s)}{-\tilde{\beta}s} \quad (37)$$

Again we differentiate both sides (with $\tilde{\beta} = \frac{\beta}{N}$ substitution):

$$i' = -\frac{1}{\tilde{\beta}} \left(\frac{(s'' + Ds')s - (s' + Ds - DN)s'}{s^2} \right), \quad (38)$$

$$i' = -\frac{1}{\tilde{\beta}} \left(\frac{s''}{s} - \frac{s'^2 - DN s'}{s^2} \right) \quad (39)$$

Now we can insert (37) into (35):

$$i' = (\tilde{\beta}s - \gamma - D) \left(\frac{s' - D(N - s)}{-\tilde{\beta}s} \right) \quad (40)$$

$$i' = -\frac{1}{\tilde{\beta}s} (\tilde{\beta}s - \gamma - D)(s' + Ds - DN) \quad (41)$$

And comparing (41) and (39) we get next equation:

$$ss'' - s'^2 + (-\tilde{\beta}s^2 + (\gamma + D)s + DN)s' - \tilde{\beta}Ds^3 - (\tilde{\beta}DN + D(\gamma + D))s^2 - (\gamma + D)DNs = 0 \quad (42)$$

We can see that coefficients at s^3 , s^2 and s are *consts* so we substitute them with $A = -\tilde{\beta}D$, $B = -(\tilde{\beta}DN + D(\gamma + D))$, $C = -(\gamma + D)DN$:

$$ss'' - s'^2 + (-\tilde{\beta}s^2 + (\gamma + D)s + DN)s' + As^3 + Bs^2 + Cs = 0 \quad (43)$$

Now we introduce a function $\phi = \frac{dt}{ds}$, so the equation will be:

$$-\phi' - \frac{\phi}{s} + (-\tilde{\beta}s^2 + (\gamma + D)s + DN)\frac{\phi^2}{s} + (As^2 + Bs + C)\phi^3 = 0 \quad (44)$$

And by substitution $v = \phi s$, $v' = \phi's + \phi$ the following equation occur:

$$v' + (-\tilde{\beta} + \frac{(\gamma + D)}{s} + \frac{DN}{s^2})v^2 + (A + \frac{B}{s} + \frac{C}{s^2})v^3 = 0 \quad (45)$$

Then we assume these three substitutions:

$$\left\{ \begin{array}{l} r(s) = \frac{1}{v(s)}, \end{array} \right. \quad (46)$$

$$\left\{ \begin{array}{l} \xi(s) = (\tilde{\beta} - \frac{(\gamma + D)}{s} - \frac{DN}{s^2}), \end{array} \right. \quad (47)$$

$$\left\{ \begin{array}{l} \eta(s) = -(A + \frac{B}{s} + \frac{C}{s^2}) \end{array} \right. \quad (48)$$

and we get Abel-type first-order second-kind equation:

$$rr' = \xi(s)r + \eta(s) \quad (49)$$

5 Modeling COVID-19 pandemic

5.1 Definitions

In order to create lifelike COVID-19 infection model we need to introduce more parameters in our model and consider different spread cases.

- S – Susceptible. Includes all individuals who are not infected but are susceptible to contract the disease.
 - E – Exposed. Includes all persons who are exposed to infection, but not yet infections. Some of them might fall ill and some may not. We introduced that component due to incubation period of COVID-19 pandemic (approximated 5-6 days by WHO).
 - I_s – Symptomatic infected. It includes all the individuals who were symptomatic and infectious. They have approached some health-care facility, but have not yet been quarantined. This compartment was brought into the picture considering concerning news from worst-affected counties like Italy and the U.S. of the hospitals and health-care centers getting filled up very fast during this pandemic. A significant portion of the infected individuals may not be quarantined in case of the health-care system collapses.
 - I_{as} – Asymptomatic infected. It includes all those people who are affected but asymptomatic before they either recover or die or get permanently disabled. This compartment was formalized keeping after reports of various authentic studies claiming that around 30-40 percent of the infected individuals remain asymptomatic.
 - Q – Quarantined. Includes all persons who are currently under quarantine in health-care facility.
 - Q' – In intensive care unit. Includes the quarantined patients who had to be moved to ICU after their condition worsened.
 - C – Carrier. This compartment includes individuals who have left quarantine after being tested negative but actually have not fully healed. So they can possibly infect other susceptible individuals. They eventually either fall sick again or recover from disease. We introduced it due to multiple cases of re-infection being reported from countries like South Korea, China and Japan. It is safe to assume that one of the following happened:
 1. Medical inefficiency (inaccurate test result)
 2. Loss of immunity after recovery and subsequent re-infection
- For the first possibility we introduced the C compartment. To account for the second possibility we have kept the transition from R to S compartment respectively.
- R_{wd} – Recovered without disability. It includes all individuals who have recovered from infection without any disability and they can no longer infect any other individual
 - D – Deceased. Includes all deaths during the pandemic.
 - R_d – Recovered with disability. In this compartment we have kept all people who have recovered from the infection, can no longer infect anyone else but have been permanently disabled post recovery.

5.2 Formualting differential equations

Now we can define differential equations corresponding to described model above.

$$\frac{dS}{dt} = -\alpha \frac{S(I_s + I_{as} + C)}{N} + gR_{wd} \quad (50)$$

Where α is the disease transmission rate (same as $\beta = mp$ in SIR model) and g is the rate at which a fraction of recovered individuals lose their immunity.

$$\frac{dE}{dt} = \alpha \frac{S(I_s + I_{as} + C)}{N} - \mu E \quad (51)$$

If average incubation period is taken to be an exponential distribution μ .

$$\frac{dI_s}{dt} = r\mu E - \varepsilon I_s + fC - \zeta_1 I_s - \eta_1 I_s \quad (52)$$

Where f is rate at which a fraction of carriers gets re-infected. Here $0 \leq r \leq 1$ is a number which shows how many individuals in the exposed compartment moves to I_s compartment rather than I_{as} one. And $\varepsilon, \zeta_1, \eta_1$ are the rates at which infected individuals get quarantined, deceased, disabled respectively.

$$\frac{dI_{as}}{dt} = (1-r)\mu E - \beta_3 I_{as} - \zeta_3 I_{as} - \eta_3 I_{as} \quad (53)$$

Where β_3, ζ_3 and η_3 are the recovery rate, death rate and disability rate of asymptomatic individuals respectively.

$$\frac{dQ}{dt} = \varepsilon I_s - \beta_1 Q - vQ - pQ - \zeta_2 Q - \eta_2 Q \quad (54)$$

Where v, p, ζ_2 and η_2 are the rates at which the quarantined individuals go to carrier state, ICU, deceased and disabled compartments respectively. β_1 is the recovery rate for quarantined people.

$$\frac{dQ'}{dt} = pQ - \beta_4 Q' - \zeta_5 Q' - \eta_5 Q' \quad (55)$$

Where β_4, ζ_5 and η_5 are the rate with which individuals in ICU recover, die and get disabled respectively.

$$\frac{dC}{dt} = vQ - fC - \beta_2 C - \zeta_4 C - \eta_4 C \quad (56)$$

Where β_2, ζ_4 and η_4 are the rate with which carrier individuals recover, die and get disabled silently, respectively.

$$\frac{dR_{wd}}{dt} = \beta_1 Q + \beta_3 I_{as} + \beta_2 C - gR_{wd} \quad (57)$$

$$\frac{dD}{dt} = \zeta_1 I_s + \zeta_2 Q + \zeta_3 I_{as} + \zeta_4 C + \zeta_5 Q' \quad (58)$$

$$\frac{dR_d}{dt} = \eta_1 I_s + \eta_2 Q + \eta_3 I_{as} + \eta_4 C + \eta_5 Q' \quad (59)$$

System of these equations can be solved using numerical methods.

5.3 Parameters selection

Table 1: Initial condtions

Parameter	Value
N	83783945
S	83689432
E	45061
I_s	67123
I_{as}	11410.91
Q	12753.37
Q'	2528
C	5610
R_{wd}	19800
D	1584
R_d	13200

Table 2: Other parameters

Parameter	Value	Parameter	Value
α	0.8	v	0.05
g	0.0001	p	0.25
μ	7	ζ_2	0.125
f	0.05	η_2	0.6
r	0.14	β_1	0.1
ε	0.2	β_4	0.05
ζ_1	0.021	ζ_5	0.5
η_1	0.1977	η_5	0.45
β_3	0.1428	β_2	0.25
ζ_3	0.003	ζ_4	0.05
η_3	0.057	η_4	0.01

We decided to use Germany as an example for modeling epidemics. These values were actual on April 5, 2020. And we take into account further statistics to fine tune the parameters to real life.

5.4 Obtained result

placeholder

5.5 Comparing with real life situation

placeholder

6 References and data sources

- Exact analytical solutions of the Susceptible-Infected-Recovered (SIR) epidemic model and of the SIR model with equal death and birth rates. Tiberiu Harko, Francisco S. N. Lobo, M. K. Mak.

- A Mathematical Model of Epidemics—A Tutorial for Students. Yutaka Okabe and Akira Shudo. Department of Physics, Tokyo Metropolitan University, Hachioji, Tokyo 192-0397, Japan.
- Modelling and simulation of COVID-19 propagation in a large population with specific reference to India. Ashish Menon, Nithin K Rajendran, Anish Chandrachud, Girish Setlur. Department of Physics, Indian Institute of Technology Guwahati.
- Statistics of the COVID-19 pandemic in Germany: https://en.wikipedia.org/wiki/Statistics_of_the_COVID-19_pandemic_in_Germany
- Close to 17 percent of patients recovered from COVID-19 could still carry virus: <https://www.sciencedaily.com/releases/2020/10/201028134029.htm>
- Flattening the disability curve: Rehabilitation and recovery after COVID-19 infection: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7211743/>
- Complete COVID-19 dataset maintained by Our World in Data: <https://github.com/owid/covid-19-data/tree/master/public/data/>