

The Representer Theorem for Kernel Based Support Vector Machines

Aaron Avram

May 27 2025

Introduction

I will outline a proof of the representer theorem for reproducing kernel Hilbert spaces over $\mathbb{R}$, and then will use it in conjunction with the Support Vector Machine methods I have developed thus far to create a kernel based support vector machine.

Lemma

First I will prove a lemma:

A Hilbert space $\mathcal{H}$ of functions from a set X to $\mathbb{R}$ (or $\mathbb{C}$) is a reproducing kernel Hilbert space if and only if $\exists K_x \in \mathcal{H}$ such that $\forall f \in \mathcal{H} \langle f, K_x \rangle = f(x)$.

Proof. $\implies$ By definition a Hilbert space H is a reproducing kernel Hilbert space if $\forall x \in X$ the linear functional $L_x : \mathcal{H} \rightarrow \mathbb{R}$ given by $L_x(f) = f(x)$ is continuous. Thus L_x is bounded on $\mathcal{H}$ as continuity implies boundedness, so we can apply Riesz's representation theorem to find a unique $g_x \in \mathcal{H}$ such that $L_x(f) = \langle f, g_x \rangle$. Setting $K_x = g_x$, it is clear that K_x is well defined as g_x is unique by Riesz's and $\langle f, K_x \rangle = \langle f, g_x \rangle = f(x)$. Thus, K_x is a reproducing kernel of $\mathcal{H}$.

$\Leftarrow$ Suppose there exists a $K_x \in \mathcal{H}$ such that $\langle f, K_x \rangle = f(x)$. Then we can define a linear functional $L_x : X \rightarrow \mathbb{R}$ $L_x(f) = \langle f, K_x \rangle$. By Cauchy-Schwarz, $|\langle f, K_x \rangle| \leq \|f\|_{\mathcal{H}}^2 \|K_x\|_{\mathcal{H}}^2$. Thus, $L_x(f)$ is bounded. Now we must show that L_x is continuous. Given $\epsilon > 0$ pick $\delta = \frac{\epsilon}{\|K_x\|_{\mathcal{H}}^2}$. Then for $g \in \mathcal{H}$ such that $\|f - g\|_{\mathcal{H}}^2 < \delta$ we have:

$$\begin{aligned} \|L_x(f) - L_x(g)\| &= |L_x(f - g)| \text{ as } L_x \text{ is a linear functional} \\ &= |\langle f - g, K_x \rangle| \\ \text{By Cauchy Schwarz we have:} \\ &\leq \|f - g\|_{\mathcal{H}}^2 \|K_x\|_{\mathcal{H}}^2 \\ &< \frac{\epsilon}{\|K_x\|_{\mathcal{H}}^2} \|K_x\|_{\mathcal{H}}^2 \\ &= \epsilon \end{aligned}$$

Thus, L_x is continuous and H is a reproducing kernel Hilbert space as required. $\square$

The Representer Theorem

Let $\mathcal{H}$ be a reproducing kernel Hilbert space of functions over $\mathbb{R}$ with domain V , a finite dimensional vector space over $\mathbb{R}$ and a set of inputs $X = \{x^{(1)}, \dots, x^{(m)}\} \subset V$ and outputs $Y = \{y^{(1)}, \dots, y^{(m)}\} \subset \mathbb{R}$. Now let $J : \mathcal{H} \rightarrow \mathbb{R}$ given by $J(f) = \sum_{i=1}^m \langle f(x^{(i)}), y^{(i)} \rangle + \lambda \|f\|_{\mathcal{H}}^2$ be a regularized risk function with respect to f . Then, $\forall f \in \mathcal{H} \exists \alpha \in \mathbb{R}^m$ such that $J(f) \leq \sum_{i=1}^m \alpha_i K(x, x^{(i)})$.