# Computational Metaphysics 1

# Denise Erfurt, Martin Lundfall 26. April 2016

### Excercise 1)

**a**)

"The ship is huge and it is blue."

$$Huge(the\_ship) \land Blue(the\_ship)$$
 (1)

b)

"I'm sad if the sun does not shine."

$$\neg Sun\_is\_shining \to Sad(I)$$
 (2)

 $\mathbf{c})$ 

"Either it's raining or it is not."

$$Is\_raining \lor Is\_not\_raining$$
 (3)

d)

"I'm only going if she is going!"

$$I\_am\_going \leftrightarrow She\_is\_going$$
 (4)

**e**) "Everyone loves chocolate or ice cream."  $\forall x Is\_someone(x) \rightarrow Loves\_ice\_cream(x) \lor Loves\_chocolate(x)$ (5)f) "There is somebody who loves ice cream and loves chocolate as well."  $\exists xIs\_someone(x) \land Loves\_ice\_cream(x) \land Loves\_chocolate(x)$ (6) $\mathbf{g}$ "Everyone has got someone to play with."  $\forall x \exists y Is\_someone(x) \land Is\_someone(y) \land Can\_play\_with(x,y)$ (7)h) "Nobody has somebody to play with if they are all mean."  $\forall x Is\_someone(x) \land Is\_mean(x) \rightarrow \neg \exists y \land Is\_somenone(y) \land Can\_play\_with(x,y)$ (8)i) "Cats have the same annoying properties as dogs."  $\forall P \forall cat \forall dog(Is\_annoying(P) \land Is\_cat(cat) \land Is\_dog(dog)) \rightarrow (P(cat) \leftrightarrow P(dog)))$ 

### Excercise 2)

a)

propositional

(9)

### b)

higher-order

#### **c**)

first-order

### d)

higher-order

## Excercise 3)

### **a**)

$$A \wedge B \to C, B \to A, B \vdash C$$
 (10)

1 B

ass.

 $B \to A$ 2

ass.

A3

mp(1,2)

4  $A \wedge B$  conjI(1,3)

 $A \wedge B \to C$  ass. 5

6 C mp(4,5)

## **b**)

$$A \vdash B \to A \tag{11}$$

 $1 \quad A$ 

ass.

 $\begin{array}{c|cc} 2 & B & hyp. \\ \hline 3 & B \to A & impI(1,2) \end{array}$ 

$$A \to (B \to C) \vdash B \to (A \to C) \tag{12}$$

$$\begin{array}{c|cccc}
1 & B & hyp. \\
2 & A & hyp. \\
3 & A \rightarrow (B \rightarrow C) & ass. \\
4 & B \rightarrow C & mp(2,3) & (13) \\
5 & C & mp(1,4) & \\
6 & A \rightarrow C & impI(2,5) & \\
7 & B \rightarrow (A \rightarrow C) & impI(1,6) & 
\end{array}$$

#### d)

$$\neg A \vdash A \to B \tag{14}$$

1 
$$\neg A$$
 ass.

$$\vdash A \lor \neg A \tag{16}$$

$$\begin{array}{c|cccc}
1 & & \neg(A \lor \neg A) & hyp. \\
2 & & A & hyp. \\
3 & & A \lor \neg A & disjI(2) \\
4 & & & notE(1,3) \\
5 & A \lor \neg A & ccontr(1,4)
\end{array} \tag{17}$$

#### Excercise 4)

 $\mathbf{a}$ 

$$\frac{\frac{[A]}{A} \text{ id}}{A \to A} \text{ impI} \tag{18}$$

b)

$$\frac{\frac{[A]}{A} \text{ id}}{\frac{B \to A}{A} \text{ impI}}$$

$$\frac{A \to (B \to A)}{A \to (B \to A)} \text{ impI}$$
(19)

Note that A follows independently of B, so in particular, it follows from B. We can always add arbitrary assumptions, even if our conclusions do not need them.

**c**)

$$\frac{[A]^{1}}{A} \text{ id} \quad [A \to B]^{3} \text{ mp} \quad \frac{[A]^{1}}{A} \text{ id} \quad [A \to (B \to C)]^{2} \text{ mp}$$

$$\frac{B}{B \to C} \text{ mp}$$

$$\frac{C}{A \to C} \text{ impI}_{1}$$

$$\frac{(A \to B) \to (A \to C)}{(A \to B) \to (A \to C)} \text{ impI}_{2}$$

$$\frac{(A \to B) \to (A \to C)}{(A \to B) \to (A \to C)} \text{ impI}_{2}$$
(20)

$$\frac{[B]^{1} \frac{[\neg A]^{2} [\neg A \rightarrow \neg B]^{3}}{\neg B} \text{mp}}{\frac{\frac{\bot}{A} \text{ccontr}_{2}}{B \rightarrow A} \text{impI}_{1}} 
\frac{(\neg A \rightarrow \neg B) \rightarrow (B \rightarrow A)}{(\neg A \rightarrow \neg B)} \text{impI}_{3}$$
(21)