

The diagram model of linear dependent type theory

Martin Lundfall

Stockholm University

martin@dapp.org

Abstract

We present a type theory dealing with non-linear, “ordinary” dependent types (which we will call *cartesian*), and *linear types*, where both constructs may depend on terms of the former. In the interplay between these, we find the new type formers $\prod_{x:A} B$ and $\prod_{x:A} B$, akin to Π and Σ , but where the dependent type B , (and therefore the resulting construct) is a linear type. These can be seen as internalizing universal and existential quantification over linear propositions, respectively. We also consider two modalities, M and L , transforming linear types into cartesian types and vice versa.

The theory is interpreted in a split comprehension category $\pi : \mathcal{T} \rightarrow \mathcal{C}^{\rightarrow}$ [2], accompanied by a split monoidal fibration, $q : \mathcal{L} \rightarrow \mathcal{C}$. We interpret \mathcal{C} as a category of contexts, which for any $\Gamma \in \mathcal{C}$, determines the fibers \mathcal{T}_{Γ} and \mathcal{L}_{Γ} . We interpret \mathcal{T}_{Γ} as category of the cartesian types over Γ , and \mathcal{L}_{Γ} as the monoidal category of linear types in Γ . In this setting, the type formers $\prod_{x:A}$ and $\prod_{x:A}$ are understood as right and left adjoints of the monoidal reindexing functor $\pi_A^* : \mathcal{L}_{\Gamma} \rightarrow \mathcal{L}_{\Gamma.A}$ corresponding to the weakening projection $\pi_A : \Gamma.A \rightarrow \Gamma$ in \mathcal{C} . The operators M and L give rise to a fiberwise adjunction $L \dashv M$ between \mathcal{L} and \mathcal{T} , where we understand the traditional exponential modality as the comonad $! = LM$.

We provide a model of this theory called the *Diagram model*, which extends the groupoid model of dependent type theory [1] to accomodate linear types. Here, cartesian types over a context Γ are interpreted as a family of groupoids indexed over the groupoid Γ , while linear types are interpreted as diagrams over groupoids, $A : \Gamma \rightarrow \mathcal{V}$ in any symmetric monoidal category \mathcal{V} . We show that the *diagrams model* can under certain conditions support a linear analogue of the univalence axiom, and provide some discussion on the higher-dimensional nature of linear dependent types.

2012 ACM Subject Classification Theory of computation \rightarrow Linear logic; Theory of computation \rightarrow Type theory

Keywords and phrases Dependent type theory, linear type theory, diagram model, monoidal categories, groupoid model

Digital Object Identifier 10.4230/LIPIcs.CVIT.2016.23

Related Version http://kurser.math.su.se/pluginfile.php/16103/mod_folder/content/0/2017/2017_47_report.pdf

Acknowledgements I want to thank Peter Lumsdaine for his constant guidance and inspiration.

References

- 1 Martin Hofmann and Thomas Streicher. The groupoid interpretation of type theory. In *Twenty-five years of constructive type theory (Venice, 1995)*, volume 36 of *Oxford Logic Guides*, pages 83–111. Oxford Univ. Press, New York, 1998.



© Martin Lundfall;
licensed under Creative Commons License CC-BY

42nd Conference on Very Important Topics (CVIT 2016).

Editors: John Q. Open and Joan R. Access; Article No. 23; pp. 23:1–23:1

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

23:2 **Models of linear dependent type theory**

- 2 Bart Jacobs. Comprehension categories and the semantics of type dependency. *Theoret. Comput. Sci.*, 107(2):169–207, 1993. URL: [https://doi.org/10.1016/0304-3975\(93\)90169-T](https://doi.org/10.1016/0304-3975(93)90169-T).