

# Math 3770H: Assignment II

Jeremy Favro (0805980)  
Trent University, Peterborough, ON, Canada

October 5, 2025

**Problem 1.** Write the function

$$f(z) = z + \frac{1}{z} \quad (z \neq 0)$$

in the form  $f(z) = u(r, \theta) + iv(r, \theta)$

**Solution 1.** Writing  $f$  first in polar form to switch to a dependence on  $r = |z|, \theta = \arctan(\operatorname{Im} z / \operatorname{Re} z)$ ,

$$z = re^{i\theta} = r \cos \theta + ir \sin \theta.$$

This is actually really nice here as it means we can easily write the inverse without using a fraction,

$$z^{-1} = r^{-1}e^{-i\theta} = r^{-1} \cos(-\theta) + ir^{-1} \sin(-\theta).$$


Which can be reworked further using the even/odd nature of  $\cos$  and  $\sin$  respectively giving:

$$\begin{aligned} f(z) &= r \cos \theta + ir \sin \theta + r^{-1} \cos \theta - ir^{-1} \sin \theta \\ &= r \cos \theta + r^{-1} \cos \theta + ir \sin \theta - ir^{-1} \sin \theta \\ &= (r + r^{-1}) \cos \theta + (r - r^{-1}) \sin \theta \end{aligned}$$

**Problem 2.** Sketch the region onto which the sector  $r \leq 1$ ,  $0 \leq \theta \leq \pi/4$  is mapped by the transformation


(a)  $w = z^2$ ;      (b)  $w = z^3$ ;      (c)  $w = z^4$ .

**Solution 2.** Generally the transform  $w = z^n$  will give  $w = r^n e^{in\theta}$ , just by exponentiation laws. Our original region looks like:




Because our region only extends out to  $r = 1$  the radius won't change.


(a)



(b)



(c)



**Problem 3.** Use definition (2), Sec.15, of the limit to prove that

$$(a) \lim_{z \rightarrow z_0} \operatorname{Re} z = \operatorname{Re} z_0; \quad (b) \lim_{z \rightarrow z_0} \bar{z} = \overline{z_0}; \quad (c) \lim_{z \rightarrow 0} \frac{\bar{z}^2}{z} = 0.$$

**Solution 3.**

(a) Let  $\epsilon > 0$ . Suppose  $|z - z_0| < \delta$ . If we take  $z = x + iy$  and  $z_0 = x_0 + iy_0$  then we have

$$|\operatorname{Re} z - \operatorname{Re} z_0| = |x - x_0| \leq |z - z_0|.$$

We can make that last statement because

$$|x - x_0| \leq |x - x_0 + iy - iy_0| = |z - z_0|.$$

This means that

$$|\operatorname{Re} z - \operatorname{Re} z_0| = |z - z_0| \leq \delta$$

by the  $\epsilon - \delta$  definition we are using. Having this means that provided  $|z - z_0| \leq \delta = \epsilon$ , then  $|\operatorname{Re} z - \operatorname{Re} z_0| < \epsilon$  by transitivity.

(b) Let  $\epsilon > 0$ . Suppose  $|z - z_0| < \delta$ . We have

$$|\bar{z} - \overline{z_0}| = |\overline{z - z_0}| = |z - z_0| < \delta$$

all by properties of the complex conjugate. So provided  $|z - z_0| < \delta = \epsilon$ , then  $|\bar{z} - \overline{z_0}| < \epsilon$ .

(c) Let  $\epsilon > 0$ . Suppose  $|z - z_0| < \delta$ . We have

$$\left| \frac{\bar{z}^2}{z} - \frac{\overline{z_0}^2}{z_0} \right| = \left| \frac{\bar{z}\bar{z}}{z} - \frac{\overline{z_0}z_0}{z_0} \right|$$

$$\lim_{z \rightarrow 0} = 0.$$

**Problem 4.** With the aid of the theorem in Sec. 17, show that when

$$T(z) = \frac{az + b}{cz + d} \quad (ad - bc) \neq 0$$

(a)  $\lim_{z \rightarrow \infty} T(z) = \infty$  if  $c = 0$ ;

(b)  $\lim_{z \rightarrow \infty} T(z) = \frac{a}{c}$  and  $\lim_{z \rightarrow -d/c} T(z) = \infty$  if  $c \neq 0$

**Solution 4.**

**Problem 5.** Use the method in Example 2, Sec. 19, to show that  $f(z)$  does not exist at any point  $z$  when

(a)  $f(z) = \operatorname{Re} z$ ; (b)  $f(z) = \operatorname{Im} z$ .

**Solution 5.**

**Problem 6.** Use the theorem in Sec. 24 to show that each of these functions is differentiable in the indicated domain of definition, and also to find  $f'(z)$ :

(a)  $f(z) = 1/z^4$  ( $z \neq 0$ );

(b)  $f(z) = e^{-\theta} \cos(\ln r) + ie^{-\theta} \sin(\ln r)$  ( $r > 0, 0 < \theta < 2\pi$ ).

**Solution 6.**

**Problem 7.** With the aid of the theorem in Sec. 21, show that each of these functions is nowhere analytic:

(a)  $f(z) = xy + iy$ ; (b)  $f(z) = 2xy + i(x^2 - y^2)$ ; (c)  $f(z) = e^y e^{ix}$ .

**Solution 7.**

**Problem 8.A.** Let the function  $f(z) = u(x, y) + iv(x, y)$  be analytic in a domain  $D$ , and consider the families of level curves  $u(x, y) = c_1$  and  $v(x, y) = c_2$ , where  $c_1$  and  $c_2$  are arbitrary real constants. Prove that these families are orthogonal. More precisely, show that if  $z_0 = (x_0, y_0)$  is a point in  $D$  which is common to two particular curves  $u(x, y) = c_1$  and  $v(x, y) = c_2$  and if  $f'(z_0) \neq 0$ , then the lines tangent to those curves at  $(x_0, y_0)$  are perpendicular.

**Problem 8.B.** Sketch the families of level curves of the component functions  $u$  and  $v$  when  $f(z) = 1/z$ , and note the orthogonality described in Exercise 2.

**Solution 8.A.**

**Solution 8.B.**