

§1 IMO

Problem 1.1 (IMO 2019 P1)

Find all $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that,

$$f(2x) + 2f(y) = f(f(x + y))$$

Proof. Let $P(x, y)$ the condition generated by x, y . Then,

$$\begin{cases} P(x, 0) : f(2x) + 2f(0) = f(f(x)) \\ P(0, x) : f(0) + 2f(x) = f(f(x)) \end{cases}$$

Combining these two we obtain that,

$$f(2x) = 2f(x) - f(0)$$

Now let us consider the following,

$$P(x, 1) : f(2x) + 2f(1) = f(f(x + 1))$$

However from the previous observations we can say that,

$$f(f(x + 1)) = 2f(x + 1) + f(0)$$

Thus,

$$\begin{aligned} 2f(x) + 2f(1) - f(0) &= 2f(x + 1) + f(0) \\ f(x + 1) &= f(x) + f(1) - f(0) \end{aligned}$$

Thus, we see that $f(x) = f(0) + x \cdot C$ where C is some constant (specifically $f(1) - f(0)$) for all $x \in \mathbb{Z}$. Now let us consider,

$$P(x, -x) : f(2x) + 2f(-x) = f(f(0))$$

We can rewrite this as,

$$\begin{aligned} f(0) + 2xC + 2f(0) - 2xC &= f(0) + f(0) \cdot C \\ \implies 2f(0) &= f(0) \cdot C \end{aligned}$$

This implies that either $f(0) = 0$ or $C = 2$. Let us consider both cases,

- If $f(0) = 0$, then, $f(x) = cx$ and it is simple to verify that then $f(x)$ must equal $2x$ or 0 .
- If $C = 2$, then,

$$\begin{aligned} f(0) + 4x + 2(f(0) + 2y) &= f(f(0) + 2(x + y)) = f(0) + 2f(0) + 4(x + y) \\ 4x + 2f(0) + 4y &= 2f(0) + 4x + 4y \end{aligned}$$

Thus, we see the only solutions to this functional equation are $f(x) = 2x + c$ and $f(x) = 0$. \square

Problem 1.2 (IMO 2022 P2)

Find all $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for each $x \in \mathbb{R}^+$ there exists exactly one $y \in \mathbb{R}^+$ such that,

$$xf(y) + yf(x) \leq 2$$

Proof. Obviously $f(x) = \frac{1}{x}$ works (from AM-GM).

Let us say that $x \sim y$ if and only if the condition from the problem statement is true. Then obviously this is commutative, meaning that if $x \sim y$ then $y \sim x$.

Lemma 1.1 If $f(x) \leq \frac{1}{x}$, then, $x \sim x$.

This is simple to see due to,

$$2xf(x) \leq 2x \cdot \frac{1}{x} = 2$$

However, notice that if $x \sim y$, then,

$$xf(y) + yf(x) \leq 2$$

and in it must be that $f(x) > \frac{1}{x}$ and $f(y) > \frac{1}{y}$ which would imply that by AM-GM,

$$xf(y) + yf(x) = \frac{x}{y} + \frac{y}{x} \geq 2$$

Thus we obtain that for all $x \in \mathbb{R}^+$ it must be that $x \sim x$. Thus that,

$$f(x) \leq \frac{1}{x}$$

for all $x \in \mathbb{R}^+$.

Notice that if $f(a) < \frac{1}{a}$, then, if,

$$f(a) = \frac{1}{a} - \epsilon$$

then,

$$xf(a) + af(x) = x \left(\frac{1}{a} - \epsilon \right) + af(x) \leq x \left(\frac{1}{a} - \epsilon \right) + \frac{a}{x} \stackrel{?}{\leq} 2$$

However the last inequality for $\epsilon > 0$ has multiple solutions for x . Because,

$$x^2 \left(\frac{1}{a} - \epsilon \right) + a - 2x \leq 0$$

The only way for this inequality to have a single solution for $x \in \mathbb{R}^+$ is if the equality has only one solution, thus the discriminant is zero,

$$\Delta = 4 - 4 \cdot a \cdot \left(\frac{1}{a} - \epsilon \right)$$

which is obviously never 0 for $\epsilon > 0$. Thus because for each x there is only one such y that $x \sim y$ it must mean that,

$$f(x) = \frac{1}{x}$$


for all $x \in \mathbb{R}^+$.

□

Problem 1.3 (IMO 2014 P4)

Let P and Q be on segment BC of an acute triangle $\triangle ABC$ such that $\angle PAB = \angle BCA$ and $\angle CAQ = \angle ABC$. Let M and N be the points on AP and AQ , respectively, such that P is the midpoint of AM and Q is the midpoint of AN . Prove that the intersection of BM and CN is on the circumference of triangle $\triangle ABC$.

Proof. Let $\triangle ABC$ be the reference triangle in barycentric coordinates.



Then we can calculate the point P , because $\triangle ABP \sim \triangle CBA$ and $\triangle ACQ \sim \triangle BCA$. Thus,

$$\begin{aligned} \frac{BA}{BC} &= \frac{BP}{BA} \\ \Rightarrow BP &= \frac{BA^2}{BC} = \frac{c^2}{a} \end{aligned}$$

consequently, $P = \left(0 : a - \frac{c^2}{a} : \frac{c^2}{a}\right)$, analogously we get, $Q = \left(0 : \frac{b^2}{a} : a - \frac{b^2}{a}\right)$. Now we can calculate points M and N ,

$$M = (-a^2 : 2a^2 - 2c^2 : 2c^2)$$

$$N = (-a^2 : 2b^2 : 2a^2 - 2b^2)$$

Now we simply need to intersect BM and CN which is not hard to do considering that both are cevians in the triangle. It is not hard to see that the following point satisfies these conditions,

$$X = (-a^2 : 2b^2 : 2c^2)$$

Now we just need to check that this point lies on the circumcircle of $\triangle ABC$ which is given by,

$$a^2yz + b^2xz + c^2xy = 0$$

Thus,

$$a^2yz + b^2xz + c^2xy = 4a^2b^2c^2 - 2a^2b^2c^2 - 2a^2b^2c^2 = 0$$

Consequently, it must be that $X \in (\triangle ABC)$. □

Problem 1.4 (IMO 2022 Shortlist, G2)

In the acute-angled triangle $\triangle ABC$, the point F is the foot of the altitude from A , and P is a point on the segment AF . The lines through P parallel to AC and AB meet BC at D and E , respectively. Points $X \neq A$ and $Y \neq A$ lie on the circles ABD and ACE , respectively, such that $DA = DX$ and $EA = EY$. Prove that B, C, X and Y are concyclic.

Proof. There are a lot of observations that can be done in this problem, however this is the solution I have.



Lemma 1.2 The second intersection of (ABD) and (AEC) lies on AH .

This is true due to basic angle chase, assume that F lies on AH and (ABD) , let us prove that then $AECF$ is cyclic. By Power of the Point,

$$HA \cdot HF = HD \cdot HD = HE \cdot HC$$

the last is obviously true because $PE \parallel AB$ and $PD \parallel AC$. Now let us intersect BX with AF at a point W , then the condition that $BXCY$ is cyclic by the radical center theorem is equivalent to showing that W, C and Y are colinear.

Lemma 1.3 BC is the angle bisector of $\angle ABW$

This is because,


$$\angle CBW = \angle XAD = \angle AXD = \angle ABC$$

because F is the midpoint of the arc between A and X per definition. Thus, W is the reflection of A over H . The exact same argument for C shows that BC is the angle bisector of $\angle YCA$ and thus, CY must intersect AF at W . \square

Problem 1.5 (IMO 2012 P1)

Given triangle ABC the point J is the center of the excircle opposite the vertex A . This excircle is tangent to side BC at M , and to lines AB and AC at K and L , respectively. Lines LM and BJ meet at F , and lines KM and CJ meet at G . Let S be the point of intersection of lines AF and BC , and let T be the point of intersection of lines AG and BC . Prove that M is the midpoint of ST .

Proof. Pascal's theorem hints us towards checking whether $FAGKJL$ is cyclic.



Notice, that because $KM \perp FJ$ and $JG \perp ML$, it must be that M is the orthocenter of FGJ . Thus, $FG \parallel BC$, consequently,

$$\angle GFJ = \angle BCJ = \angle MKJ$$

Thus, $FKJG$ is cyclic, analogously we conclude that $FKJLG$ is cyclic and by Pascal's theorem it must be that A lies on this circle as well, thus $FAGLJK$ is cyclic.

Now notice,

$$\angle FGA = \angle ALF = \angle CML = \angle GFL$$

the last step comes from $FG \parallel BC$. Thus, $AT \parallel FM$, the same for $AG \parallel FM$. Consequently, $AGMF$ is a parallelogram. Because of that it must be that FG is the midline of $\triangle AST$ and M is the midpoint of the side ST . \square

Problem 1.6 (IMO 2022 Shortlist C1)

A ± 1 -sequence is a sequence of 2022 numbers a_1, \dots, a_{2022} , each equal to either $+1$ or -1 . Determine the largest C such that, for any ± 1 -sequence, there exists an integer k and indices $1 \leq t_1 \leq \dots \leq t_k \leq 2022$ so that $t_{i+1} - t_i \leq 2$ for all i , and

$$\left| \sum_{i=1}^k a_{t_i} \right| \geq C$$

Proof. WLOG, let there be more 1 than -1 in the ± 1 -sequence. Then, let us consider the following algorithm,

Algorithm 1.4 If the next element is -1 , then skip to the next one and include it. If the next element is 1 , then include it. Continue this process until you have went through the entire sequence.

This algorithm will guarantee that we skip at least $\lceil \frac{X}{2} \rceil$, where X is the number of -1 in the sequence. Thus,

$$\left| \sum_{i=1}^k a_{t_i} \right| \geq (2022 - X) - \left\lfloor \frac{X}{2} \right\rfloor \geq (2022 - 1011) - \left\lfloor \frac{1011}{2} \right\rfloor = 506$$

Consequently, it must be that $C \geq 506$. Now consider the following sequence,

$$1, 1, -1, -1, 1, 1, \dots, -1, -1, 1, -1$$

It alternates between $1, 1$ and $-1, -1$ until the very end, where it is 1 and -1 .

Notice that in each consecutive pair of identical numbers at least one of the numbers is present in our sequence. Thus, no matter our choice of indices, the total sum is bounded by,

$$\left| \sum_{i=1}^k a_{t_i} \right| \leq 2 \cdot \frac{1010}{2} - \frac{1010}{2} + 1 = 2 \cdot 505 - 505 + 1 = 506$$

Thus, it must be that $C = 506$. □