

# Master Theorem: Practice Problems and Solutions

## Practice Problems

For each of the following recurrences, give an expression for the runtime  $T(n)$  if the recurrence can be solved with the Master Theorem. Otherwise, indicate that the Master Theorem does not apply.

1.  $T(n) = 3T(n/2) + n^2$

2.  $T(n) = 4T(n/2) + n^2$

3.  $T(n) = T(n/2) + 2^n$

4.  $T(n) = 2^n T(n/2) + n^n$

5.  $T(n) = 16T(n/4) + n$

6.  $T(n) = 2T(n/2) + n \log n$

---

<sup>1</sup>most of the time,  $k = 0$

$$7. T(n) = 2T(n/2) + n/\log n$$

$$8. T(n) = 2T(n/4) + n^{0.51}$$

$$9. T(n) = 0.5T(n/2) + 1/n$$

$$10. T(n) = 16T(n/4) + n!$$

$$11. T(n) = \sqrt{2}T(n/2) + \log n$$

$$12. T(n) = 3T(n/2) + n$$

$$13. T(n) = 3T(n/3) + \sqrt{n}$$

$$14. T(n) = 4T(n/2) + cn$$

$$15. T(n) = 3T(n/4) + n \log n$$

$$16. T(n) = 3T(n/3) + n/2$$

$$17. T(n) = 6T(n/3) + n^2 \log n$$

$$18. T(n) = 4T(n/2) + n/\log n$$

$$19. T(n) = 64T(n/8) - n^2 \log n$$

$$20. T(n) = 7T(n/3) + n^2$$

$$21. T(n) = 4T(n/2) + \log n$$

$$22. T(n) = T(n/2) + n(2 - \cos n)$$

## Solutions

1.  $T(n) = 3T(n/2) + n^2 \implies T(n) = \Theta(n^2)$  (Case 3)
2.  $T(n) = 4T(n/2) + n^2 \implies T(n) = \Theta(n^2 \log n)$  (Case 2)
3.  $T(n) = T(n/2) + 2^n \implies \Theta(2^n)$  (Case 3)
4.  $T(n) = 2^n T(n/2) + n^n \implies$  Does not apply ( $a$  is not constant)
5.  $T(n) = 16T(n/4) + n \implies T(n) = \Theta(n^2)$  (Case 1)
6.  $T(n) = 2T(n/2) + n \log n \implies$  Does not apply  
(no polynomial separation between  $f(n)$  and the number of leaves.)
7.  $T(n) = 2T(n/2) + n/\log n \implies$  Does not apply (non-polynomial difference between  $f(n)$  and  $n^{\log_b a}$ )
8.  $T(n) = 2T(n/4) + n^{0.51} \implies T(n) = \Theta(n^{0.51})$  (Case 3)
9.  $T(n) = 0.5T(n/2) + 1/n \implies$  Does not apply ( $a < 1$ )
10.  $T(n) = 16T(n/4) + n! \implies T(n) = \Theta(n!)$  (Case 3)
11.  $T(n) = \sqrt{2}T(n/2) + \log n \implies T(n) = \Theta(\sqrt{n})$  (Case 1)
12.  $T(n) = 3T(n/2) + n \implies T(n) = \Theta(n^{\lg 3})$  (Case 1)
13.  $T(n) = 3T(n/3) + \sqrt{n} \implies T(n) = \Theta(n)$  (Case 1)
14.  $T(n) = 4T(n/2) + cn \implies T(n) = \Theta(n^2)$  (Case 1)
15.  $T(n) = 3T(n/4) + n \log n \implies T(n) = \Theta(n \log n)$  (Case 3)
16.  $T(n) = 3T(n/3) + n/2 \implies T(n) = \Theta(n \log n)$  (Case 2)
17.  $T(n) = 6T(n/3) + n^2 \log n \implies T(n) = \Theta(n^2 \log n)$  (Case 3)
18.  $T(n) = 4T(n/2) + n/\log n \implies T(n) = \Theta(n^2)$  (Case 1)
19.  $T(n) = 64T(n/8) - n^2 \log n \implies$  Does not apply ( $f(n)$  is not positive)
20.  $T(n) = 7T(n/3) + n^2 \implies T(n) = \Theta(n^2)$  (Case 3)
21.  $T(n) = 4T(n/2) + \log n \implies T(n) = \Theta(n^2)$  (Case 1)
22.  $T(n) = T(n/2) + n(2 - \cos n) \implies$  Does not apply. We are in Case 3, but the regularity condition is violated. (Consider  $n = 2\pi k$ , where  $k$  is odd and arbitrarily large. For any such choice of  $n$ , you can show that  $c \geq 3/2$ , thereby violating the regularity condition.)