

General Kernel Spectral Methods for Equilibrium Measures

MMSC Dissertation



Peter Julius Waldert
Mathematical Institute
University of Oxford

- ▶ Simulate Many-Particle-Systems and find their Equilibrium Distribution
- ▶ Interactions through (power-law) Attraction-Repulsion Potentials

$$K(r) = \frac{r^\alpha}{\alpha} - \frac{r^\beta}{\beta} \quad \alpha, \beta \in \mathbb{R}.$$

- ▶ Each particle $i = 1, \dots, N$ at position \mathbf{x}_i and time t follows

$$\frac{d^2 \mathbf{x}_i}{dt^2} = f\left(\left\|\frac{d\mathbf{x}_i}{dt}\right\|_2\right) \frac{d\mathbf{x}_i}{dt} - \frac{1}{N} \sum_{j=1, i \neq j}^N \nabla K(\|\mathbf{x}_i - \mathbf{x}_j\|_2),$$

(Gutleb, Carrillo and Olver [2020](#); Gutleb, Carrillo and Olver [2021](#)).

- [1] Timon S. Gutleb, José A. Carrillo and Sheehan Olver. ‘Computation of Power Law Equilibrium Measures on Balls of Arbitrary Dimension’. In: *arXiv* (Sept. 2021). DOI: [10.1007/s00365-022-09606-0](https://doi.org/10.1007/s00365-022-09606-0). eprint: [2109.00843](https://arxiv.org/abs/2109.00843).
- [2] Timon S. Gutleb, José A. Carrillo and Sheehan Olver. ‘Computing Equilibrium Measures with Power Law Kernels’. In: *arXiv* (Oct. 2020). DOI: [10.1090/mcom/3740](https://doi.org/10.1090/mcom/3740). eprint: [2011.00045](https://arxiv.org/abs/2011.00045).