

General Kernel Spectral Methods for Equilibrium Measures

Seminar Talk at the University of Graz




Peter Julius Waldert

Mathematical Institute
University of Oxford


supervised by
Dr. Timon Gutleb and Prof. José Carrillo

9th of November, 2023

- Find the equilibrium distribution $\rho(\mathbf{x})$ of a many-particle system.



$N_p = 8$ particles interacting with one another through the potential $K(r)$.



Plot of attractive-repulsive potential functions $K(r) = \frac{r^\alpha}{\alpha} - \frac{r^\beta}{\beta}$ for different α, β .

- Model interactions through (power law) attraction-repulsion potentials¹

for example, $K(r) = \frac{r^\alpha}{\alpha} - \frac{r^\beta}{\beta}$ with parameters $\alpha, \beta \in \mathbb{R} \setminus \{0\}$.

- Each particle $i = 1, \dots, N$ at position $\mathbf{x}_i \in \mathbb{R}^d$ and time $t \in \mathbb{R}^+$ follows

$$\frac{d^2 \mathbf{x}_i}{dt^2} = f \left(\left\| \frac{d\mathbf{x}_i}{dt} \right\|_2 \right) \frac{d\mathbf{x}_i}{dt} - \frac{1}{N} \sum_{j=1, i \neq j}^N \nabla K(\|\mathbf{x}_i - \mathbf{x}_j\|_2),$$

for reference see, for example, [1, 2]. For now, we only consider the case without an external potential $V(\mathbf{x})$.

¹If the repulsive term is stronger (so $\beta > \alpha$), there is no equilibrium distribution as particles simply continue repelling each other out to infinity.




Figure: Comparison of different interaction potentials.




Figure: $N_p = 120$ particles in $d = 2$ dimensions.




Figure: Self-propelled particles in a reflective box $[-1, 1]^3$.

Every particle i at position $\mathbf{x}_i \in \mathbb{R}^d$ with velocity $\mathbf{v}_i \in \mathbb{R}^d$ is updated using

$$\begin{aligned}\mathbf{x}_i(t + \tau) &= \mathbf{x}_i(t) + \tau \cdot \mathbf{v}_i(t + \tau/2), & \text{for } t = 0, \tau, \dots, \\ \mathbf{v}_i(t + \tau/2) &= \mathbf{v}_i(t - \tau/2) + \tau \cdot \mathbf{f}[\mathbf{x}_i(t), t], & \text{for } t = \tau, 2\tau, \dots, \\ \mathbf{v}_i(\tau/2) &= \mathbf{v}_i(0) + \frac{\tau}{2} \cdot \mathbf{f}[\mathbf{x}_i(0), 0], & \text{for } t = 0,\end{aligned}$$

where $\mathbf{f}_i[\mathbf{x}_i(t), t] \in \mathbb{R}^d$ denotes the acceleration (sum of contributions of all forces divided by particle mass m_i) at time t .

In fewer words... a particle at position $\mathbf{x} \in \mathbb{R}^d$ with velocity $\mathbf{v} \in \mathbb{R}^d$





Figure: Visualisation of the Leapfrog integration method, position and velocity are updated at times shifted by $\tau/2$, half the timestep.



The positional distribution approached by $N_p = 250$ particles.

The total potential energy of an N_p -particle system is then given by

$$E = \sum_{i=1}^{N_p} \sum_{j=1, j \neq i}^{N_p} K(\|\mathbf{x}_i - \mathbf{x}_j\|_2) ,$$

which, in the continuous limit as $N_p \rightarrow \infty$, becomes

$$E = \frac{1}{2} \iint K(\|\mathbf{x} - \mathbf{y}\|_2) \, \mathrm{d}\rho(\mathbf{x}) \, \mathrm{d}\rho(\mathbf{y}) ,$$

where $\mathrm{d}\rho = \rho(\mathbf{x})\mathrm{d}\mathbf{x}$ is a measure (the equilibrium distribution) chosen such that

$$M = \int \mathrm{d}\rho = \int_{\text{supp}(\rho)} \rho(\mathbf{x}) \, \mathrm{d}\mathbf{x} = 1 .$$

Definition (Equilibrium Measure)

For a given pairwise interaction potential $K : \mathbb{R} \rightarrow \mathbb{R}$, the equilibrium measure $\hat{\rho} : D \rightarrow \mathbb{R}$ with $D \subseteq \mathbb{R}^d$ is a measure chosen such that

$$U_K[\hat{\rho}] := \frac{1}{2} \iint K(\|\hat{\mathbf{x}} - \hat{\mathbf{y}}\|_2) \, \mathrm{d}\hat{\rho}(\hat{\mathbf{x}}) \, \mathrm{d}\hat{\rho}(\hat{\mathbf{y}}),$$

is minimised, where $\mathrm{d}\hat{\rho} = \hat{\rho}(\hat{\mathbf{x}})\mathrm{d}\hat{\mathbf{x}}$ [2].

Definition (Equilibrium Measure)

For a given pairwise interaction potential $K : \mathbb{R} \rightarrow \mathbb{R}$, the equilibrium measure $\hat{\rho} : D \rightarrow \mathbb{R}$ with $D \subseteq \mathbb{R}^d$ is a measure chosen such that

$$U_K[\hat{\rho}] := \frac{1}{2} \iint K(\|\hat{\mathbf{x}} - \hat{\mathbf{y}}\|_2) d\hat{\rho}(\hat{\mathbf{x}}) d\hat{\rho}(\hat{\mathbf{y}}),$$

is minimised, where $d\hat{\rho} = \hat{\rho}(\hat{\mathbf{x}})d\hat{\mathbf{x}}$ [2].

For example: in a two-particle system, $\hat{\rho}(\hat{\mathbf{x}}) = \delta(\hat{\mathbf{x}} - \hat{\mathbf{p}}_1) + \delta(\hat{\mathbf{x}} - \hat{\mathbf{p}}_2)$ to see that $U_K[\hat{\rho}] = K(0) + K(\|\hat{\mathbf{p}}_1 - \hat{\mathbf{p}}_2\|_2)$, and hence, the total energy becomes

$$E = \frac{m}{2} \left(\|\hat{\mathbf{v}}_1\|_2^2 + \|\hat{\mathbf{v}}_2\|_2^2 \right) + K(0) + K(\|\hat{\mathbf{p}}_1 - \hat{\mathbf{p}}_2\|_2).$$

- ▶ In order to find $\rho(\mathbf{x})$, we consider an ansatz of the following form:

$$\rho(\mathbf{x}) = \sum_{n=0}^{\infty} c_n \varphi_n(\mathbf{x}), \quad c_n \in \mathbb{R},$$

with which we construct a spectral method for the numerical solution of the above integral equation.

- ▶ Minimization routine of E over coefficients in ρ , as a subroutine of outer minimisation over the bounds of the box (simpler case: use $[-r, r]$, $r \in \mathbb{R}^+$).
- ▶ By construction, we find that we do not need an iterative approach for the inner optimisation routine.
- ▶ The outer minimisation can be performed using known methods from continuous optimisation.

Jacobi polynomials $P_n^{(a,b)}(x)$ are orthogonal on $[-1, 1]$ w.r.t. the weight function

$$w^{(a,b)}(x) = (1-x)^a(1+x)^b,$$

so they satisfy

$$\int_{-1}^1 (1-x)^a(1+x)^b P_n^{(a,b)} P_m^{(a,b)} dx = \frac{2^{a+b+1} \Gamma(a+n+1) \Gamma(b+n+1)}{n!(a+b+2n+1) \Gamma(a+b+n+1)} \delta_{n,m},$$

with $a, b > -1$, which uniquely determines $P_n^{(a,b)}(x)$. The special case of $a = b$ corresponds to the ultraspherical or Gegenbauer polynomials, while the case $a = b = 0$ corresponds to the Legendre polynomials [3].

- This basis yields a **sparse**, and in particular, **banded** operator.

Definition (Particle Density Distribution Problem)

Given an interaction kernel $K : \mathbb{R}^+ \rightarrow \mathbb{R}$, the density distribution problem is to find the equilibrium measure $\hat{\rho} : B_R(\mathbf{0}) \rightarrow \mathbb{R}$ of mass $M = 1$ on a d -dimensional ball of radius $R \in \mathbb{R}^+$ that minimises the total potential $U_K[\hat{\rho}]$.

Definition (Particle Density Distribution Problem)

Given an interaction kernel $K : \mathbb{R}^+ \rightarrow \mathbb{R}$, the density distribution problem is to find the equilibrium measure $\hat{\rho} : B_R(\mathbf{0}) \rightarrow \mathbb{R}$ of mass $M = 1$ on a d -dimensional ball of radius $R \in \mathbb{R}^+$ that minimises the total potential $U_K[\hat{\rho}]$.

As mentioned above, we will consider a **spectral method** to solve this problem. More precisely, using the following ansatz:

$$\rho(\mathbf{x}) := \left(1 - \|\mathbf{x}\|_2^2\right)^{m - \frac{\alpha+d}{2}} \sum_{k=0}^{N-1} \rho_k P_k^{\left(m - \frac{\alpha+d}{2}, \frac{d-2}{2}\right)} \left(2 \|\mathbf{x}\|_2^2 - 1\right), \quad (1)$$

with $P_k^{(a,b)}$ the k th Jacobi polynomial and ρ_k its corresponding coefficient.

Definition (Power Law Operator \mathcal{Q}^β)

The power law operator $\mathcal{Q}^\beta : \mathcal{L} \rightarrow \mathcal{L}$ is given by

$$\mathcal{Q}^\beta[\rho](\mathbf{x}) := \int \|\mathbf{x} - \mathbf{y}\|_2^\beta \, d\rho(\mathbf{y}) = \int_{\text{supp}(\rho)} \|\mathbf{x} - \mathbf{y}\|_2^\beta \rho(\mathbf{y}) \, d\mathbf{y}.$$

Applied to the ansatz given in (1), we can evaluate the appearing integrals **explicitly**. For the attractive-repulsive interaction kernel $K_{\alpha,\beta}(r) = \frac{r^\alpha}{\alpha} - \frac{r^\beta}{\beta}$, the matrix representation of the operator becomes

$$Q_{\alpha,\beta} := \frac{R^{\alpha+d}}{\alpha} Q^\alpha - \frac{R^{\beta+d}}{\beta} Q^\beta. \quad (2)$$

Theorem (Power Law Potential of the n th Jacobi Polynomial)

On the d -dimensional unit ball $B_1(\mathbf{0})$ the power law potential, with power $\alpha \in (-d, 2 + 2m - d)$, $m \in \mathbb{N}_0$ and $\beta > -d$, of the n th weighted radial Jacobi polynomial $(1 - \|\mathbf{y}\|_2^2)^{m - \frac{\alpha+d}{2}} P_n^{(m - \frac{\alpha+d}{2}, \frac{d-2}{2})} (2\|\mathbf{y}\|_2^2 - 1)$ reduces to a Gaussian hypergeometric function as follows:

$$\begin{aligned} I_{m,n}^{\alpha,\beta}(\mathbf{x}) &= \int_{B_1(\mathbf{0})} \|\mathbf{x} - \mathbf{y}\|_2^\beta (1 - \|\mathbf{y}\|_2^2)^{m - \frac{\alpha+d}{2}} P_n^{(m - \frac{\alpha+d}{2}, \frac{d-2}{2})} (2\|\mathbf{y}\|_2^2 - 1) d\mathbf{y} \\ &= \frac{\pi^{d/2} \Gamma(1 + \frac{\beta}{2}) \Gamma(\frac{\beta+d}{2}) \Gamma(m+n - \frac{\alpha+d}{2} + 1)}{\Gamma(\frac{d}{2}) \Gamma(n+1) \Gamma(\frac{\beta}{2} - n + 1) \Gamma(\frac{\beta-\alpha}{2} + m+n+1)} {}_2F_1 \left(n - \frac{\beta}{2}, -m - n + \frac{\alpha-\beta}{2}; \frac{d}{2}; \|\mathbf{x}\|_2^2 \right). \end{aligned}$$

Lemma (Mass of the Solution)

For a given solution $\rho : B_1(\mathbf{0}) \rightarrow \mathbb{R}$, its mass $M \in \mathbb{R}$ can be evaluated explicitly, provided the appropriate ansatz, an expansion of weighted radial Jacobi polynomials with coefficients ρ_k . The mass is given by

$$M[\rho[\boldsymbol{\rho}]] = \int_{\text{supp}(\rho)} \rho(y) \, dy = \frac{\pi^{d/2} \Gamma(a+1)}{\Gamma(a+d/2+1)} \rho_0,$$

so solely depending on the first coefficient ρ_0 .




Figure: The attractive and repulsive operators (matrices), the (absolute) matrix values are shown in a \log_{10} colour scale. Due to the choice of basis, the attractive operator is exactly banded.




Figure: The analytical solution $\rho(x)$ compared to the (spectral method) solutions $\rho_N(x)$ of different order.




Figure: Simulation output with $N_p = 150$ particles using the same attractive-repulsive kernel $K_{\alpha, \beta}(r)$.

We express the general kernel as a polynomial, through reprojection from the Jacobi polynomials,

$$K_G(r) = \sum_{l=0}^{G-1} g_l r^l \approx K(r), \quad \mathbf{g} := (g_0, \dots, g_{G-1})^T \in \mathbb{R}^G. \quad (3)$$

The operator can then be expressed as

$$\mathcal{Q}_G[\hat{\rho}](\hat{\mathbf{x}}) = \int_{B_R(\mathbf{0})} \sum_{l=0}^{G-1} g_l \|\hat{\mathbf{x}} - \hat{\mathbf{y}}\|_2^l \hat{\rho}(\hat{\mathbf{y}}) \, \mathrm{d}\hat{\mathbf{y}} = \sum_{l=0}^{G-1} g_l R^{l+d} \mathcal{Q}^l[\hat{\rho}](\mathbf{x}).$$




Figure: Comparison of simulation output with the $G = 8$ general kernel solver's equilibrium measure $\rho_{12}(r)$ at R given by the simulator.




Figure: Growth of the 2-norm condition number $\kappa_2(Q)$ of the attractive-repulsive operators $Q^{(\alpha)}$, $Q^{(\beta)}$ and $Q_{\alpha, \beta}$ for growing system size N .




Figure: Absolute value of the solution coefficients ρ_k with and without Tikhonov regularisation after the solution of a 200×200 linear system.

Contributions:

- ▶ Original implementation of simulator (+GUI, in C++) and solver (in Julia).
- ▶ General kernel spectral method + implementation.
- ▶ Lemma for an initial guess of R .

Challenges:

- ▶ Proving Theorem 4.3, the power law integral of the Jacobi polynomials.
- ▶ Synchronisation of simulation and solver (implementations).
- ▶ Correct parameter choices for the general kernel spectral method.

Future Work:

- ▶ Describing phase-space distributions in a self-propulsion setup.
- ▶ Modelling (constrained) boundaries and boundary conditions.

Thank you.

- [1] Timon S. Gutleb, José Antonio Carrillo and Sheehan Olver. ‘Computing equilibrium measures with power law kernels’. In: *Math. Comput.* 91.337 (Sept. 2022), pp. 2247–2281. ISSN: 0025-5718. DOI: [10.1090/mcom/3740](https://doi.org/10.1090/mcom/3740).
- [2] Timon S. Gutleb, José Antonio Carrillo and Sheehan Olver. ‘Computation of Power Law Equilibrium Measures on Balls of Arbitrary Dimension’. In: *Constr. Approx.* (Dec. 2022), pp. 1–46. ISSN: 1432-0940. DOI: [10.1007/s00365-022-09606-0](https://doi.org/10.1007/s00365-022-09606-0).
- [3] F.W.J. Olver, A.B.O. Daalhuis, D.W. Lozier, B.I. Schneider, R.F. Boisvert, C.W. Clark, B.R. Miller and B. V. Saunders. *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov>. Dec. 2018. DOI: [10.1023/A:1022915830921](https://doi.org/10.1023/A:1022915830921). (Visited on 18/08/2023).




Figure: Surface plot of the total potential energy $E(\alpha, R)$, the blue scatter points represent the fit $E(\alpha, R; \mathbf{x}_{\min})$.