

Abstract

To be written.

This MMSC thesis will further explore general kernel spectral methods for finding equilibrium measures where initial progress made in [Gutleb, Carrillo and S. Olver 2020](#) and [Gutleb, Carrillo and S. Olver 2021](#).

Keywords: Equilibrium Measures


Languages: C++, Julia, Python

Contents

1	Introduction	3
2	Spectral Method	6
2.1	Derivation of Operator	6
3	Conclusion	8
	Acronyms, Definitions and Theorems	9
	Bibliography	10
	List of Figures and Tables	11
A	Supplemental Proofs	12

Chapter 1

Introduction



(a) $N = 8$ particles interacting with one another through the potential $K(r)$. **(b)** Plot of attractive-repulsive potential functions $K(r) = \frac{r^\alpha}{\alpha} - \frac{r^\beta}{\beta}$ for different α, β .

Cf. Figure 1.1a and Figure 1.1b.

All plots and figures in this thesis were generated using the Makie visualisation tool (Danisch and Krumbiegel 2021), an open-source package available for the Julia computing language (Bezanson et al. 2017).

Just Notes

This chapter's purpose is the collection of notes, and it will not be included in the final dissertation.

Special Functions we like

Pochhammer's falling symbol $(x)_n := \prod_{k=0}^{n-1} (x - k)$.

Pochhammer's rising symbol $(x)^n := \prod_{k=0}^{n-1} (x + k)$.

Generalised hypergeometric series

$${}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; z) := \sum_{n=0}^{\infty} \frac{(a_1)_n \cdots (a_p)_n}{(b_1)_n \cdots (b_q)_n} \frac{z^n}{n!}.$$

(Gaussian) Hypergeometric function

$${}_2F_1(a, -n; c; z) = \sum_{j=0}^n (-1)^j \binom{n}{j} \frac{(a)_j}{(c)_j} z^j.$$

(A special case of the hypergeometric series with $p = 2$, $q = 1$).

Jacobi (=hypergeometric) polynomials

$$P_n^{(\alpha, \beta)}(z) := \frac{(\alpha + 1)_n}{n!} {}_2F_1\left(-n, 1 + \alpha + \beta + n; \alpha + 1; \frac{1}{2}(1 - z)\right).$$

Gegenbauer (=ultraspherical) polynomials

$$C_n^{(\lambda)}(z) := \frac{(2\lambda)_n}{n!} {}_2F_1\left(-n, 2\lambda + n; \lambda + \frac{1}{2}; \frac{1 - z}{2}\right) = \frac{(2\lambda)_n}{(\lambda + \frac{1}{2})_n} P_n^{(\lambda-1/2, \lambda-1/2)}(x).$$

They satisfy a three-term recurrence relation (as all orthogonal polynomials do!)

$$\begin{aligned} C_0^{(\lambda)}(x) &= 1 \\ C_1^{(\lambda)}(x) &= 2\lambda x \\ (n+1)C_{n+1}^{(\lambda)}(x) &= 2(n+\lambda)x C_n^{(\lambda)}(x) - (n+2\lambda-1)C_{n-1}^{(\lambda)}(x). \end{aligned}$$

From Wikipedia: In spectral methods for solving differential equations, if a function is expanded in the basis of Chebyshev polynomials and its derivative is represented in a Gegenbauer/ultraspherical basis, then the derivative operator becomes a diagonal matrix, leading to fast banded matrix methods for large problems (S. Olver and Townsend 2013).

Three-term recurrence relationship F. Olver et al. 2018, p. 18.9.1:

$$xC_n^{(\lambda)}(x) = \frac{(n+2\lambda-1)}{2(n+\lambda)}C_{n-1}^{(\lambda)}(x) + \frac{n+1}{2(n+\lambda)}C_{n+1}^{(\lambda)}(x). \quad (1.1)$$

1.0.1 Theorem: Two term recurrence of Q^α

The integral operator

$$Q^\alpha[u](x) = \int_{-1}^1 |x-y|^\alpha u(y) dy$$

satisfies a two-term recurrence relationship when acting on the ultraspherical polynomials $C_n^{(\lambda)}(y)$ with weight $w(y) = (1-y^2)^{\lambda-\frac{1}{2}}$ such that

$$xQ^\alpha[wC_n^{(\lambda)}](x) = \kappa_1 Q^\alpha[wC_{n-1}^{(\lambda)}](x) + \kappa_2 Q^\alpha[wC_{n+1}^{(\lambda)}](x),$$

where $n \geq 2$ and with the constants

$$\begin{aligned} \kappa_1 &= \frac{(n-\alpha-1)(2\lambda+n-1)}{2n(\lambda+n)}, \\ \kappa_2 &= \frac{(n+1)(2\lambda+n+\alpha+1)}{2(\lambda+n)(2\lambda+n)}. \end{aligned}$$

Chapter 2

Spectral Method

2.0.1 Theorem: Integration Theorem that needs a name

On the d -dimensional unit ball B_1 the power law potential, with power $\alpha \in (-d, 2 + 2m - d)$, $m \in \mathbb{N}_0$ and $\beta > -d$, of the n -th weighted radial Jacobi polynomial

$$(1 - |y|^2)^{m - \frac{\alpha + d}{2}} P_n^{(m - \frac{\alpha + d}{2}, \frac{d - 2}{2})}(2|y|^2 - 1)$$

reduces to a Gaussian hypergeometric function as follows:

$$\begin{aligned} & \int_{B_1} |x - y|^\beta (1 - |y|^2)^{m - \frac{\alpha + d}{2}} P_n^{(m - \frac{\alpha + d}{2}, \frac{d - 2}{2})}(2|y|^2 - 1) dy \\ &= \frac{\pi^{d/2} \Gamma(1 + \frac{\beta}{2}) \Gamma(\frac{\beta + d}{2}) \Gamma(m + n - \frac{\alpha + d}{2} + 1)}{\Gamma(\frac{d}{2}) \Gamma(n + 1) \Gamma(\frac{\beta}{2} - n + 1) \Gamma(\frac{\beta - \alpha}{2} + m + n + 1)} {}_2F_1 \left(n - \frac{\beta}{2}, \quad -m - n + \frac{\alpha - \beta}{2}; \frac{d}{2}; |x|^2 \right). \end{aligned}$$

2.1 Derivation of Operator

Considering the operator $\hat{Q}^\beta[\rho]$ as in [[Theorem 2.16]], from the [[Ansatz]] $\rho(x)$ we have

$$\hat{Q}^\beta(x) = \sum_{k=1}^N \rho_k \int \|x - y\|^\beta (1 - \|y\|^2)^a P_k^{(a,b)}(2\|y\|^2 - 1) dy$$

Yeah what is that




Figure 2.1: The attractive and repulsive operators

Chapter 3

Conclusion

In the present thesis, we explored the interesting realm of particle-particle interactions. Next to the written part, the reader will find an implementation of the particle simulator, including a Graphical User Interface (GUI), as well as the numerical solver.

Acronyms, Definitions and Theorems

GUI Graphical User Interface

8

Definitions

Theorems

1.0.1	Two term recurrence of Q^α	5
2.0.1	Integration Theorem that needs a name	6

Lemmata

Remarks

Bibliography

- Bezanson, Jeff, Alan Edelman, Stefan Karpinski and Viral B Shah (2017). ‘Julia: A fresh approach to numerical computing’. In: *SIAM review* 59.1, pp. 65–98. URL: <https://doi.org/10.1137/141000671>.
- Danisch, Simon and Julius Krumbiegel (2021). ‘Makie.jl: Flexible high-performance data visualization for Julia’. In: *Journal of Open Source Software* 6.65, p. 3349. DOI: [10.21105/joss.03349](https://doi.org/10.21105/joss.03349). URL: <https://doi.org/10.21105/joss.03349>.
- Gutleb, Timon S., José A. Carrillo and Sheehan Olver (Oct. 2020). ‘Computing Equilibrium Measures with Power Law Kernels’. In: *arXiv*. DOI: [10.1090/mcom/3740](https://doi.org/10.1090/mcom/3740). eprint: [2011.00045](https://arxiv.org/abs/2011.00045).
- (Sept. 2021). ‘Computation of Power Law Equilibrium Measures on Balls of Arbitrary Dimension’. In: *arXiv*. DOI: [10.1007/s00365-022-09606-0](https://doi.org/10.1007/s00365-022-09606-0). eprint: [2109.00843](https://arxiv.org/abs/2109.00843).
- Olver, F.W.J., A.B.O. Daalhuis, D.W. Lozier, B.I. Schneider, R.F. Boisvert, C.W. Clark, B.R. Miller and B. V. Saunders (eds.) (Dec. 2018). *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov>. (Visited on 11/11/2020).
- Olver, Sheehan and Alex Townsend (Aug. 2013). ‘A Fast and Well-Conditioned Spectral Method’. In: *SIAM Rev.* URL: <https://epubs.siam.org/doi/10.1137/120865458>.

List of Figures and Tables

List of Figures

2.1 The attractive and repulsive operators	7
--	---

List of Tables

Appendix A – Supplemental Proofs