

A tale of two primes

Bert

November 30, 2025

Some text about the article.

Definition

This is a definition.

Some text about the article.

Definition

This is a definition.

Example

This is an example of a block.

Some text about the article.

Definition

This is a definition.

Example

This is an example of a block.

Euclid's theorem

This is a theorem.

1 Element

- 1 Element
- 2 Element

- 1 Element
- 2 Element
- 3 Element

Sets

A **set** is a collection of objects.

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z .

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ".

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

\mathbb{Q}

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

$$\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

$$\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\} \quad (\text{"rational numbers"})$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

$$\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\} \quad (\text{"rational numbers"})$$

$$\mathbb{R}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

$$\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\} \quad (\text{"rational numbers"})$$

$$\mathbb{R} = \{\text{decimal numbers}\}$$

Sets

A **set** is a collection of objects. For example:

$$Z = \{\text{cow}, \text{pig}, \text{elephant}\}.$$

We call the objects in Z the **elements** of Z . We write

$$\text{cow} \in Z$$

with "cow is an element of Z ". Frequently encountered sets are

$$\mathbb{N} = \{1, 2, 3, \dots\} \quad (\text{"natural numbers"})$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} \quad (\text{"integer numbers"})$$

$$\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\} \quad (\text{"rational numbers"})$$

$$\mathbb{R} = \{\text{decimal numbers}\} \quad (\text{"real numbers"})$$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$f'(x)$$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$f'(x) = g'(x) \cdot h(x) +$$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$f'(x) = g'(x) \cdot h(x) + g(x) \cdot h'(x)$$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$\begin{aligned} f'(x) &= g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 2x \cdot \sin(x) + \end{aligned}$$

- $[a, b]$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$\begin{aligned} f'(x) &= g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 2x \cdot \sin(x) + x^2 \cdot \cos(x). \end{aligned}$$

- $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\},$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$\begin{aligned} f'(x) &= g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 2x \cdot \sin(x) + x^2 \cdot \cos(x). \end{aligned}$$

- $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\},$
- (a, b)

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$\begin{aligned} f'(x) &= g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 2x \cdot \sin(x) + x^2 \cdot \cos(x). \end{aligned}$$

- $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\},$
- $(a, b) = \{x \in \mathbb{R} : a < x < b\},$

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$\begin{aligned} f'(x) &= g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 2x \cdot \sin(x) + x^2 \cdot \cos(x). \end{aligned}$$

- $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\},$
- $(a, b) = \{x \in \mathbb{R} : a < x < b\},$
- (a, ∞)

The derivative of $f(x) = g(x) \cdot h(x)$, with $g(x) = x^2$ and $h(x) = \sin(x)$ equals

$$\begin{aligned} f'(x) &= g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 2x \cdot \sin(x) + x^2 \cdot \cos(x). \end{aligned}$$

- $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$,
- $(a, b) = \{x \in \mathbb{R} : a < x < b\}$,
- $(a, \infty) = \{x \in \mathbb{R} : x > a\}$.

A helpful tool for mathematicians.