

# Beamer example - CSD rework

Usage of the theme U{iBCSD

**Veljko Lipovac**



UNIVERSITY OF BERGEN

*Center for Modeling of Coupled Subsurface Dynamics*

VISTA



# Table of contents

## 1 Overview

## 2 Branding

- Logos

## 3 Mathematics

- Theorem
- Example

## 4 Highlighting

## 5 Lists

## 6 Effects

## 7 References

# UiB Emblem

For displaying the the UiB emblem at the lower right corner, use the command `\showlogo` outside the frame environment.

Use `\hidelogo` to remove it from a frame.



# UiB-CSD logo

For displaying the the UiB emblem at the lower right corner, use the command `\showlogo` outside the frame environment.

Use `\hidelogo` to remove it from a frame.

For displaying the the UiB emblem with CSD lettering, use the command `\showCSDlogo` outside the frame environment.



# UiB-CSD and VISTA logo

For displaying the the UiB emblem with CSD lettering and additionally the VISTA logo, use the command `\showVISTACSDlogo` outside the frame environment.

Logos are hidden by default from the beginning.

The last logo command will hold for all following frames until new command is given.



# Mathematics


## Theorem (Fermat's little theorem)

*For a prime  $p$  and  $a \in \mathbb{Z}$  it holds that  $a^p \equiv a \pmod{p}$ .*

## Proof.

The invertible elements in a field form a group under multiplication. In particular, the elements

$$1, 2, \dots, p-1 \in \mathbb{Z}_p$$

form a group under multiplication modulo  $p$ . This is a group of order  $p-1$ . For  $a \in \mathbb{Z}_p$  and  $a \neq 0$  we thus get  $a^{p-1} = 1 \in \mathbb{Z}_p$ . The claim follows. 

# Mathematics

## Example

The function  $\varphi: \mathbb{R} \rightarrow \mathbb{R}$  given by  $\varphi(x) = 2x$  is continuous at the point  $x = \alpha$ , because if  $\epsilon > 0$  and  $x \in \mathbb{R}$  is such that  $|x - \alpha| < \delta = \frac{\epsilon}{2}$ , then

$$|\varphi(x) - \varphi(\alpha)| = 2|x - \alpha| < 2\delta = \epsilon.$$

**Highlighting**



# Highlighting

Sometimes it is useful to highlight certain words in the text.

## Important message

If a lot of text should be highlighted, it is a good idea to put it in a box.

It is easy to match the colour theme.

# Lists

- Bullet lists are marked with a red box.
- 1 Numbered lists are marked with a white number inside a red box.

Description highlights important words with red text.

Items in numbered lists like 1 can be referenced with a red box.

## Example

- Lists change colour after the environment.

# Effects

## 1 Effects that control

Use textblock for arbitrary placement of objects.



# Effects

- 1 Effects that control
- 2 when text is displayed

Use **textblock** for arbitrary placement of objects.

## Theorem

*This theorem is only visible on slide number 2.*

# Effects

- 1 Effects that control
- 2 when text is displayed
- 3 are specified with `<>` and a list of slides.

Use **textblock** for arbitrary placement of objects.



# Effects

- 1 Effects that control
- 2 when text is displayed
- 3 are specified with `<>` and a list of slides.

Use **textblock** for arbitrary placement of objects.

# Effects

- 1 Effects that control
- 2 when text is displayed
- 3 are specified with `<>` and a list of slides.

Use **textblock** for arbitrary placement of objects.

It creates a box with the specified width (here in a percentage of the slide's width) and upper left corner at the specified coordinate (x, y) (here x is a percentage of width and y a percentage of height).

# References I



Hartshorne, R.  
*Algebraic Geometry*.  
Springer-Verlag, 1977.



Helsø, M.  
'Rational quartic symmetroids'.  
*Adv. Geom.*, 20(1):71–89, 2020.



Helsø, M. and Ranestad, K.  
*Rational quartic spectrahedra*, 2018.  
<https://arxiv.org/abs/1810.11235>

- ▶ Atiyah, M. and Macdonald, I.

*Introduction to commutative algebra*.  
Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills,  
Ont., 1969



# References II

[5] Artin, M.

‘On isolated rational singularities of surfaces’.

*Amer. J. Math.*, 80(1):129–136, 1966.