

# HW4

Christian Sallinger

20 4 2021

## 1. The mean of independent normal distributions

(a) Show that the moment generating function (mgf) of  $X \sim \mathcal{N}(\mu, \sigma^2)$  is of the form

$$M_X(t) = e^{\mu t + \frac{\sigma^2 t^2}{2}}.$$

(b) Let  $X \sim \mathcal{N}(\mu, \sigma^2)$  and let  $Y = aX + b$  with fixed real constants  $a$  and  $b$ . Show that  $Y \sim \mathcal{N}(a\mu + b, a^2\sigma^2)$ .

(c) Let  $X_1, \dots, X_n$  be independent identically distributed random variables with  $X_1 \sim \mathcal{N}(\mu, \sigma^2)$ . Show that the mean  $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$  is also normally distributed and  $\bar{X} \sim \mathcal{N}(\mu, \frac{\sigma^2}{n})$ .

**Solution:**

The mgf is given via  $M_X(t) = \mathbb{E}(e^{tx}) = \int_{-\infty}^{\infty} e^{tx} f_X(x) dx$ . We will show this first for  $Z \sim \mathcal{N}(0, 1)$ :

$$M_Z(t) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{tx} e^{-x^2/2} dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{(2tx-x^2)/2} dx = \dots$$

We can substitute  $y = x - t$  and get

$$\dots = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{(2t(y+t)-(y+t)^2)/2} dy = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{(t^2-y^2)/2} dy = e^{t^2/2}$$

Where the last equality follows from the well known calculus result  $\int_{\mathbb{R}} e^{x^2/2} dx = \sqrt{2\pi}$ .

We will show in (b), that  $\sigma Z + \mu \sim \mathcal{N}(\mu, \sigma^2)$ . We can now use the properties of the mfg to show the wanted result:

$$M_{\sigma Z + \mu}(t) = e^{\mu t} M_Z(\sigma t) = e^{\mu t} e^{\frac{\sigma^2 t^2}{2}}$$

which is exactly what we wanted to show.

(b) To show this we use our transformation theorem: The function

$$\begin{aligned} g : (-\infty, \infty) &\rightarrow (-\infty, \infty) \\ x &\mapsto ax + b \end{aligned}$$

is invertible with differentiable inverse  $h(x) = \frac{x-b}{a}$  as long as  $a \neq 0$ . Since  $Y$  would be constant for  $a = 0$  we will not consider this case anyway. For the pdf of  $Y$  we now get:

$$\begin{aligned}
f_Y(y) &= f_X(h(y))|h'(y)| = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{((y-b)/a-\mu)^2}{2\sigma^2}} \frac{1}{a} \\
&= \frac{1}{\sqrt{2\pi a^2 \sigma^2}} e^{-\frac{(y-b-\mu a)^2}{2\sigma^2 a^2}} = \frac{1}{\sqrt{2\pi a^2 \sigma^2}} e^{-\frac{(y-(\mu a+b))^2}{2\sigma^2 a^2}}
\end{aligned}$$

So we can see that  $Y \sim \mathcal{N}(a\mu + b, a^2\sigma^2)$  really holds.

(c) Since the random variables are independent we know from the lecture that the mgf of  $\bar{X}$  is given by

$$M_{\bar{X}}(t) = \left(M_{X_1}\left(\frac{t}{n}\right)\right)^n.$$

We have already seen in (a) what the mgf of  $X_1$  looks like, so we can plug it in here:

$$\left(M_{X_1}\left(\frac{t}{n}\right)\right)^n = \left(e^{t\mu/n + \frac{\sigma^2 t^2}{2n^2}}\right)^n = e^{t\mu + \frac{\sigma^2 t^2}{n}}$$

Since the mgf exists in a neighbourhood of zero the distribution is uniquely determined by its mgf and as we can see this mgf corresponds to a normal distribution with mean  $\mu$  and variance  $\frac{\sigma^2}{n}$ , which is exactly what we wanted to show.

## 2. Sum of two independent distributions

(a) Let  $X \sim \mathcal{P}(\lambda_1)$  and  $Y \sim \mathcal{P}(\lambda_2)$  be two independent Poisson random variables. Show that

$$X + Y \sim \mathcal{P}(\lambda_1 + \lambda_2).$$

(b) Let  $U$  and  $V$  be two independent random variables with exponential distribution  $\exp(\lambda)$ . Show that

$$\begin{aligned}
U + V &\sim \text{Gamma}(2, \lambda) \quad \text{and} \\
\min\{U, V\} &\sim \exp(2\lambda).
\end{aligned}$$

*Hint:* It is useful to use moment generating functions. Recall, the pdf of a random variable  $X \sim \text{Gamma}(\alpha, \frac{1}{\beta})$  is

$$f(x) = \begin{cases} \frac{x^{\alpha-1} e^{-\frac{x}{\beta}}}{\Gamma(\alpha)\beta^\alpha}, & x > 0 \\ 0, & x \leq 0 \end{cases}$$

and its mgf is of the form  $\left(\frac{1}{1-\beta t}\right)^\alpha$  for  $t \leq \frac{1}{\beta}$ . Particularly, the pdf of a random variable  $X \sim \exp(\lambda) = \text{Gamma}(1, \frac{1}{\lambda})$  is of the form

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x > 0 \\ 0, & x \leq 0 \end{cases}$$

**Solution:** (a) We recall that the pmf of a random variable  $Z \sim \mathcal{P}(\lambda)$  is given by

$$f_Z(k) = \frac{\lambda^k}{k!} e^{-\lambda}.$$

Now we can use the convolution formula for discrete random variables:

$$\begin{aligned}
 f_{X+Y}(k) &= \sum_{y \in \{0, \dots, k\}} f_Y(y) f_X(k-y) = \sum_{y \in \{0, \dots, k\}} \frac{\lambda_2^y}{y!} e^{-\lambda_2} \frac{\lambda_1^{k-y}}{(k-y)!} e^{-\lambda_1} \\
 &= e^{-(\lambda_1 + \lambda_2)} \sum_{y \in \{0, \dots, k\}} \frac{\lambda_2^y \lambda_1^{k-y}}{y!(k-y)!} = e^{-(\lambda_1 + \lambda_2)} \frac{1}{k!} \sum_{y \in \{0, \dots, k\}} \binom{k}{y} \lambda_2^y \lambda_1^{k-y} \\
 &= e^{-(\lambda_1 + \lambda_2)} \frac{(\lambda_1 + \lambda_2)^k}{k!}
 \end{aligned}$$

(b) Here we use the convolution formula for continuous random variables:

$$\begin{aligned}
 f_{U+V}(z) &= \int_{\mathbb{R}} f_U(z-y) f_V(y) dy = \lambda^2 \int_{\mathbb{R}} e^{-\lambda(z-y)} e^{-\lambda y} 1_{(0, \infty)}(z-y) 1_{(0, \infty)}(y) dy \\
 &= \lambda^2 \int_0^z e^{-\lambda z} dy = \lambda^2 z e^{-\lambda z}, \quad z > 0
 \end{aligned}$$

This is what we wanted to show, since  $\Gamma(2) = 1$ . For the second part we use the general formula for the cdf of the minimum of two independent, identically distributed random variables from last week. First we define  $Z = \min\{U, V\}$ . Also recall that the cdf for a random variable  $x \sim \exp(\lambda)$  is given by

$$F_X(x) = \begin{cases} 1 - e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases}$$

So we get:

$$F_Z(z) = 2F_U(z) - F_U(z)^2 = 2(1 - e^{-\lambda z}) - (1 - e^{-\lambda z})^2 = 2 - 2e^{-\lambda z} - 1 + 2e^{-\lambda z} - e^{-2\lambda z} = 1 - e^{-2\lambda z}, \quad z \geq 0$$

Comparing with the cdf above we see that  $\min\{U, V\} \sim \exp(2\lambda)$  really holds.

### 3. Real roots

Let  $A, B$  and  $C$  be independent random variables, uniformly distributed on  $(0, 1)$ .

(a) What is the probability that the quadratic equation  $Ax^2 + Bx + C = 0$  has real roots?

(b) Consider the following code in R. What does it do and how is it related to your solution in part (a)?

```
n = 10000
a = runif(n)
b = runif(n)
c = runif(n)
sum(b^2 > 4*a*c)/n
```

```
## [1] 0.2529
```

*Hint:* In HW2/ex. 3(b) we showed that if  $X$  has uniform  $(0, 1)$  distribution then  $-\log X$  has exponential distribution  $\exp(1)$ . In an analogue way, one can prove that  $-s \log X \sim \exp(\frac{1}{s})$  for any  $s > 0$ . Also, in HW4/ex. 2(b) we proved that the sum of two independent exponential distributions is a gamma distribution. Namely, if  $X \sim \exp(1)$  and  $Y \sim \exp(1)$  are independent then  $X + Y \sim \text{Gamma}(2, 1)$ .

**Solution:**

Using the "Grosse Lösungsformel" we see that the quadratic equation has real roots iff  $B^2 \geq 4AC$  holds. By use of the monotony of the logarithm as well as random variables  $Z \sim \exp(\frac{1}{2})$ ,  $U, V \sim \exp(1)$  and  $G \sim \text{Gamma}(2, 1)$  we follow the hint:

$$\begin{aligned}
\mathbb{P}(B^2 \geq 4AC) &\iff \mathbb{P}(\log(B^2) \geq \log(4AC)) \\
&\iff \mathbb{P}(-2\log(B) \leq -\log(4AC)) \iff \mathbb{P}(-2\log(B) \leq -\log(4) - \log(A) - \log(C)) \\
&\iff \mathbb{P}(Z - (-\log(A) - \log(C)) \leq -\log(4)) \iff \mathbb{P}(Z - (U + V) \leq -\log(4)) \\
&\iff \mathbb{P}(Z - G \leq -\log(4))
\end{aligned}$$

To get the pdf of  $Z - G$  we use the convolution formula:

$$\begin{aligned}
f_{Z+(-G)}(z) &= \int_{\mathbb{R}} f_Z(z-y) f_G(-z) dy = -\frac{1}{2} \int_{\mathbb{R}} e^{-\frac{z-y}{2}} y e^y 1_{(-\infty, 0)}(y) 1_{[0, \infty)}(z-y) dy = -\frac{1}{2} \int_{-\infty}^{\min\{z, 0\}} e^{-\frac{z-y}{2}} y e^y dy \\
&= \begin{cases} -\frac{(6z-4)e^z}{18}, & z < 0 \\ \frac{2e^{\frac{z}{2}}}{9}, & z \geq 0 \end{cases}
\end{aligned}$$

Now we can calculate the probability:

$$\mathbb{P}(Z - G \leq -\log(4)) = F_{Z-G}(-\log(4)) = \int_{-\infty}^{-\log(4)} f_{Z-G}(z) dz \approx 0.2544$$

- (b) The command "runif" creates a vector of length  $n$  with samples taken from an uniformly distributed random variable. In the sum we get 1 if  $b_j^2 > 4a_j c_j$  for the  $j$ -th entry of the vectors and 0 otherwise. So in the sum we have the total number of times the inequality holds for the  $n$ -samples. Dividing by  $n$  then gives the mean. The value gives us a good approximation of the real probability because by the strong law of large numbers it converges to it. To see this we define a random variable  $X$  that is 1 if  $B^2 > 4AC$  and 0 otherwise. The expectation is

$$\mathbb{E}(X) = 0 \cdot \mathbb{P}(X = 0) + 1 \cdot \mathbb{P}(X = 1) = \mathbb{P}(B^2 > 4AC)$$

#### 4. Sum and average

Let  $X$  be a random variable with  $\mathcal{N}(5, 2^2)$ . Let  $X_1, X_2, \dots, X_{50}$  be independent identically distributed copies of  $X$ . Let  $S$  be their sum and  $\bar{X}$  their average, i.e.

$$S = X_1 + \dots + X_{50} \quad \text{and} \quad \bar{X} = \frac{1}{50}(X_1 + \dots + X_{50}).$$

- (a) Plot the density and the distribution function for  $X$  using R.  
(b) What are the expectation and the standard deviation of  $S$  and  $\bar{X}$ ?  
(c) Generate a sample of 50 numbers from  $\mathcal{N}(5, 2^2)$ . Plot the histogram for this sample. Do the same for a sample of 500 numbers from  $\mathcal{N}(5, 2^2)$ .


**Solution:** (a)

```

x <- seq(-4, 14, 0.1)
y <- dnorm(x, 5, 4)
plot(x, y, type="l", main = "pdf of the normal-distribution", xlab = "x", ylab = "value of pdf"
     , xaxp = c(-4, 14, 18))
abline(v = 5)


```

### pdf of the normal-distribution



```
z <- pnorm(x,5,4)
plot(x,z,type = "l", main = "cdf of the normal-distribution",xlab = "x", ylab = "Value of cdf"
,xaxp = c(-4, 14, 18))
```

### cdf of the normal-distribution



(b) We first calculate the expectations:

$$\mathbb{E}(S) = \sum_{i=1}^{50} \mathbb{E}(X_i) = 50 \cdot 5 = 250$$

$$\mathbb{E}(\bar{X}) = \frac{1}{50} \mathbb{E}(S) = 5$$


For the variances, since the random variables are independent, we get:

$$\mathbb{V}ar(S) = \sum_{i=1}^{50} \mathbb{V}ar(X_i) = 200$$


$$\mathbb{V}ar(\bar{X}) = \left(\frac{1}{50}\right)^2 \mathbb{V}ar(S) = \frac{200}{2500} = \frac{8}{100}$$

(c)

```
x = rnorm(50,5,4)
hist(x, main = "histogram of 50 samples")
```



```
y = rnorm(500,5,4)
hist(y, main = "histogram of 500 samples")
```



## 5. Central Limit Theorem

Let  $\bar{X}_1$  and  $\bar{X}_2$  be the means of two independent samples of size  $n$  from the same population with variance  $\sigma^2$ . Use the Central limit theorem to find a value for  $n$  so that

$$P(|\bar{X}_1 - \bar{X}_2| < \frac{\sigma}{50}) \approx 0.99.$$

Justify your calculations.

**Solution:**

We first define  $\bar{X}_n := \bar{X}_1 - \bar{X}_2 = \frac{1}{n} \sum_{i=1}^n (X_{1,i} - X_{2,i})$ . We now want to calculate the mean of  $X_i$  for some  $i$ :

$$\mathbb{E}(X_i) = \mathbb{E}(X_{1,i}) - \mathbb{E}(X_{2,i}) = \mu - \mu = 0$$

Since they are independent samples we can also calculate the variance rather easily.

$$\text{Var}(X_i) = \text{Var}(X_{1,i} - X_{2,i}) = \text{Var}(X_{1,i}) + \text{Var}(X_{2,i}) = 2\sigma^2$$

The central limit theorem now states that the cdf of  $\sqrt{n} \cdot \frac{\bar{X}_n}{\sqrt{2}\sigma}$  converges to the cdf of the standard normal distribution. So we can use  $\Phi$  as an approximation.



$$\begin{aligned}
\mathbb{P}(|\bar{X}_1 - \bar{X}_2| < \frac{\sigma}{50}) &= \mathbb{P}(-\frac{\sigma}{50} < \bar{X}_n < \frac{\sigma}{50}) = \mathbb{P}(-\frac{1}{50} < \frac{\sqrt{2}}{\sigma\sqrt{2}}\bar{X}_n < \frac{1}{50}) \\
&= \mathbb{P}(-\frac{\sqrt{n}}{50\sqrt{2}} < \frac{\sqrt{n}}{\sigma\sqrt{2}}\bar{X}_n < \frac{\sqrt{n}}{50\sqrt{2}}) \approx \Phi\left(\frac{\sqrt{n}}{50\sqrt{2}}\right) - (1 - \Phi\left(\frac{\sqrt{n}}{50\sqrt{2}}\right)) = 2\Phi\left(\frac{\sqrt{n}}{50\sqrt{2}}\right) - 1
\end{aligned}$$

Now we want to find  $n$  so that

$$2\Phi\left(\frac{\sqrt{n}}{50\sqrt{2}}\right) - 1 \approx 0.99 \iff \Phi\left(\frac{\sqrt{n}}{50\sqrt{2}}\right) \approx 0.995 \iff \frac{\sqrt{n}}{50\sqrt{2}} = 2.58 \iff n = 33282$$