Électrochimiothérapie des métastases rachidiennes et osseuses complexes : aspects techniques, balistique et sécurité neurologique

Titre original : Electrochemotherapy in Spine Metastases: A Case Series Focused on Technical Aspects, Surgical Strategies and Results

Auteurs : Angelini A, D'Amico A et al. | **Revue :** *Diagnostics (Basel, Switzerland)* (**IF : 3.0**) | **Date :** Avr. 2024 | **PMID :** 38732352 | **Modalité :** Scanner TDM

1. Pertinence Clinique & Lien avec l'Agenda Interventionnel (Cochin)

Fournit les abaques techniques de placement des électrodes et de sécurité neurologique lors des procédures d'ECT réalisées en Radiologie B en collaboration avec l'HEGP.

2. Synthèse Clinique en Français

Résumé exécutif : Étude technique détaillant la mise en œuvre de l'électrochimiothérapie (ECT) pour les métastases osseuses à haut risque neurologique ou mécanique. Grâce à son mécanisme non thermique, l'ECT permet de traiter des lésions au contact direct d'éléments nerveux ou articulaires là où la radiofréquence et la cryoablation seules exposeraient à des complications.

Analyse détaillée : Cet article analyse les paramètres pratiques d'implantation des électrodes d'électroporation dans l'os pathologique, la vérification de la couverture du champ électrique et les résultats cliniques sur la douleur (échelle EVA) et le contrôle tumoral local en imagerie. L'étude souligne l'importance de la synchronisation anesthésique (curarisation profonde au moment des impulsions électriques et limitation de la FiO2 sous bléomycine) et confirme l'absence de nécrose thermique des tissus sains environnants.

3. Protocole Technique & Paramètres Opératoires

Planification TDM de la géométrie des aiguilles • Curarisation per-procédurale pour limiter les secousses musculaires lors des impulsions • FiO2 < 30 % pendant l'anesthésie générale (prévention de la toxicité pulmonaire de la bléomycine) • Contrôle IRM à 6–12 semaines.

4. Points Clés à Retenir pour la Pratique

- Précaution anesthésique essentielle : maintenir une FiO2 minimale (< 30 %) lors de l'AG après administration de bléomycine.
- Permet une destruction tumorale sélective au contact immédiat des structures nerveuses ou cutanées (pied, cheville, épidure).
- Diminution rapide et durable de l'EVA douleur et réduction de la consommation d'opioïdes dès le premier mois.
- Peut être associée dans le même temps opératoire à une cryoablation centrale (debulking) avec couronne périphérique d'ECT.

5. Original PubMed Abstract (Texte Intégral de l'Abstract)

Metastases are complications of primary tumors due to prolonged cancer survival and have become an important issue for oncological patients and the most frequent cause of death and disability. Bone metastases occur at a later stage of cancer disease, and the spine is the most frequent site. To date, the aim of the treatment of metastases remains to be the control of disease and provide a satisfactory quality of life. The decision making of treatment is influenced by several factors such as the status of the primary disease, the

number of metastases, site involvement, and the performance status of the patients. For this reason, the treatment of metastases is challenging and undergoes constant development. Therefore, alternative techniques with respect to surgery, which is the first option but not always practicable, and radiochemotherapy are attractive. Lately, electrochemotherapy has emerged as an innovative method for treating various primary and metastatic solid tumors, showing promising outcomes in terms of inducing tumor tissue necrosis and alleviating symptoms. This technique uses electric pulses to increase the uptake of chemotherapy by tumor cells. Despite the initial enthusiasm and good results in the treatment of bone tumors, relatively few papers have described its use in spine metastases. Therefore, we conducted a systemic review of this intriguing topic while also reporting our experience in the use of electrochemotherapy for the treatment of spine metastases.

---

6. Full-Text Sections (Europe PMC Open Access JATS XML)

1. Introduction

Bone is the third most common site of metastases following the lung and liver, and the spine is the most common site for bone metastasis [1]. The highest incidence of spine metastases occurs from 40 to 65 years of age [2]. The thoracic spine is the most common site of involvement (70%), followed by the lumbar spine (20%), the cervical spine, and the sacrum [3]. The vertebral body is involved in 80% of cases, with posterior elements being affected in 20%, and most metastases are osteolytic (95%) [4,5]. The paramount purpose of spine metastasis treatment is to correct mechanical instability, resolve eventual neurological compression, and improve quality of life [6].

Treatment in spine metastases has changed over the years, and several techniques are described in the literature [6,7,8,9]. In this scenario, recently, electrochemotherapy (ECT) has been demonstrated to be an efficient treatment for spine metastases [10]. This technique combines systemic chemotherapy, usually bleomycin, and electric pulses delivered locally at the tumor site. The intriguing mechanism of this technique is characterized by the temporary permeabilization of cell membranes of tumoral tissue that allow bleomycin to diffuse inside cells and increase its toxicity. It is also used in different settings such as benign tumors of the spine [11]. Several studies have reported the efficacy and feasibility of this technique in the treatment of bone metastasis [12], but papers about this topic focused on spine metastasis are sparse, and to date, a standardized protocol has not been described. Therefore, a systematic review of the literature on the use of ECT in spine metastasis was conducted. Additionally, we report our initial experience in the treatment of spine metastases with ECT.

2. Material and Methods

Systematic review of the literature: Given the sparse evidence about ECT in metastatic spine disease, a systematic literature review was performed according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. The protocol of this review was not registered on prospective registers; however, the systematic review was conducted as follows: A literature search was performed on the PubMed database in October 2023. PubMed was queried without backward limits with the following terms: ((spine) AND (metastasis)) AND ((ECT); (ECT) AND (spine)); ((ECT) and (metastasis)). Reference lists of all publications were also screened for missed results. Exclusion criteria were inaccessibility to full text, non-English and non-Italian language, studies not pertinent to ECT in spine disease by “title/abstract”, and studies lacking relevant clinical data (clinical presentation, demographics, surgical procedure, and radiological findings). Review articles were also excluded.

Data extracted from each study included article type, class of evidence if reported, the spine side involved, the type of primary tumor, ECT technique, and functional outcome at the last follow-up. Clinical outcome was evaluated as pain relief, measured by a VAS score ranging from 0 (no pain) to 10 (worst pain ever) and the American Spinal Injury Association (ASIA) impairment scale [13].

Our protocol and technical aspects: While under general anesthesia, the patient is positioned prone, followed by the execution of a posterior surgical approach involving laminectomy at the targeted level. This approach

ensures sufficient exposure of the pedicle area. Particular attention is paid to avoiding damage to nerve roots and the dural sac. Two stainless steel needle electrodes, each measuring 1.8 mm in diameter with sharpened tips, are then inserted into both pedicles of the affected vertebra, with an additional set of four electrodes placed in the pedicles of the adjacent proximal and distal vertebrae. These placements are guided by fluoroscopy and/or neuronavigation techniques. The electrodes are linked to the independently regulated generator outputs of the Cliniporator Vitae (IGEA, Carpi, Italy). This Cliniporator Vitae unit functions as a pulse generator, featuring six electrically isolated outputs, each capable of delivering up to 3000 V with a maximum current of 50 A. It administers eight rectangular electrical pulses, each lasting 100 ms with a rise time of 1 ms, at a pulse repetition rate of 4 Hz [14]. Systemic bleomycin (Bristol-Myers Squibb, Princeton, NJ, USA) is administered intravenously (i.v.) via push injection, typically over a period of 30 to 45 s, at a dosage of 26 mg. This injection is conducted following electrode connections eight minutes before the delivery of electrical pulses. The administration of bleomycin adheres to the ESOPE protocol, which specifies doses of 10 or 15 mg/m2 [15].

Following an 8 min interval, pulse delivery is initiated, with continuous monitoring to ensure the efficacy of the applied electrical field, set at 1.5 A. Both current and voltage levels are meticulously measured and recorded with a precision exceeding 3%, facilitating precise control over pulse delivery and enabling thorough evaluation post-treatment. Throughout the administration of electrical pulses, the patient undergoes partial curarization.

3. Results

The literature search provided the following results: ((spine) AND (metastasis)) AND ((ECT) 3 results; (ECT) AND (spine)) 5 results; ((ECT) and (metastasis)) 137 results. Through this literature search, we retrieved a total of 145 papers.

After careful screening following the PRISMA flowchart and considering the purpose of this systematic review and the exclusion criteria, only four studies [12,16,17,18] were retrieved and assessed for eligibility (Figure 1; Table 1).

PRISMA flow diagram of the literature review.

(*) See Discussion section for more details. ** One patient died of disease 4 months after treatment and another one after 3 months due to renal failure.

4. Case Series

A 75-year-old female Caucasian patient with no history of oncologic disease presented with swelling on the anterior surface of the left leg. Imaging studies revealed a soft tissue suprafascial mass diagnosed as high-grade myxofibrosarcoma (G3 at FNCLCC classification) upon histopathologic evaluation of percutaneous tru-cut biopsy. Subsequently, the patient underwent surgical excision and soft-tissue reconstruction with a myocutaneous flap and cutaneous Thiersch graft. Local radiotherapy was completed with a dosage of 36 Gy within 6 months postoperatively. Six months after the completion of radiotherapy, the patient presented with suspected micronodularity in both lungs and a small lytic lesion in the L3 vertebra (Figure 2), not associated with low back pain (VAS 1).

Case 1. A 75-year-old female with small lytic metastasis from high-grade myxofibrosarcoma in the L3 vertebra. The lesion was confined to the vertebral body (black arrow) without the involvement of the posterior vertebral wall. White arrow in sagittal view corresponds to the axial level.

A decision was made to conduct a short follow-up re-evaluation, which was agreed upon by the patient. Three months later, a new CT scan revealed that the lesion in the L3 vertebra had increased in volume (Figure 3a,b), and MR imaging showed partial compression of the medullary canal (Figure 3c,d). FDG-PET/CT was performed, showing increased uptake (SUV 12) (Figure 3e). The patient reported a significant worsening of pain, with pain radiating to the left thigh, but there were no ongoing peripheral neurological deficits (ASIA scale E).

Case 1 at 3 months of follow-up: (a) Sagittal and (b) corresponding axial CT scans show a lytic lesion in the L3 vertebra with posterior erosion of the cortex, involving the spinal canal. (c) Sagittal T1-weighted MR imaging with contrast medium shows a homogeneous lesion at the L3 level with mixed hyperintense signal, whereas in (d) sagittal T2-weighted MRI, the lesion appears hypointense. (e) FDG-PET/CT on axial view shows the intense focal area of 18-FDG activity in the L3 vertebral body.

Afterward, the patient was referred to our oncologic spine surgery department, where a percutaneous biopsy of the L3 vertebral body was performed. The results of histopathological examination and immunohistochemistry indicated metastatic high-grade myxofibrosarcoma. Considering the patient's oncologic scores, which suggested the need for locoregional treatment at the L3 level, surgical planning began. Notably, there were no signs of spine instability or significant neurological deficits detected at that time, but the growth of the lesion was concerning.

Subsequently, a wide laminectomy procedure coupled with stabilization utilizing pedicled screws and ECT was proposed as a local/palliative treatment approach. This strategy aimed to slow down rapid disease advancement while minimizing the potential for intolerable morbidity associated with extensive surgical resection. The patient consented to this option after being fully informed. The surgery was performed with the patient positioned prone under the navigation and fluoroscopic guidance (Figure 4).

Intraoperative pictures: (a) positioning of the patient in the operating room. The devices for intraoperative navigation (white arrow) and the electrodes for intraoperative evoked potentials (black arrow) are visible; (b) sterile field; (c) precise alignment of needle electrodes surrounding the metastatic lesion was attained under the guidance of navigation and fluoroscopy; (d) the electrodes were connected to the pulse generator; (e) the distance between each couple of electrodes was registered and (f) reported on the device; (g) after ECT, electrodes were replaced by pedicled screws and laminectomy for spinal decompression was performed; (h) dura covered with bone graft.

Follow-up assessments, comprising both clinical and radiological examinations, were scheduled at one- and four-week post-surgery, followed by three-month intervals thereafter. The evaluation of the patient's reported outcomes using the Visual Analog Scale (VAS) revealed a statistically significant reduction from 10 to 2 ($p < 0.001$). However, no significant changes were noted in the patient's neurological status, and there were no local or general side effects observed either immediately after waking or during postoperative follow-up. Imaging restaging conducted 3 months after surgery (Figure 5) demonstrated no signs of implant loosening, and the size of the tumor at the L3 vertebra remained stable. Unfortunately, the patient passed away at 4 months of follow-up due to respiratory failure caused by the progression of metastatic disease to the lungs.

5. Discussion

Undoubtedly, ECT is an important option to manage primary tumors and metastases [19]. Okino et al. [20,21] and Mir et al. [22] provided initial descriptions of a technique involving the local application of electric current to enhance the uptake of chemotherapy drugs, typically bleomycin, into tumor cells. This approach has proved promising regardless of tumor histology [23]. Recently, ECT has been used in more than 83 centers all over Europe [24], and studies have shown that it can also potentially be used in the treatment of bone tumors and bone metastases. To date, its use in clinical practice as a new option for metastasis management is increasing; however, while several studies have been conducted about this technique in the treatment of bone metastasis, few papers have described its use and results in spine metastases [12,16,17,18]. In fact, from our systematic review, only four articles described the use of ECT in spine metastases [12,16,17,18].

Spine metastases are the most common bone tumors and a frequent occurrence in the spine. Therefore, in managing this challenging issue, ECT would work together with medical therapy, surgery, and chemo- and radiotherapy to enhance the quality of life for these patients. The primary goal of treating patients with spinal metastases is palliative, focusing on improving pain control, which is often the foremost concern for these individuals, while also aiming to preserve spinal function and stability [25]. Traditionally, medical therapy, surgery, chemotherapy, and radiotherapy are valid options for managing oncological patients with a risk–benefit–cost ratio that is not always favorable [26]. For this reason, new strategies of treatment with low

comorbidity are attractive. In this scenario, ECT is a new tissue-sparing treatment with selective destruction of tumor cells. In 2015, Gasbarrini et al. [12] were the first to report the use of ECT in the palliative treatment of spinal metastases. They described a case involving a 51-year-old female with melanoma metastasis in the L5 spine. This option was chosen based on oncological scores [27,28]. The electrodes were placed directly in the L5 vertebral body before decompression. The chemotherapy drug delivered was systemic bleomycin. The procedure was tolerated without side effects. Response assessment included percutaneous vertebral biopsy at 3 months (95% of tumor necrosis occurred) and PET/TC at 6 months. Then, the patient underwent radiotherapy for a suspected local relapse identified on the PET/TC scan. In 2018, Cindri■ et al. [16] proposed a new way to deliver electrical pulses near tumor tissue, describing two cases of spine metastases treated by ECT with a transpedicular approach. The authors demonstrated nearly 100% coverage of tumor volume with the electric field delivered via a transpedicular approach [16]. Moreover, when the tumor was located in the anterior part of the vertebral body, no electroporation damage was observed on spinal cord tissue [16]. In this situation, the electrodes have more distance to the target, and a higher electrical pulse is needed to provide electroporation effects on tumor tissue. This is an important concept when treating spine metastases because the potential damage to the spinal cord and roots needs to be considered. Cornelis et al. [17] reported their experience of percutaneous image-guided ECT in two patients with metastatic epidural spinal cord compression from breast and lung cancer. Both had previously undergone chemotherapy, and radiotherapy was performed to relieve back pain with no good result. Thus, ECT was proposed as an alternative palliative treatment. A transpedicular technique was used, and electric pulses were delivered near tumor tissue by inserting electrodes in the affected vertebral body and at the above and below levels with a CT-guided procedure. Electric pulses were delivered 8 min after the intravenous administration of bleomycin. The treatment was well tolerated, and no side effects occurred. MRI confirmed the tumor's local response, and a biopsy at 3 months of follow-up showed >95% necrosis of tumor tissue [17]. Recently, Deschamps et al. [18] reported the most numerous series of spine metastatic diseases treated with ECT and bleomycin. The authors reported a total of 40 patients with metastatic epidural spinal cord compression. All patients in this series received radiotherapy, and 12 patients had a previous surgery. ECT was performed with the percutaneous image CT-guided technique, as described by Cornelis [17], with pain relief obtained prevalently in the first month after the procedure and neurological improvement in the third month of follow-up [18]. The study's data analysis revealed that only 21 out of 40 patients were evaluated at the three-month follow-up, which dampens the enthusiasm regarding the described results [18].

ECT for spine metastases presents effective outcomes but several inherent risks and challenges. Firstly, the proximity of the spinal cord and nerve roots to metastatic lesions increases the potential for neurological damage during the procedure [29]. This risk necessitates careful planning and precise electrode placement to minimize the likelihood of adverse events. A transpedicular ECT procedure combined with vertebral screw fixation in selected patients with tumor volume contained within the vertebral body minimizes the risk for neural damage [16,17,18]. There are important limits in this technique such as the position and angles of electrodes in pedicles and the difficulty of a follow-up examination of the treated lesion with standard imaging techniques due to pedicle screws [16]. A recent study confirmed that the deflection of needles alters the distribution of the electric field, reducing the efficacy of ECT [30]. This aspect could be overcome by the introduction of long-needle variable electrode geometry [31]. Secondly, spine metastases often involve critical structures such as vertebral bodies and the surrounding vasculature. ECT in this context requires meticulous attention to avoid injury to these structures, which could lead to consequences such as spinal cord compression or vascular compromise. Additionally, the efficacy of ECT may be limited in cases where metastases are located in regions of the spine that are difficult to access or where electrode placement is challenging. This may necessitate alternative treatment approaches or a combination of therapies to achieve optimal outcomes while mitigating risks. Close monitoring and follow-up are essential to detect early recurrence or disease progression and adjust treatment strategies accordingly. Overall, while ECT can be a valuable treatment option for spine metastases, its implementation requires careful consideration of the specific risks and challenges associated with this anatomical location, as well as a multidisciplinary approach involving oncologists, neurosurgeons, and interventional radiologists to ensure patient safety and treatment efficacy.

6. Conclusions

According to the literature review and our experience, ECT enhances the toxicity of bleomycin administrated systematically, changing the permeability of membrane cells. This effect is possible due to the modification of membrane cell polarization induced by electric pulses delivered near the tumor site with several techniques. ECT does not modify the overall survival of oncologic patients but is an alternative option for the treatment of spine metastases. Pain is a paramount issue in bone metastatic patients, and the findings of all studies in this review and our experience concur on the capability of ECT to improve neurological conditions and achieve pain relief.

Indeed, ECT is a safe, feasible, and minimally invasive rescue option for oncological patients with spine metastases and not an alternative to surgical resection; it has been shown to improve pain and neurological symptoms in both the short term and long term.

Author Contributions

Conceptualization, project administration, A.A. and A.D.; data curation and formal analysis, S.P., R.S. and G.B.; supervision, L.D. and P.R.; writing—original draft preparation, A.D. and G.D.R.; writing—review and editing, A.A. and A.D. All authors have read and agreed to the published version of the manuscript.

Informed Consent Statement

Written informed consent has been obtained from the patients to publish this paper at the time of admission to our Institute. However, all pictures have been reported anonymized.