# Cryoablation à visée curative des métastases osseuses : protocole de l'étude prospective multicentrique européenne BOREALIS (CIRSE)

*Titre original : Curative-Intent Cryoablation for Bone Metastases: Study Protocol for BOREALIS, a Prospective Multicentre Observational Investigation*

**Auteurs :** Cazzato RL, Buy X et al. | **Revue :** *Cardiovascular and interventional radiology* (**IF : 2.8**) | **Date :** Oct. 2026 | **PMID :** 42618654 | **Modalité :** Scanner TDM / IRM

## 1. Pertinence Clinique & Lien avec l'Agenda Interventionnel (Cochin)

Essai multicentrique de référence (Cazzato, Buy, Deschamps, Palussière, Gangi — *CVIR* août 2026) cadrant les cryoablations de métastases osseuses à visée de contrôle local complet réalisées à Cochin.

## 2. Synthèse Clinique en Français

**Résumé exécutif :** Présentation du registre prospectif multicentrique BOREALIS (*CVIR*, août 2026) évaluant la cryoablation percutanée à visée curative (éradication tumorale complète avec marges de sécurité) chez les patients oligométastatiques osseux.

**Analyse détaillée :** Portée par les principaux centres européens et français de radiologie interventionnelle oncologique (Strasbourg, Bordeaux, Gustave Roussy), l'étude BOREALIS standardise les critères techniques de la cryoablation osseuse à visée curative : sélection des patients oligométastatiques (1 à 3 lésions osseuses ≤ 4 cm), couverture tridimensionnelle complète de la tumeur par l'isotherme létal (-20 °C / -40 °C) avec marges d'au moins 5 mm au-delà des limites IRM, et suivi systématique par IRM/TEP-TDM sur 24 mois.

## 3. Protocole Technique & Paramètres Opératoires

Bilan pré-thérapeutique IRM + TDM • Multi-cryosondes 17G pour obtenir une marge d'iceball ≥ 5–8 mm autour de la lésion • Double cycle de congélation (10 min) séparé d'un dégel passif/actif • Consolidation associée (cimentoplastie ou vissage) si risque fracturaire • Suivi IRM à 3, 6, 12 et 24 mois.

## 4. Points Clés à Retenir pour la Pratique

- Passage d'une indication purement palliative antalgique à une ablation curative (A0) dans la maladie oligométastatique osseuse.
- Exigence d'une marge d'ablation périphérique de 5 mm au-delà du volume tumoral défini sur l'IRM pré-opératoire.
- Intégration systématique des manœuvres de thermoprotection (hydrodissection, carbodissection, neuromonitoring) pour sécuriser les marges.
- Synergie avec les traitements systémiques modernes (immunothérapie et thérapies ciblées).

## 5. Original PubMed Abstract (Texte Intégral de l'Abstract)

Purpose Cryoablation (CA) is a well-established, safe and effective palliative treatment for painful bone metastases (BM). Data exploring its curative potential (i.e. radical clearing of the target lesion) is however very scarce, mostly coming from small retrospective studies. The BOne metastases REgistry for patients undergoing cryoAbLation with curatIve purpoSe (BOREALIS) aims to address this shortcoming by collecting prospective, multicentric real-world data on the outcomes of curative-intent CA for bone metastases. Methods BOREALIS is an observational cohort study designed to follow for 1 year up to 50 participants with BM ( Conclusion BOREALIS intends to provide insights into the effectiveness of curative-intent CA for BM to inform

the development of future clinical guidelines. BOREALIS is registered on ClinicalTrial.gov (NCT06480955).

---

## 6. Full-Text Sections (Europe PMC Open Access JATS XML)

### Introduction

As systemic therapies continue to extend survival, an increasing number of patients present with oligometastatic (≤ 5 lesions) or oligoprogressive bone disease, highlighting the growing need for local treatment strategies delivered with curative intent at the treated site. In this context, cryoablation (CA) has been applied selectively [1]; however, high-quality prospective data evaluating its locally curative potential remain limited. Existing studies are constrained by small sample sizes, retrospective designs, and substantial heterogeneity in patient selection, treatment protocols, and follow-up methods [2–8]. This lack of robust prospective evidence limits the ability to accurately assess the oncologic performance of CA and prevents the establishment of standardized indications for its use with a locally curative intent for bone metastases (BM).

The BOREALIS registry seeks to address this evidence gap by generating rigorous multicentre prospective data on patients undergoing CA for oligometastatic or oligoprogressive bone disease. Its primary objective is to evaluate the 1-year local non-progressive disease rate, providing a clinically meaningful measure of local tumor control.

### Study Design and Site Selection

BOREALIS is an investigator-initiated, prospective, multicentre, single-cohort observational study. It is sponsored by the Cardiovascular and Interventional Radiological Society of Europe (CIRSE) and overseen by a scientific Steering Committee composed of experts in interventional radiology, oncology, and nuclear medicine, with extensive experience in the management of BM. The study is conducted in compliance with the Declaration of Helsinki, Good Clinical Practice, and all relevant European and national regulations and has received all necessary approvals from national or local ethics committees and, where applicable, regulatory authorities.

Sites in Europe and North America with long-standing (> 10 years) experience in CA of BM and a minimum annual volume of 20 bone CA procedures during the preceding 2 years were invited to participate, with seven sites across five countries expected to contribute. Patient enrolment began in April 2025.

### Study Population

Patients with BM from oligometastatic or bone oligo-progressive disease from solid tumours scheduled to receive curative-intent CA per standard-of-care shall be recruited upon giving informed consent. Eligibility criteria are listed in Table 1. Principal Investigators (PIs) are instructed to offer participation in BOREALIS to all eligible patients and to document any refusals or reasons for non-eligibility as appropriate. Table 1Patient inclusion and exclusion criteriaInclusion Criteria1. Bone metastases from oligometastatic disease (< 5 bone/visceral metastases) or bone oligo-progressive disease (visceral or osseous multi-metastatic disease stable under systemic treatment except 1–3 progressing bone metastases) from solid tumours2. Number of target bone metastases ≤ 33. Size of target bone metastases < 5 cm (largest diameter)4. Referral to local ablative treatment by multidisciplinary tumour board (MDTB)5. Treatment by percutaneous cryoablation with curative intent (i.e., complete target bone metastases destruction) assessed as technically feasible by the IR in charge6. Procedure performed with a cryoablation system from Boston ScientificExclusion Criteria1. < 18 years old2. Incapacity or refusal to give informed consent3. Ongoing pregnancy4. Health status not compatible with the minimum meaningful follow-up period (i.e., 12 months): – Karnofsky Performance Scale < 60%, or – Bone metastases originating from rapidly evolving tumour histologies not allowing reasonable life expectancy of at least 12 months, or – Life expectancy < 12 months5. Infection of treatment site or systemic infection6. Uncorrectable coagulopathy7. Haematological disease (including multiple myeloma and plasmacytoma)8. Concomitant radiation therapy or radiation therapy within 12 weeks before the planned cryoablation procedure

Patient inclusion and exclusion criteria

4. Health status not compatible with the minimum meaningful follow-up period (i.e., 12 months):

– Karnofsky Performance Scale < 60%, or

– Bone metastases originating from rapidly evolving tumour histologies not allowing reasonable life expectancy of at least 12 months, or

**Data Collection and Quality Control**

Baseline data, obtained within 6 weeks before the planned CA procedure, includes demographics (age and sex), performance status (ECOG scale) and relevant medical history (primary origin, date of diagnosis and TNM classification, number and location of extra-osseus metastases, and details on prior or concomitant systemic bone-protective or anti-cancer treatments). Information specific to the target BM is also captured: date of diagnosis, location, radiographic feature (lytic, sclerotic, mixed, CT-occult), and history of prior local treatments. Upon completion of the planned CA session, procedural details, including information on any adjunctive biopsy or bone consolidation, are recorded for each target BM, together with the corresponding DICOM files. Follow-up data is collected at 1, 3, 6 and 12 months. Safety data is recorded from the CA procedure throughout the study, and PROMs are planned to be captured at each follow-up time point.

All data and images are collected via an electronic data capture (EDC) system (REDCap Cloud, nPhase, Inc, Version 1.7.3) through electronic case report forms (eCRFs) which incorporate pre-defined data-type constraints, range checks, and automated alerts to minimize missing or inconsistent entries and ensure data completeness and uniformity. A regular remote monitoring scheme is implemented for each participating site to verify data quality, identify potential issues related to patient enrolment and data collection, and ensure compliance with the study protocol and applicable guidelines and regulations. Source data verification may be performed as needed.

**Outcome Measures**

A list of all outcome measures is summarized in Table 2. Response to treatment will be assessed by blinded central image review (BCIR) of standard-of-care CE-MRI, CE-CT or PET-CT scans. Unless indicated otherwise, all time-to-event outcome measures will be evaluated from the time of the first CA session.
Table 2Primary and secondary outcome measures in BOREALISOutcome measureWhenReported perReported byEffectiveness - PrimaryLocal progression-free survival rate12 monthsLesionBCIREffectiveness - SecondaryComplete local tumour response12 monthsLesionBCIRDuration of response (PR or CR)Time-to-eventLesionBCIRFracture-free survivalTime-to-eventLesionInvestigator/BCIRFracture-free survival rate12 monthsLesionInvestigator/BCIRTarget bone metastasis progression-free survivalTime-to-eventPatientBCIRBone-disease progression-free survivalTime-to-eventPatientInvestigatorExtraosseous disease progression-free survivalTime-to-eventPatientInvestigatorOverall progression-free survivalTime-to-eventPatientInvestigatorOverall survivalTime-to-eventPatientInvestigatorSystemic therapy-free survivalTime-to-eventPatientInvestigatorFeasibilityTechnical feasibilityTreatment dayLesionInvestigatorFeasibility of patient managementUntil end of studyPatientInvestigatorSafetyIncidence and severity of PCs and major AEsUntil end of studyPatientInvestigatorPROMsBrief Pain Inventory – Short form1, 3, 6 and 12 monthsPatientPatientEORTC QLQ-C30 questionnaire1, 3, 6 and 12 monthsPatientPatientPCs: procedural complications, AEs: adverse events, PROMs: patient-reported outcome measures, EORTC: European Organisation for Research and Treatment of Cancer, BCIR: blinded central image review. Duration of response will be measured from the time of first documented PR or CR. All other time-to-event outcome measures will be assessed from the time of the first CA session

Primary and secondary outcome measures in BOREALIS

PCs: procedural complications, AEs: adverse events, PROMs: patient-reported outcome measures, EORTC: European Organisation for Research and Treatment of Cancer, BCIR: blinded central image review. Duration of response will be measured from the time of first documented PR or CR. All other time-to-event outcome measures will be assessed from the time of the first CA session

Evaluating local treatment response in BM remains challenging due to limitations of existing response criteria. Size-based approaches, such as RECIST [9], are inadequate for bone lesions where size is not always a reliable indicator of tumour viability and treatment response is more often reflected by structural, density, or metabolic changes. Structural or qualitative criteria, such as the MD Anderson criteria [10], lack clear quantitative thresholds, resulting in inter-observer variability and limited reproducibility. While criteria based on tumour viability or metabolic activity are central to response assessment, they often fail to capture relevant morphologic changes such as clinically significant increases in lesion volume [11–13]. Overall, these limitations highlight the need for an updated assessment framework that reflects imaging modalities currently used in clinical practice and integrates complementary morphologic and viability-related features with standardized, reproducible thresholds.

To address this need, the BOREALIS Steering Committee developed criteria for assessing treatment response of BM following CA (Table 3). These criteria combine anatomical (lesion volume) and viability/metabolic features (contrast enhancement or radiotracer uptake) with clear quantitative thresholds for the standard response categories: complete response (CR), partial response (PR), stable disease (SD), and progressive disease (PD). The BOREALIS criteria were designed to remain applicable in real-world clinical settings, allowing sufficient flexibility to accommodate variability in imaging protocols across centres and the diversity of primary tumour origins. For instance, no restriction is imposed on the type of radiotracer used in PET-CT, as its selection should be guided by tumour histology and institutional practice. Examples of BM responses to CA are presented in Figs. 1, 2 and 3 and in Supplementary Figs. 1, 2. Table 3BOREALIS Tumour response criteria for bone metastasesSizeCE-CT / CE-MRIPET-CTCRStable or decreased lesion volumeANDAbsence of contrast enhancementorAbsence of radiotracer uptakePRStable or decreased lesion volumeAND ≥ 30% decrease of viable lesion volumeor ≥ 25% reduction in SUVmaxSDAny case that does not meet the criteria for CR, PR, PDPDLesion volume increased ≥ 20%OR ≥ 20% increase of viable lesion volumeor > 25% increase in SUVmaxCR: complete response, PR: partial response, SD: stable disease, PD: progressive disease, SUV: standardised uptake value. Assessment of tumour response shall be done per target lesion, ideally using the same imaging modality. CR and PR shall be assessed taking baseline as reference, PD shall be assessed taking as reference the best response since the first CA sessionFig. 1Example of complete response (CR) assessed by CE-MRI. 55-year-old female patient with two single, biopsy-proven, bone metastases of breast cancer in the right iliac bone. A Axial contrast-enhanced T1-weighted subtracted sequence showing the two metastases (arrows) respectively sized 1.6 and 1.2 cm (volumes: 1.9 and 0.5 cm3, respectively). B Given the oligo-metastatic scenario, the patient underwent percutaneous cryoablation with local curative intent; of notice the large iceball (arrows) encompassing the two metastases. C–D Axial T2-STIR sequences obtained at 1-month follow-up showing size stability (1.6 and 1.2 cm; volumes 1.9 and 0.5 cm3, respectively) for both metastases (arrows); and E axial contrast-enhanced T1-weighted subtracted sequence obtained at the same time interval, showing a large area of necrosis (arrows) encompassing the two metastases (i.e. no contrast-enhancement of the two target metastases), thus being consistent with a complete response (CR)Fig. 2Example of partial response (PR) assessed by CE-MRI. 72-year-old male patient with a single biopsy-proven metastasis of prostate cancer in the left ischiatic bone. A Axial unenhanced T1 and B axial contrast-enhanced T1-weighted subtracted sequences showing the target metastasis (arrows) sized 1.9 cm (volume 4.5 cm3). C Given the oligo-metastatic scenario, the patient underwent percutaneous cryoablation with local «curative intent»; of notice the iceball (arrows) extending in the nearby soft-tissues; and the large hydrodissection (dotted arrows) to protect the sciatic nerve (arrowhead). D Axial T1-weighted sequence obtained at 1-month follow-up showing lesion stability in terms of size, and E axial contrast-enhanced T1-weighted subtracted sequence obtained at the same time interval showing a large area of necrosis (arrow) along with a persistent peripheral pseudo-nodular contrast-enhancement (sized 1.5 cm; volume 1.5 cm3, corresponding to a decrease of viable tumor volume larger than 30%) on the medial side of the ischiatic bone (arrow head), thus being consistent with a partial response (PR)Fig. 3Example of progressive disease (PD) assessed by CE-CT. 53-year old male patient with a biopsy-proven metastasis of kidney cancer with sarcomatoid components (ISUP 4) of the right ischiatic bone, progressing despite systemic treatment. All the other known metastases were stable, thus being consistent with an oligoprogressive bone disease. A Axial

computed tomography image showing the target metastasis (arrow), sized 2.7 cm (volume 13.8 cm3). B Given the single oligo-progression, the patient underwent percutaneous cryoablation with local curative intent; of note, the iceball (arrows) extending in the nearby soft-tissues, thus largely encompassing the target metastasis. C Axial CT image obtained at 3-month follow-up showing increased sizes (sized 9 cm; volume 265.3 cm3, corresponding to a volume increase significantly larger than 20%) of the target metastasis (arrows), thus being consistent with a progressive disease (PD)

BOREALIS Tumour response criteria for bone metastases

CR: complete response, PR: partial response, SD: stable disease, PD: progressive disease, SUV: standardised uptake value. Assessment of tumour response shall be done per target lesion, ideally using the same imaging modality. CR and PR shall be assessed taking baseline as reference, PD shall be assessed taking as reference the best response since the first CA session

Example of complete response (CR) assessed by CE-MRI. 55-year-old female patient with two single, biopsy-proven, bone metastases of breast cancer in the right iliac bone. A Axial contrast-enhanced T1-weighted subtracted sequence showing the two metastases (arrows) respectively sized 1.6 and 1.2 cm (volumes: 1.9 and 0.5 cm3, respectively). B Given the oligo-metastatic scenario, the patient underwent percutaneous cryoablation with local curative intent; of notice the large iceball (arrows) encompassing the two metastases. C–D Axial T2-STIR sequences obtained at 1-month follow-up showing size stability (1.6 and 1.2 cm; volumes 1.9 and 0.5 cm3, respectively) for both metastases (arrows); and E axial contrast-enhanced T1-weighted subtracted sequence obtained at the same time interval, showing a large area of necrosis (arrows) encompassing the two metastases (i.e. no contrast-enhancement of the two target metastases), thus being consistent with a complete response (CR)

**Statistical Analysis**

Descriptive statistics will be mainly used to report the outcome measures. The target of 42–50 participants was not determined through a formal sample size calculation but was assessed by the Steering Committee as a sufficiently large sample to provide meaningful information on the treatment's effectiveness while remaining achievable within the planned enrolment period. To understand the level of uncertainty of the collected estimate, a method involving simulation and bootstrap resampling was employed, assuming a 1-year LPFS rate of 80%. With a sample size of 50 participants, a 95% confidence interval (CI) of 68%–90% can be expected, corresponding to a 20% probability of observing a local tumour control rate below 75%.

All participants who were planned to receive cryoablation will be included in the intent-to-treat (ITT) population, regardless of whether they received the planned procedure. In contrast, the modified intent-to-treat (mITT) population will only include participants who received cryoablation.

The results of the primary outcome measure (12-month LPFS rate) and its 95% confidence interval will be derived using the Kaplan–Meier estimate, on a per-lesion basis. Lesions without documented progression will be censored at the last assessment at which their status was known, including cases where patients died without prior lesion progression. For time-to-event outcome measures, median time-to-event and 95% CI will be calculated using the Kaplan–Meier estimator. The impact of different covariates on LPFS and all other time-to-event outcome measures will be assessed using a stratified log-rank test and a univariable Cox proportional hazards regression. Covariates with p-values < 0.1 and < 20% missing data will subsequently be included in a multivariable Cox proportional hazards regression model. Variables with p-values < 0.05 will be retained after backwards elimination, or if they contribute meaningfully to the model's predictive performance. All other statistical tests will be based on a significance level of 5%, and estimates will be presented with 95% bilateral CIs. No adjustment for multiple testing will be done. For all time-to-event analyses, data will be censored at the last time point at which the patient's status was assessed.

No imputation of missing data is planned for the analyses, and missing observations will be presented as percentage of the total.

Given the absence of planned confirmatory hypothesis testing, findings should be considered exploratory. Statistical analyses will be conducted using the latest available versions of R and RStudio.

## Discussion

Cryoablation has emerged as a very promising technique for managing BM, supported by a growing body of evidence demonstrating both its safety and substantial palliative benefit, and by several notable clinical and technical advantages (analgesic effect, multi-probe technique, visualisation of the iceball on cross-sectional imaging, etc.).

Current international guidelines endorse CA as a palliative treatment option for BM [14]. However, several retrospective studies suggest that beyond pain relief, CA may also achieve meaningful and durable local tumor control, with reported rates exceeding 80% at 16–35 months [8, 15], particularly in BM sized ≤ 2 cm [4, 7]. Although these retrospective results approach those achieved by stereotactic body radiation therapy (SBRT), which yields one-year control rates around 94% [16, 17], the quality and consistency of the available CA data remain insufficient to promote guideline inclusion for curative-intent treatment.

The absence of CA from oncologic guidelines has direct implications for patient access, despite several practical and oncologic advantages. Importantly, aside from high local tumor control rates, CA enables biopsy and bone consolidation to be performed during the same procedure [18], thereby offering immediate bone stabilization when indicated. This combined approach may significantly reduce the risk of secondary pathological fractures, which were reported in only 1.8% of cases (with an overall 2.3% major complication rate) in a large series of bone tumors undergoing CA [19]. These rates contrast sharply with the fracture risks associated with SBRT: up to 18% for vertebral sites [20] and up to 10% for extra-spinal ones [21, 22]. Moreover, while SBRT remains a highly effective option, its further limitations include decreased effectiveness in certain radio-resistant histologies, restrictions on repeat treatment due to cumulative dose constraints, and the logistical burden of multi-fraction regimens [23]. By contrast, CA is repeatable and compatible with concurrent systemic or local therapies, including radiation [24].

Despite these advantages, prospective real-world data evaluating the curative potential of CA remain scarce; and existing studies are limited by retrospective design, small cohorts, heterogeneous imaging and follow-up protocols, and lack of standardized response criteria. As a result, clinicians lack definitive evidence to guide patient selection, anticipate oncologic performance, or establish CA as a standard local treatment option for oligometastatic or oligoprogressive bone disease.

BOREALIS is designed to address these gaps. By focusing on patients undergoing CA with curative intent, BOREALIS aims to provide estimates of 12-month local progression-free survival rates, assessed using newly developed, ablation-specific tumor response criteria that integrate anatomical and functional imaging metrics.

## Conclusions

BOREALIS is designed to generate high-quality, prospective evidence to better define the role of CA in the contemporary management of BM requiring local control. By integrating clinical, procedural, radiological, and patient-reported outcomes, the study aims to inform future guideline development and clarify the position of CA among available therapeutic options for patients with oligometastatic or oligoprogressive bone disease.