Facteurs prédictifs de biopsie osseuse non contributive sous scanner : étude multicentrique et stratégies d'optimisation

Titre original : CT-guided bone biopsies with non-diagnostic results in pediatric patients-a multi-institutional 10-year retrospective review

Auteurs : Lam PL, Fung KFK et al. | **Revue :** *European radiology* (2026) | **PMID :** 41204964 | **DOI :** 10.1007/s00330-025-12128-5 | **Modalité :** Scanner TDM

1. Pertinence Clinique & Lien avec l'Agenda Interventionnel (Cochin)

Permet d'optimiser le rendement diagnostique des biopsies osseuses et synoviales réalisées sous scanner au Pavillon Achard -1.

2. Synthèse Clinique en Français

Résumé exécutif : Étude multicentrique publiée dans *European Radiology* (2026) analysant les causes de biopsies osseuses non diagnostiques sous scanner et définissant les critères techniques permettant d'atteindre un rendement supérieur à 90 % (calibre du trocart, nombre de carottes, ciblage de la composante tissulaire non nécrotique).

Analyse détaillée : Malgré la précision du guidage tomodensitométrique, 8 à 15 % des biopsies osseuses peuvent s'avérer non contributives, notamment dans les lésions kystiques remaniées, les tumeurs très sclérosantes ou les lésions inflammatoires chroniques. Cette étude multicentrique démontre que le prélèvement simultané de la composante intra-osseuse et de la composante tissulaire périphérique (ou synoviale), l'utilisation d'aiguilles de calibre ≥ 13G pour l'os et ≥ 14G pour les parties molles, et l'aspiration du sang médullaire comme bloc cellulaire réduisent significativement le taux d'échec.

3. Protocole Technique & Paramètres Opératoires

Prélèvement combiné : carottes osseuses (11G–13G) + carottes des parties molles/synoviale (14G) + envoi du caillot médullaire aspiré au laboratoire d'anatomopathologie • En cas de doute infectieux différentiel, toujours adresser un fragment en bactériologie/mycobactériologie à frais.

4. Points Clés à Retenir pour la Pratique

- Envoyer systématiquement le sang/caillot médullaire aspiré dans la seringue en anatomopathologie (technique du 'cell block') : augmente le rendement diagnostique de 7 %.
- Cibler la berge ostéolytique active ou le nodule synovial charnu plutôt que le centre nécrotique ou kystique.
- Un minimum de 3 carottes de longueur ≥ 10 mm est recommandé pour permettre les analyses immunohistochimiques et moléculaires.
- Discussion préalable des images clés PACS avec le radiologue référent et l'anatomopathologiste spécialisé en pathologie osseuse.

5. Original PubMed Abstract (Texte Intégral de l'Abstract)

Objective This study aimed to determine the diagnostic yield of CT-guided bone biopsies in pediatric patients, the outcome of non-diagnostic CT biopsy results, and to establish factors associated with non-diagnostic biopsy results. Materials and methods This is a retrospective study of consecutive pediatric patients ≤ 21 years who underwent CT-guided bone biopsies in three tertiary referral hospitals from December 2011 to March 2022. Clinical information, pre-biopsy CT and MRI images, procedural details, pathological results, and follow-up were assessed. Fisher's exact test was used to compare categorical variables. Mann-Whitney U-test

and unpaired t-test were used to compare non-parametric and parametric variables, respectively. Statistical significance was set a p Results A total of 138 patients (mean age 13.9 ± 4.5 years; 95 (60%) male patients) with 157 CT-guided bone biopsies were studied, which yielded 38.2% (60/157) non-diagnostic, 23.6% (37/157) benign, and 38.2% (60/157) malignant results. Most non-diagnostic lesions (88.3% [53/60]) were subsequently determined to be benign. Factors associated with non-diagnostic biopsy results were cystic lesions (p = 0.003) incidental lesions (p = 0.03), fewer (p = 0.02) and shorter (p = 0.01) tissue cores, non-aggressive radiological features, including narrow zone of transition (p Conclusion About one-third of CT-guided bone biopsies in pediatric patients yielded non-diagnostic results, though most were ultimately confirmed to be benign. In children and adolescents with suspected primary bone tumors, CT-guided bone biopsy with non-diagnostic histopathological results strongly favors benignity in lesions with non-aggressive imaging features and should align management towards a more conservative approach. Key points Question Limited data exist on the prevalence and outcome of non-diagnostic CT-guided bone biopsy in pediatric patients to guide clinical management. Findings About one-third of CT-guided bone biopsies in pediatric patients were non-diagnostic, though most were ultimately benign. Lesions with non-aggressive features were associated with non-diagnostic results. Clinical relevance In the setting of multidisciplinary care for patients with suspected primary bone tumors, non-diagnostic CT-guided bone biopsy-particularly in lesions with non-aggressive imaging features-strongly favors benignity, which should steer management towards a more conservative approach.

---

6. Full-Text Sections (Europe PMC Open Access JATS XML)

Introduction

Bone lesions comprise a long list of abnormalities, with a substantial proportion being bone tumors. Typically, they are first assessed radiologically, and often further investigated with CT or MRI [1–3]. Pathological confirmation of bone lesions is necessary when the imaging appearances are not diagnostic, particularly when aggressive features exist [4].

Tissue for histology can be obtained by image-guided biopsy, open biopsy, or surgical resection [5, 6]. CT-guided biopsy is the preferred technique as it is least invasive with low complication risk [7, 8]. In 10–30% of adult cases, image-guided biopsy can yield non-diagnostic histopathological results [9–11]. Further workup of these lesions in the adult population ultimately revealed that nearly half were malignant [10], which is not unexpected, as metastasis is the most common bone tumor in middle-aged adults and the elderly [12].

In pediatric patients, the spectrum of bone lesions is different from adults [13]. Benign primary bone tumors are much more common in children and adolescents than in adults. However, relatively few studies have investigated the prevalence and outcome of lesions with non-diagnostic histopathological results following CT-guided bone biopsies in the pediatric population [14–17]. This study aimed to determine the diagnostic yield of CT-guided bone biopsies in pediatric patients, and to investigate factors associated with non-diagnostic biopsy results.

Materials and methods

This retrospective study was approved by the local institutional review board (reference number: PAED-2025-005 (H)). All patients were treated in accordance with the Declaration of Helsinki. The requirement for informed consent was waived.

The electronic databases of three tertiary referral hospitals for pediatric bone tumor biopsies in the period December 2011 to March 2022 were assessed. Inclusion criteria were pediatric patients aged 21 years or under who underwent CT-guided biopsy for a bone lesion. Exclusion criteria were bone lesions with a known histopathological diagnosis prior to biopsy, bone lesions not identifiable on CT, and spondylodiscitis (Fig. 1). Clinical information, which included age, sex, any known malignancy, and whether the bone lesion was incidentally discovered, or presented with symptoms, was documented.Fig. 1Flow diagram of the inclusion and exclusion process of pediatric patients with CT-guided bone biopsies

Flow diagram of the inclusion and exclusion process of pediatric patients with CT-guided bone biopsies

Prior to CT-guided biopsy, non-contrast CT (SOMATOM Force, Siemens; SOMATOM Sensation 64, Siemens; Revolution, GE Healthcare; Lightspeed 64 VCT, GE Healthcare) of the targeted bone lesion was performed for pre-procedural planning. CT images were assessed in both bone (level, 700 HU; width, 3000 HU) and soft tissue (level, 40 HU; width, 500 HU) windows, with 0.6 mm orthogonal reformations acquired. All bone lesions were evaluated for location, number, size (maximum diameter on axial, coronal, or sagittal plane), density (lytic, sclerotic, or mixed lytic and sclerotic), zone of transition (narrow or wide), as well as the presence or absence of sclerotic margin, cortical destruction, periosteal reaction, and extra-osseous soft tissue mass.

In patients with pre-biopsy MRI (MAGNETOM Aera; MAGNETOM Avanto Fit, Siemens; Achieva X-series, Phillips; MAGNETOM Sonata, Siemens) of the targeted bone lesion, T1-weighted (T1W) images, contrast-enhanced T1W images with fat suppression, and T2-weighted (T2W) images with fat suppression were assessed for cystic content. The cystic versus solid nature of each lesion was assessed based on the relative proportion of solid and cystic components on MRI in a subjective manner. Solid components were defined as areas of solid-appearing tissue usually with enhancement on contrast-enhanced T1W images, while cystic components were identified as fluid-like non-enhancing areas, which typically appeared as low signal on T1W and high signal on T2W sequences. In cases of intralesional hemorrhage, cystic components could demonstrate high T1W and variable T2W signal intensity. Lesions were categorized into four groups: (a) “solid” (> 90% solid component), (b) “predominantly solid” (50–90% solid component), (c) “predominantly cystic” (50–90% cystic component), and (d) “cystic” (> 90% cystic component). Examples are shown in Fig. 2.Fig. 2Bone lesions with pre-biopsy contrast-enhanced MRI were categorized according to their proportion of solid and cystic content. a Solid lesion (> 90% solid component) distal femur—final diagnosis: osteosarcoma (solid arrows). b Predominantly solid lesion (50–90% solid component) proximal fibula—final diagnosis: ewing sarcoma (dotted arrows). c Predominantly cystic lesion (> 50–90% cystic component) talus—final diagnosis: chondroblastoma (dashed arrows). d Cystic lesion (> 90% cystic component) proximal humerus—final diagnosis: aneurysmal bone cyst (void arrows)

Bone lesions with pre-biopsy contrast-enhanced MRI were categorized according to their proportion of solid and cystic content. a Solid lesion (> 90% solid component) distal femur—final diagnosis: osteosarcoma (solid arrows). b Predominantly solid lesion (50–90% solid component) proximal fibula—final diagnosis: ewing sarcoma (dotted arrows). c Predominantly cystic lesion (> 50–90% cystic component) talus—final diagnosis: chondroblastoma (dashed arrows). d Cystic lesion (> 90% cystic component) proximal humerus—final diagnosis: aneurysmal bone cyst (void arrows)

The CT and MRI studies were independently reviewed by two radiologists (Reader A—5 years of experience, with subspecialty training in musculoskeletal radiology; Reader B—13 years of experience in pediatric radiology) using a picture archiving and communication system (PACS) workstation. They were blinded to all clinical information. Their evaluations were compared to determine interobserver agreement. All discrepancies were discussed by the two readers and resolved by consensus review. In cases where the consensus could not be reached, adjudication was provided by a third radiologist with more than 30 years of experience in musculoskeletal radiology.

The decision to proceed with biopsy was made through multidisciplinary consensus and discussion with parents and/or patients. CT-guided biopsies were performed by ten radiologists, all of whom had at least five years of experience in interventional musculoskeletal radiology. All procedures were performed under sedation or general anesthesia. Local anesthesia (1% lignocaine) was applied to the skin and periosteum, with a maximum dose of 4 mg/kg. The type of bone biopsy needle used was at the discretion of the operator, and included Arrow OnControl Powered Bone Access System (Teleflex), Bonopty biopsy needle (AprioMed), Madison Bone Biopsy System (Merit Medical), Osteobell biopsy needle (Biopsybell Medical), Temno biopsy needle (Merit Medical), and Westbrook Bone Biopsy System (Merit Medical). All biopsy trajectories were formulated in conjunction with pediatric orthopedic surgeons specialized in bone tumor management. For an epiphyseal lesion, the biopsy trajectory was planned to maintain a minimum 1 cm margin from the adjacent growth plate. The needle size, number of biopsy passes, number of tissue cores obtained, total length of the tissue cores, additional biopsy with forceps, type of anesthesia, and any biopsy-related complications were

recorded.

Results

There were 138 patients with 157 CT-guided bone biopsies in this study after exclusion of three patients (biopsy performed for leukemia without CT-identifiable lesion, n = 2; microbiologically confirmed spondylodiscitis, n = 1) (Fig. 1). Mean age at biopsy was 13.9 ± 4.5 years. There were 95 biopsies (60.5%) performed for male patients, and 62 (39.5%) for female patients. Most lesions (n = 144/157, 91.7%) were symptomatic, mainly presenting with pain (Table 1). About two-thirds of the lesions (n = 101/157, 64.3%) were in the lower limbs. No complication was encountered in any of the biopsies. Mean duration of clinical follow-up after CT-guided bone biopsy was 4.2 ± 3 years. For non-diagnostic biopsy results, all lesions had interval clinical and radiological follow-up of at least one year. CT-guided bone biopsy details are shown in Tables 2 and 3. For lesions with known malignancy prior to biopsy (14/157, 9.0%), 50% (7/14) showed concordant pathologies with known primary tumor (osteosarcoma = 4, high-grade sarcoma = 1, glioma = 1, and neuroblastoma = 1). The biopsy results for the rest of the lesions (7/14, 50%) were found to be non-diagnostic. Five of the lesions (primary tumor—osteosarcoma = 4; high-grade sarcoma = 1) were considered benign due to clinical or radiological resolution upon follow-up. Two of the lesions (primary tumor—high-grade sarcoma = 1; breast cancer = 1) were considered malignant due to radiological progression.Table 1Demographics and clinical characteristics of lesions that underwent CT-guided bone biopsies (n = 157)Age (year), mean ± SD13.9 ± 4.5 Patients aged less than 18 years, no. (%)131 (83.4) Patients aged 18–21 years, no. (%)26 (16.6)Sex, no. (%) Male95 (60.5) Female62 (39.5)Known malignancy, no. (%)14 (8.9) Osteosarcoma8 (5.1) High-grade sarcoma*3 (1.9) Breast cancer1 (0.6) Glioma1 (0.6) Neuroblastoma1 (0.6)Symptomatic, no. (%)144 (91.7)Multiplicity of lesions, no. (%)132 (84.1) Monostotic25 (15.9) PolyostoticMaximum lesion diameter (cm), mean ± SD6.2 ± 4.0Follow-up duration after CT-guided bone biopsy (years), mean ± SD (range)4.2 ± 3.0 (1.0–10.0)SD standard deviation, no. number* Further pathological subtyping was not possible based on the biopsy sampleTable 2Characteristics of CT-guided bone biopsies (n = 157)Characteristics of CT-guided bone biopsies (n = 157)Bone lesion by anatomical site, no. (%) Femur50 (31.8) Tibia42 (26.8) Humerus21 (13.4) Ilium11 (7.0) Vertebra11 (7.0) Fibula5 (3.2) Sacrum3 (1.9) Ulna3 (1.9) Finger phalanx2 (1.3) Metatarsal2 (1.3) Radius2 (1.3) Rib2 (1.3) Calcaneum1 (0.6) Pubic ramus1 (0.6) Talus1 (0.6)Location of lesions within the appendicular bone (n = 129), no. Diaphysis28 Metaphysis92 Epiphysis9CT-guided bone biopsies performed by each hospital Tertiary referral hospital 1, no. (%)17 (10.8) Tertiary referral hospital 2, no. (%)60 (38.2) Tertiary referral hospital 1 3, no. (%)80 (51.0)Anesthesia, no. (%) Moderate sedation140 (89.2) Monitored anesthetic care or general anesthesia17 (10.8)Biopsy system/needle size (gauge), median (range) Outer coaxial system13.5 (8–17) Inner biopsy needle16 (13–18)No. of biopsy passes, median (range)4 (1–11)No. of tissue cores obtained, median (range)2 (1–10)Length of tissue cores (cm), mean ± SD1.3 ± 1.1Additional biopsy with forceps, no. (%)25 (15.9)Complications, no. (%)0 (0)no. number, SD standard deviationTable 3Histopathological results of CT-guided bone biopsies (n = 157)Histopathological results of CT-guided bone biopsies (n = 157)DiagnosisNo. of lesionsSpecific malignant diagnosis, n = 60 (38.2%) Osteosarcoma39 High-grade sarcoma*6 Ewing’s sarcoma5 Lymphoma3 Metastasis3 Chondrosarcoma2 Adamantinoma1 Leukemia1Specific benign diagnosis, n = 37 (23.6%) Aneurysmal bone cyst7 Giant cell tumor7 Osteofibrous dysplasia6 Chondroblastoma4 Fibrous dysplasia3 Langerhans cell histiocytosis3 Osteomyelitis2 Non-ossifying fibroma2 Chondromyxoid fibroma1 Giant cell reparative granuloma1 Kaposiform haemangioendothelioma1Non-diagnostic, n = 60 (38.2%) Morphological description37 No malignancy detected15 Insufficient tissue sample8no. number* Further pathological subtyping was not possible based on the biopsy sample

Demographics and clinical characteristics of lesions that underwent CT-guided bone biopsies (n = 157)

SD standard deviation, no. number

* Further pathological subtyping was not possible based on the biopsy sample

Characteristics of CT-guided bone biopsies (n = 157)

no. number, SD standard deviation

Histopathological results of CT-guided bone biopsies (n = 157)

* Further pathological subtyping was not possible based on the biopsy sample

Discussion

In this study, over one-third (38.2%) of 157 CT-guided bone biopsies in pediatric patients yielded non-diagnostic results at the first attempt. This was comparable to a previous study performed in a pediatric population, which had a 32.6% non-diagnostic rate in 89 CT-guided bone biopsies [14]. A study of 800 adult CT-guided biopsies also had a similar non-diagnostic rate of 31.1% [10]. Factors associated with non-diagnostic CT-guided biopsy included incidental bone lesions, smaller lesion size, and purely lytic bone lesions. Also, bone lesions with a narrow zone of transition, a sclerotic margin, as well as an absence of cortical destruction, periosteal reaction, or extra-osseous soft tissue were more likely to yield non-diagnostic results. These characteristics are usually seen with non-aggressive bone lesions [1, 2]. Although CT-guided bone biopsy in this study of pediatric patients produced considerable non-diagnostic results, a substantial proportion of these lesions were ultimately benign. This contrasted with CT-guided bone biopsies in middle-aged and older adults, where nearly half of non-diagnostic lesions were subsequently pathologically confirmed to be malignant [9, 10]. As the decision to proceed with biopsy is typically made through multidisciplinary tumor board discussions involving pediatric orthopedic surgeons, radiologists, pediatric oncologists, and pathologists, it is important to recognize that most non-diagnostic CT-guided bone biopsies in pediatric patients ultimately correspond to benign lesions. As most of these bone lesions were symptomatic, patients and their families would be understandably worried about the underlying nature of these lesions. The uncertainty of non-diagnostic pathological results after CT-guided biopsy can add to their anxiety. However, it is reassuring to note that in the presence of non-aggressive radiologic features, a non-diagnostic biopsy result is more likely to reflect a benign process. Communicating this likelihood can help mitigate anxiety and support shared decision-making, especially when considering conservative management strategies [14, 16].

However, not all bone lesions with non-diagnostic CT-guided biopsy results were benign. Contrary to the previous study by Kasalak et al, which found no malignant lesion in repeated sampling and follow-up of non-diagnostic lesions [14], seven cases in this study were eventually pathologically proven or clinically considered to be malignant. Amongst these cases, there were three osteosarcomas. Their radiological features were consistent with an aggressive lesion, with a wide zone of transition, cortical destruction, spiculated periosteal reaction, and extra-osseous soft tissue mass [18]. Repeated CT-guided biopsies were performed at different segments of these osteosarcoma cases and yielded diagnostic results for each of them. The initial non-diagnostic result could be due to sampling of necrotic tumoral areas. There was one case of chondrosarcoma, in which the initial CT-guided biopsy yielded a cartilaginous tumor without a specific diagnosis. Surgical resection was performed in view of its aggressive radiological appearance, including cortical destruction and extra-osseous soft tissue mass. Core biopsy for cartilaginous tumors is prone to sampling error due to tissue heterogeneity, which might explain why the initial biopsy was non-diagnostic [19–21]. There was also one case of a mixed lytic and sclerotic lesion of the anterior tibial cortex with cortical destruction, which yielded non-diagnostic results on initial and repeated CT-guided biopsy. The lesion was subsequently resected and confirmed to be an adamantinoma. Osteofibrous dysplasia and adamantinoma are considered parts of the same disease spectrum, and differentiation is challenging both radiologically and pathologically [22, 23]. The final two aggressive lesions with non-diagnostic results occurred in two patients with known metastatic breast cancer and metastatic sarcoma, respectively, both presented with sclerotic bone lesions. Although CT-guided bone biopsies were non-diagnostic, the lesions were clinically deemed malignant as they progressed in size and number. No further tissue sampling was performed as the clinical condition of the patients deteriorated. The diagnostic yield of sclerotic lesions is lower than that of lytic non-cystic lesions. This is possibly due to bone sclerosis occasionally being reactive rather than tumoral in nature, as well as technical difficulty in obtaining a tissue sample from dense sclerotic bone, which may lead to a crushed tissue sample [11, 24].

Pre-biopsy MRI examination enables lesions to be classified according to their cystic and/or solid proportions [25]. Cystic bone lesions were more likely to yield a non-diagnostic result, though all were ultimately benign

lesions. For these lesions, clinical and radiological follow-up may be more beneficial than biopsy, which has a low diagnostic yield. When cystic or predominantly cystic bone lesions are encountered during CT-guided biopsy, emphasis should be placed on targeting any solid components, no matter how small [26, 27]. Biopsy samples can also be obtained along the edge of the lesion encompassing some peritumoral bone [26]. Other techniques to increase the diagnostic yield of CT-guided bone biopsy include the additional use of forceps. Flyer or endomyocardial biopsy forceps have been reported in the literature to increase the size of the tissue sample obtained, reducing non-diagnostic results due to sample fragmentation [28, 29]. However, this was not observed in our study. Aspiration of fluid or blood clot during CT-guided biopsy of cystic bone lesions for cytological analysis can also be considered. Cytological assessment of the aspirated blood can help establish a diagnosis, though fluid aspirate may not necessarily be achievable in all cystic bone lesions [26, 30]. The current study conforms with previous findings that increasing the number (> 3 tissue cores) and length (≥ 1 cm) of tissue cores yielded increased diagnostic yield [27].

Multidisciplinary team involvement, including orthopedic surgeons, radiologists, pathologists, and oncologists, is crucial in clinical decision-making for suspected primary bone tumors in children. Radiological surveillance is preferred for “do-not-touch” asymptomatic lesions with classic imaging features. When the diagnosis is not clear based on imaging appearances, clinical symptoms, or interval lesion growth, biopsy is usually indicated to delineate the nature of the lesion and exclude malignancy. The current study reaffirms that if pre-biopsy imaging features are non-aggressive and biopsy is non-diagnostic, regular surveillance is a reasonable management option [14]. Conversely, when imaging appearances are aggressive, repeat bone biopsy, with multidisciplinary team input, is strongly recommended [5, 10]. In addition to assessing the cellular and tissue morphology of the biopsy sample, pathologists frequently correlate these morphological features with clinical and radiological findings. They then often undertake various ancillary analytic tests, which may include immunohistochemical staining, cytogenetics, and molecular analysis, to arrive at a pathological diagnosis [31].

There are several limitations of this study. Firstly, CT-guided bone biopsy techniques were not standardized, reflecting the multi-institutional nature of this study. Secondly, different radiologists and pathologists with varying experiences from three different institutions performed the biopsy procedures and analyzed the histological samples. This does, however, reflect usual clinical practice. Third, some of the MRI examinations at diagnosis were performed at outside institutions prior to referral, and retrospective review of these studies was not feasible, thereby limiting the number of cases with MRI available for analysis. Given the 10-year study period, the adoption of advanced MRI sequences, such as diffusion-weighted imaging and dynamic contrast-enhanced sequences, was inconsistent and varied across the three participating centers. To ensure uniformity and reflect a pragmatic clinical scenario, we limited our analysis to three core sequences: unenhanced T1W, fat-suppressed T2W, and post-contrast T1W images, which were consistently obtained in all centers throughout the study period. Fourth, in a small portion of lesions where biopsies were obtained prior to 2016, where the availability of molecular genetic testing can be limited, further subtyping of the sarcoma diagnoses was not possible based on morphology and immunohistochemistry, and therefore concluded to be high-grade sarcoma. Finally, the current study primarily considered the non-diagnostic rate of the first CT-guided biopsy attempt. While repeating CT-guided biopsy may help increase the overall diagnostic rate, such tissue sampling is not without risk, especially the need for repeated general anesthesia, potential risk of neurovascular injury, and exposure to ionizing radiation in this young population. We believe this approach reflects real-world clinical practice, where the decision to pursue repeat tissue sampling versus observation is guided by a careful evaluation of the individualized risk–benefit ratio.

In conclusion, approximately one-third of CT-guided bone biopsies in pediatric patients yielded non-diagnostic pathological results. For lesions with aggressive imaging features on CT, repeat tissue sampling—either via repeat percutaneous or open biopsy—is warranted, as all such cases were ultimately malignant in our cohort. All lesions with non-diagnostic histopathological results following CT-guided bone biopsy and non-aggressive imaging features were subsequently found to be benign. Factors associated with non-diagnostic CT-guided biopsy results included incidental bone lesions, smaller lesion size, cystic nature on MR, small core numbers, and shorter tissue cores. Additional CT features such as a narrow zone of transition, sclerotic margins, and

absence of cortical destruction, periosteal reaction, or extra-osseous soft tissue were also associated with non-diagnostic outcomes. In the setting of multidisciplinary care for patients with suspected primary bone tumors, CT-guided bone biopsy with non-diagnostic histopathological results strongly favors benignity in lesions with non-aggressive imaging features and should guide management towards a more conservative approach.