

1 Exercises

Derivative Suggested Exercises I

I) Understanding the Definition

- i) Explain intuitively why the limit formula for the derivative in equation (??) is correct? (i.e. gives you the best linear approximation to the function at the point).
- ii) Give a geometric interpretation of the derivative of a function. (*Hint: Draw a graph of a function, pick a point and draw something that helps clarify what the derivative is. You should think about what the Jacobian is: can you identify it's domain and target in the picture?.*).

II) Computation Warmup

- i) Compute the derivative of $f(x) = kx$ for k any constant at some point $x_0 \in \mathbb{R}$
- ii) Compute the derivative of $f(x) = ax^2$ for a any constant, at some point $x_0 \in \mathbb{R}$

III) **Bonus:** Suppose $f : \mathbb{R} \rightarrow \mathbb{R}^n$ with $f(t) = (f_1(t), \dots, f_n(t))$. How should the Jacobian of f look like?

Derivative Suggested Exercises II

I) Computational Practice: Compute the following derivatives:

$$\begin{array}{cc} \frac{d}{dt} \sin(3t) & \frac{d}{dt} (\cos(t))^2 \\ \frac{d}{dt} \cos(t) \sin(t) & \frac{d}{dt} \exp(\cos(t)) \\ \frac{d}{dt} \frac{1}{t^n} & \frac{d}{dt} \frac{1}{\sin(t)} \end{array}$$

II) Understanding Division:

- i) Let f be a differentiable function and $f \neq 0$. Compute the following derivative:

$$\frac{d}{dt} \frac{1}{f(t)}$$

- ii) Use the result above to verify that:

$$\frac{d}{dt} \frac{f(t)}{g(t)} = \frac{f'(t)g(t) - f(t)g'(t)}{(g(t))^2}$$

III) Inverse of Differentiation

Find a function such f that the following are its derivative:

$$\frac{d}{dt} f = t^2 \quad \frac{d}{dt} f = \sin(t) \cos(t)$$

Suggested Integral Exercises

I) Compute the following integrals

$$\begin{aligned} \int_0^a dx \quad \int_0^a 3x^2 + 7 \\ \int_0^a e^{kx} dx \quad \int_0^a xe^{x^2} \\ \int_0^a x \sin(x) \quad \int_0^a \sin(x) \cos(x) \\ \int_0^a xe^x \quad \int_0^a x^2 e^x \end{aligned}$$

II) Recall that Newton's equations of motion are given by:

$$F = ma = m\ddot{x}$$

i) Consider a particle restricted to 1 dimension. Suppose we attach a string to it. Then in some coordinate system, the force acting on the particle at position x is $F = -kx$.

We define the potential energy to be:

$$U(p) = - \int_0^p F(x) dx$$

Compute $U(x)$. Then show that the total energy:

$$E(x(t)) = \frac{1}{2}m(\dot{x})^2 + U(x(t))$$

Is constant with time. Show that if E is conserved for any F that does not depend on t .

Partial Derivative Suggested Exercises

I) Let $g : \mathbb{R} \rightarrow \mathbb{R}^m$ and $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be both differentiable. Convince yourself that:

$$\left. \frac{\partial}{\partial t} f(g(t)) \right|_{t_0} = \sum_{i=1}^m \left. \frac{\partial f}{\partial x_i} \right|_{g(t_0)} \cdot \left. \frac{\partial g_i}{\partial t} \right|_{t_0}$$

II) Compute the following derivatives:

$$\begin{aligned} \frac{\partial \sin(xy)y^2}{\partial x} \quad \frac{\partial e^{x^2 \sin(y)}}{\partial y} \\ \frac{\partial}{\partial x} \frac{\partial}{\partial y} e^{x^2 \sin(y)} \quad \frac{\partial}{\partial y} \frac{\partial}{\partial x} e^{x^2 \sin(y)} \end{aligned}$$

III) Consider the function:

$$T(t) = \frac{1}{2} (\dot{x}_1^2 + \dot{x}_2^2)$$

i) Suppose you are given $r(t)$ and $\varphi(t)$ such that:

$$x = (x_1, x_2) = (r \cos \varphi, r \sin \varphi)$$

Express $T(t)$ in terms of $r(t)$ and $\varphi(t)$

IV) Recall that acceleration is:

$$a = \ddot{x} = (\ddot{x}_1, \ddot{x}_2)$$

Suppose that the particle's motion is restricted to a circle of radius r . It's path $x(t)$ is then fully determined by some function $\varphi(t)$ via:

$$x(t) = r(\cos \varphi(t), \sin \varphi(t))$$

Show that the acceleration, must be proportional to $x(t)$ (i.e. pointing along the same direction as the radius vector of x)

Suggested Exercises Functional Calculus

I) Let

$$F[f] = \int_0^1 (f(x))^2 dx$$

- i) Compute $DF_f[h]$ for arbitrary f and h
- ii) Suppose that F is restricted only to functions f such that $f(a) = f(b)$. Suppose that some function g extremizes F . Find the condition that g must satisfy.

II) Let

$$F[x_1(t), x_2(t)] = \int_0^{t_1} \sqrt{\dot{x}_1^2 + \dot{x}_2^2} dt$$

Where $x_1(0) = x_2(0) = 0$ and $x_1(t_1) = x_2(t_1) = 1$

What does this functional compute?

Find $\tilde{x}(t)$ such that $DF[\tilde{x}] = 0$. Does this answer make sense?