

WEEK
1

Foundations + Northern Hardwood Anchors

Instruction | retrieval | multi-organ practice | field-ready reference

Public learner edition

This packet contains the teaching guide, cumulative reference, and formative practice for the week. Secure graded assessments remain in the online course so unfamiliar examination specimens and answer material are not published.

Student / participant

Date started

Planned pace / study notes

Field standard

Identify the narrowest rank the visible evidence can defend. A common name OR a scientific name satisfies an ordinary identity prompt unless the activity explicitly tests nomenclature. State uncertainty instead of converting weak evidence into a confident species claim.

Use this packet alongside the online course at cochetopa.co/course/. Work the instruction first, complete retrieval without looking back, and use the secure online assessment when ready for grading.

Week 1 overview

Learning objectives

- 1. Apply dendrological vocabulary and the maximum-defensible-rank decision rule.
- 2. Identify the northern-hardwood anchor taxa from unfamiliar foliage, bark, reproductive, dormant, and whole-tree evidence where defensible.
- 3. State a nearest plausible alternative, visible separator, confidence code, and resolving view.
- 4. Retrieve canonical common and scientific names and introductory field silvics for every new taxon.
- 5. Separate American beech host identity from clean or normal bark condition.

Species introduced this week

Common name	Scientific name	Family	Tier
Sugar maple	Acer saccharum	Sapindaceae; Aceraceae in older texts	A
Red maple	Acer rubrum	Sapindaceae	A
American beech	Fagus grandifolia	Fagaceae	A
Yellow birch	Betula alleghaniensis	Betulaceae	A
Paper birch	Betula papyrifera	Betulaceae	A
Eastern hemlock	Tsuga canadensis	Pinaceae	A
American basswood	Tilia americana	Malvaceae; Tiliaceae in older texts	A
Black cherry	Prunus serotina	Rosaceae	A
Eastern hophornbeam	Ostrya virginiana	Betulaceae	A

Cumulative review

9 taxa remain eligible. Earlier species do not disappear from retrieval merely because the weekly theme changes.

Confusion sets

Yellow birch / black cherry bark: Golden horizontal curls and wintergreen twig versus burnt-chip plates, petiole glands, and almond twig odor
Request: Young and old bark; scratched twig; leaf petiole and underside; fruit

Assessment schedule

Assessment	Type	Items	Time
W01-PRACTICAL	weekly identification practical	18	35 min

Five-session field workflow

Session A - Six northern-hardwood anchors

diagnostic teaching examples plus field-decision modeling. Modalities: foliage, bark, whole tree context.

☐ Session A completed (enter X)

Session B - Anchor retrieval

closed-response batches of five to ten unfamiliar practice items. Modalities: foliage, bark, whole tree context.

☐ Session B completed (enter X)

Session C - Birch, cherry, and hophornbeam contrasts

direct comparison and isolated unfamiliar specimens. Modalities: foliage, bark, reproductive, dormant.

☐ Session C completed (enter X)

Session D - Anchor multi-organ laboratory

single-organ and partial-evidence practice stations. Modalities: foliage, bark, reproductive, dormant, whole tree context.

☐ Session D completed (enter X)

Session E - Week 1 cumulative retrieval

adaptive retrieval and remediation insertion. Modalities: foliage, bark, reproductive, dormant, whole tree context.

☐ Session E completed (enter X)

Evidence-to-decision protocol

Step	Field action
1. Inventory	Name only what is visible: organ, arrangement, attachment, surface, scale, maturity, damage, and context.
2. Generate	Name the leading identity and the nearest plausible alternative before seeking confirmation.
3. Separate	Cite the visible character that changes the decision; do not substitute habitat or color alone for structure.
4. Calibrate	Use species, genus, or insufficient evidence according to the photograph, not according to what you hope it shows.
5. Request	Ask for the single additional view with the highest expected diagnostic value.

Part I - Species instruction atlas

These profiles are labeled teaching material. Photographs come only from the teaching pool; no exact image in the formal assessment section appears here.

Sugar maple - *Acer saccharum*

New this week | Family: Sapindaceae; Aceraceae in older texts | Tier A | Priority 1 - core

Widespread from MN and the Great Lakes through southern ON/QC, NY, New England, NB, and the Allegheny region

Major shade-tolerant northern-hardwood canopy dominant; valuable timber and sap tree



Bark

(c) Loïc Mathieu, some rights reserved...
INAT_PHOTO_611772046 | cc-by



Dormant

(c) daksu, some rights reserved (CC BY)
INAT_PHOTO_612188654 | cc-by



Foliage

no rights reserved
INAT_PHOTO_669709341 | cc0

Evidence	Diagnostic interpretation
Foliage	Opposite, simple blades are palmately veined and usually five-lobed, with pointed lobes, broad rounded U-shaped sinuses, and relatively few coarse teeth. A fresh broken petiole has clear rather than milky sap; arrangement and margin pattern matter more than fall color.
Bark age series	Seedlings and saplings have smooth light-gray bark; pole trees develop long shallow furrows; mature and old boles form irregular vertical plates whose edges lift outward. Old trees can be dark and deeply fissured, so the classic plated surface is an age-series character rather than a universal texture.
Dormant	Opposite brown twigs carry a narrow, sharply pointed terminal bud with many tightly overlapping brown scales and smaller paired lateral buds. Leaf scars are opposite and usually V-shaped; the narrow conical terminal bud is a high-value separator from red and Norway maples.
Fruit / cone / seed	Small yellow-green apetalous flowers hang on long stalks before or during leaf expansion. Paired samaras mature in autumn and commonly meet at an acute angle; timing and wing angle support the identification but overlap black maple.
Whole tree / context	A large, long-lived canopy tree with a dense rounded crown in cool, fertile, well-drained northern hardwoods, commonly with beech, yellow birch, basswood, and hemlock. Site is corroboration, not a substitute for morphology.
Variation / condition	Shade leaves may be thin and shallowly lobed; vigorous shoots can carry larger or more toothed leaves. Black-maple intergradation, crown damage, old bark, lichens, wetness, and logging scars can weaken textbook characters.
Evidence limit	Opposite arrangement plus several representative leaves and pointed brown buds normally supports species. One detached atypical leaf, distant crown color, or old plated bark alone does not securely separate sugar maple from black maple, red maple, or planted Norway maple.

Confuser control

Alternative	Visible separator	Resolving view
<i>Acer rubrum</i>	Red maple usually has regular fine teeth around the lobes and into sharper sinuses, red blunt-ovoid buds, and spring-maturing samaras; sugar maple has fewer coarse teeth, rounded sinuses, narrow brown pointed buds, and autumn fruit.	An intact opposite shoot with terminal and lateral buds, the full leaf margin, and mature samaras if present.
<i>Acer nigrum</i> and <i>Acer platanoides</i>	Black maple tends toward three broad drooping lobes, a deeply cordate base, and lower-surface hair but commonly intergrades. Norway maple has milky petiole latex, large blunt buds, and samara wings spread nearly straight across.	Several mature leaf undersides and petiole bases, a fresh petiole break, buds, samaras, and site context.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Very shade tolerant; seedlings and saplings can persist for years as advance regeneration and accelerate after a gradual increase in light.
Moisture / soils	Best on cool, fertile, moist but well-drained loams with adequate calcium and magnesium; performs poorly on droughty shallow soils and is intolerant of prolonged growing-season flooding.
Geographic affinity	A cool-temperate Great Lakes, southern Canadian, northeastern, and Appalachian species whose strongest dominance is in humid continental northern hardwood climates.
Associations	A principal canopy component of sugar maple-basswood and sugar maple-beech-yellow birch forests, with hemlock, white ash, black cherry, and other mesic associates.
Regeneration	Wind-dispersed autumn samaras germinate after cold moist stratification; abundant seedlings establish beneath canopy and form advance regeneration before release.
Vegetative reproduction	Young and small stumps can sprout, and damaged seedlings readily replace shoots; root suckering is rare and seed is the principal stand-regeneration pathway.
Seed / mast	Good crops recur irregularly, often every few years; seed usually does not remain viable in the forest floor beyond the following growing season.
Establishment	Seedlings can penetrate litter and establish under shade on moist forest floors, but need sustained soil moisture, controlled competition, and eventual crown release for height growth.
Succession	Late-successional canopy dominant that can replace less-tolerant pioneers and maintain uneven-aged structures through advance regeneration and gap recruitment.
Disturbance response	Small gaps and partial cutting release established regeneration; fire, severe drought, soil compaction, root damage, and careless bole wounding are damaging.
Longevity / growth	Long-lived, commonly several centuries on good sites; slow while suppressed, then moderate and persistent in diameter growth after release.
Browse	Deer readily browse seedlings and saplings, and repeated browsing can prevent advance regeneration from reaching recruitment height.
Insects / disease	Maple decline, sugar maple borer, defoliators, cankers, and decay following wounds are important; stress interactions often matter more than a single agent.
Management	Protect roots and boles during harvest, match retention to fertile mesic sites, and evaluate browse and competing understory before relying on abundant small seedlings.
Silvicultural systems	Selection, group selection, and shelterwood approaches can work where healthy advance regeneration is distributed and released gradually; excessive high grading and abrupt exposure reduce quality and vigor.

Personal field cue for Sugar maple

learner-created retrieval hook

Red maple - *Acer rubrum*

New this week | Family: Sapindaceae | Tier A | Priority 1 - core

Nearly throughout the region from swamps to dry uplands

Ubiquitous generalist; prolific seeder and sprouter after disturbance



Bark

(c) Bryce, some rights reserved (CC BY)
INAT_PHOTO_458152552 | cc-by



Condition

(c) Steven Lamonde, some rights...
INAT_PHOTO_699498715 | cc-by



Foliage

no rights reserved
INAT_PHOTO_717573410 | cc0

Evidence	Diagnostic interpretation
Foliage	Opposite, simple blades usually have three to five lobes, angular or V-shaped sinuses, and regular serration continuing around the lobes and into the sinuses; the lower surface is often pale. Red petioles or fall color are supportive but not diagnostic.
Bark age series	Young stems are smooth and light gray; intermediate bark develops narrow vertical cracks and small scaly plates; old boles become dark, furrowed, and broken into long narrow plates that may lift at the ends. Age and site produce substantial variation.
Dormant	Opposite reddish twigs bear blunt-ovoid red buds with several pairs of scales; flower buds are often clustered and plumper than vegetative buds. The terminal bud is much shorter and blunter than sugar maple's narrow brown point.
Fruit / cone / seed	Red to yellowish flowers open well before leaf-out, and paired samaras mature and fall in spring or early summer. Early fruit timing is a strong separator from sugar and Norway maples.
Whole tree / context	A medium to large tree with a variable crown occurring from forested wetlands and riparian flats to mesic and dry uplands. Broad ecological amplitude raises plausibility but contributes little species-level proof.
Variation / condition	Lobe number, depth, pubescence, color, and bark vary widely among sites and individuals. Suppressed shoots, stump sprouts, drought, flooding, lichens, and injury can make leaves or bark atypical.
Evidence limit	A secure call combines opposite arrangement with the serrated lobe pattern, red blunt buds, spring fruit, or a second independent organ. A red-colored twig, one ambiguous maple leaf, or rough old bark alone warrants <i>Acer</i> or insufficient evidence.

Confuser control

Alternative	Visible separator	Resolving view
<i>Acer saccharum</i>	Sugar maple has fewer large teeth, usually broader rounded sinuses, narrow sharply pointed brown buds, and autumn-maturing samaras; red maple is more continuously serrate with blunt red buds and spring fruit.	The complete margin and sinuses, intact terminal buds, and fruit with season information.
<i>Acer saccharinum</i>	Silver maple generally has much deeper sinuses, narrower lobes, a strongly silvery-white lower surface, more conspicuously shaggy mature bark, and larger spring samaras.	Both leaf surfaces with scale, mature bark, and measured fruit.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Moderately shade tolerant when young and able to persist beneath partial canopy, but faster growth and canopy recruitment follow disturbance and increased light.
Moisture / soils	Occupies an exceptionally broad range from saturated swamps and floodplains to dry ridges; local seed sources are often adapted to contrasting hydrologic regimes.
Geographic affinity	One of the most geographically and climatically widespread eastern trees, common across the entire course envelope.

Dimension	Field-use summary
Associations	Occurs in red maple wetlands, northern hardwoods, oak-hardwood, pine-hardwood, aspen-birch, and many disturbed or transitional stands.
Regeneration	Produces abundant wind-dispersed spring seed that germinates promptly on moist substrates; advance seedlings and post-disturbance cohorts are both common.
Vegetative reproduction	Stumps sprout vigorously, especially when cut young, and sprouts can dominate early stand development after harvest or top-kill.
Seed / mast	Seed crops begin at relatively young ages and are frequent, often annual; short-lived seed germinates soon after spring dispersal rather than forming a persistent bank.
Establishment	Moist seedbeds and early-season light favor germination, while broad physiological tolerance and sprouting permit establishment across many disturbed settings.
Succession	A flexible early- to mid-seral generalist that can persist into later stands as a subordinate canopy tree, especially where repeated disturbance limits more demanding species.
Disturbance response	Logging, fire top-kill, agricultural abandonment, drainage change, and canopy gaps often increase red maple through seed and sprouts; severe drought or prolonged flooding response varies by ecotype.
Longevity / growth	Moderately fast when released and usually shorter-lived than sugar maple, beech, or hemlock; poor form and internal defects are common on stressed sites.
Browse	Young shoots are browsed by deer and moose, though prolific seedling and sprout production can maintain abundance under moderate pressure.
Insects / disease	Defoliators, leafhoppers, borers, cankers, decay, and fire or logging wounds affect quality; no single agent generally controls its regional abundance.
Management	Do not infer site quality from presence alone; decide whether red maple is an objective species, protect selected crop trees from injury, and account for vigorous sprouting in regeneration treatments.
Silvicultural systems	Responds to clearcut, shelterwood, group, and selection openings; outcomes depend on advance regeneration, stump size, seed source, competing vegetation, and the desired species mixture.

Personal field cue for Red maple

learner-created retrieval hook

American beech - *Fagus grandifolia*

New this week | Family: Fagaceae | Tier A | Priority 1 - core

Widespread from the Great Lakes through southern Canada, New England, NB, and the Allegheny region

Major shade-tolerant mast tree; root sprouting and disease strongly affect stand structure



Bark

(c) Nick Kleinschmidt, some rights reserved
INAT_PHOTO_650866298 | cc-by



Foliage

(c) Matt Pelikan, some rights reserved
INAT_PHOTO_602541471 | cc-by



Mixed Evidence

no rights reserved
INAT_PHOTO_631749583 | cc0

Evidence	Diagnostic interpretation
Foliage	Alternate, simple leaves are elliptic to ovate with a short petiole, straight parallel lateral veins, and one distinct marginal tooth at or near each vein ending. The surface is papery rather than corrugated, and saplings often retain tan dead leaves in winter.
Bark age series	Saplings and healthy mature trees have thin, smooth, pale bluish-gray bark, often compared with elephant hide. Scale insects, beech bark disease, cankers, callus, wounds, lichen, wetness, and old aftermath roughening can partly or almost completely obscure that anchor pattern.
Dormant	Alternate twigs carry exceptionally long, slender, sharply pointed cigar-shaped buds with many overlapping scales; the buds project away from the twig. This is one of the most reliable winter characters in the regional flora.
Fruit / cone / seed	Male flowers occur in drooping spherical clusters; paired sharply three-angled nuts mature inside a four-part prickly husk. Detached triangular nuts or open husks strongly support beech when provenance is secure.
Whole tree / context	A large tree with a broad crown, smooth-looking boles, shallow spreading roots, and frequent root-sucker thickets in mesic northern hardwoods. Dense shade and beech saplings may persist below diseased overstory trees.
Variation / condition	Host identification must be separated from diagnosis. Clean juvenile bark, normal smooth bark, scale patches, cankers, rough aftermath bark, callus, scars, wet bark, lichen, and foliage affected by beech leaf disease can all occur on <i>Fagus grandifolia</i> ; a photograph cannot establish genetic resistance.
Evidence limit	Cigar buds, vein-tooth foliage, nuts, or multiple host characters can identify beech even when bark is altered. Smooth gray bark alone is not enough, and rough bark alone cannot diagnose beech bark disease, disease stage, cause, resistance, or tolerance.

Confuser control

Alternative	Visible separator	Resolving view
<i>Ostrya virginiana</i>	Hophornbeam has shorter ovoid buds, doubly serrate corrugated leaves, narrow shreddy bark strips, and hoplike fruit; beech has very long cigar buds, a simpler vein-tooth margin, smooth host bark, and prickly nut husks.	Dormant buds, close leaf venation and margin, representative bark away from wounds, and fruit.
<i>Carpinus caroliniana</i>	American hornbeam has smooth muscle-like fluted bark but much shorter buds, doubly serrate leaves, and nutlets on three-lobed leafy bracts rather than beechnuts in prickly husks.	A full twig with buds, leaf margin, bole form, and reproductive structures.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Exceptionally shade tolerant; seedlings, saplings, and root suckers can persist under dense canopy and respond when gaps form.
Moisture / soils	Most competitive on moist, well-drained, often fertile loams and slopes; absent from persistently flooded sites and less vigorous on droughty shallow soils.

Dimension	Field-use summary
Geographic affinity	A broad eastern cool- and warm-temperate species, central to Great Lakes, Acadian, northeastern, and Allegheny northern hardwoods.
Associations	Major component of beech-sugar maple and sugar maple-beech-yellow birch forests, also associated with hemlock, red spruce, white pine, oaks, and Allegheny hardwoods.
Regeneration	Heavy nuts establish seedlings after autumn dispersal, while existing advance regeneration and root sprouts often contribute more consistently to local replacement.
Vegetative reproduction	Root suckering is important and may intensify after root injury, overstory mortality, or cutting, producing dense genetically connected thickets.
Seed / mast	Beechnut production is episodic, with irregular mast years that strongly affect wildlife food and seedling pulses; rodents and birds consume and disperse nuts.
Establishment	Seedlings tolerate litter and deep shade on mesic sites, but recruitment above browse height and into canopy positions generally depends on protection and gap release.
Succession	Late-successional, long-persistent canopy species that can dominate regeneration layers even where overstory quality is reduced by disease.
Disturbance response	Canopy gaps release suppressed stems; root damage and cutting stimulate suckers, while fire and bole injury readily damage thin bark and create decay entry points.
Longevity / growth	Long-lived and slow-growing under suppression, capable of large size but often limited in current forests by disease, defects, and repeated mortality.
Browse	Deer browse seedlings and suckers; intense browsing can simplify regeneration while beech's persistent clonal shoots remain abundant at low height.
Insects / disease	Beech scale and fungi associated with beech bark disease, beech leaf disease, cankers, decay, and defoliators are regionally important; symptoms and host identity require separate evidence.
Management	Inventory both overstory condition and regeneration origin, avoid assuming every rough-barked beech is in the same disease stage, retain valuable mast and healthy phenotypes where objectives permit, and manage dense sucker competition deliberately.
Silvicultural systems	Selection and shelterwood openings release advance beech and suckers strongly; without regeneration control they may outcompete desired associates, while sanitation decisions must consider mast, diversity, disease, and residual damage.

Personal field cue for American beech
learner-created retrieval hook

Yellow birch - *Betula alleghaniensis*

New this week | Family: Betulaceae | Tier A | Priority 1 - core

Cool mesic forests from MN/WI/MI through ON/QC, NY, New England, NB, and Allegheny highlands

Long-lived northern-hardwood dominant; establishes on decayed wood and disturbed mineral soil



Bark

(c) Amy MacPherson, some rights reserved...
INAT_PHOTO_574455120 | cc-by-sa



Foliage

(c) Alan Prather, some rights reserved...
INAT_PHOTO_705760040 | cc-by



Mixed Evidence

(c) Nash Turley, some rights reserved...
INAT_PHOTO_702302386 | cc-by

Evidence	Diagnostic interpretation
Foliage	Alternate ovate leaves have a rounded to weakly heart-shaped base, many straight lateral veins, and sharply doubly serrate margins; lower vein axils may be hairy. Foliage alone overlaps sweet birch and needs bark, twig odor, or reproductive support.
Bark age series	Seedlings are gray-brown and unremarkable; intermediate stems develop bronze to yellow-gold bark with horizontal lenticels and fine curling papery strips; old trunks become dark reddish-brown to gray and break into rough irregular plates with much less visible curl.
Dormant	Slender alternate twigs bear pointed reddish-brown buds, small spur shoots, and no true terminal bud; scratched young twig or inner bark has a wintergreen odor. Odor should be wafted rather than tasted and does not separate yellow from sweet birch.
Fruit / cone / seed	Male catkins form in advance and persist through winter; erect female strobiles mature and disintegrate into tiny winged nutlets and three-lobed scales. Complete fruit scales can help separate birches when clearly imaged.
Whole tree / context	A large long-lived canopy tree of cool moist northern hardwood and mixedwood sites, often rooted on old logs, stumps, tip-up mounds, or boulders with stilted bases after the nurse substrate decays.
Variation / condition	Classic golden curl is strongest at intermediate age. Shade, wet bark, lichens, scars, and very old plated bark change color and texture; sprouts and saplings may lack the feature learners expect.
Evidence limit	Golden curling bark plus horizontal lenticels is strong evidence, and wintergreen odor corroborates the birch group. A dark old bole, a sterile leaf, or odor alone cannot securely exclude sweet birch, black cherry, or other <i>Betula</i> .

Confuser control

Alternative	Visible separator	Resolving view
<i>Betula lenta</i>	Both can smell of wintergreen and have similar leaves, but sweet birch develops dark nonpapery bark with horizontal lenticels and later heavy plates, whereas yellow birch shows characteristic bronze-gold fine curling at intermediate age.	Intermediate-age bark, several mature leaves including lower surfaces, buds, and intact fruiting scales.
<i>Prunus serotina</i>	Young black cherry can show horizontal lenticels and old bark becomes cornflake-like, but cherry leaves have petiole glands and rusty hair along the lower midrib and fruit in fleshy racemes; yellow birch has catkins and papery curl.	Current twigs, petiole and lower midrib, intermediate bark, and fruit or catkins.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Intermediate in shade tolerance: more tolerant than paper birch and aspen, but less able than sugar maple, beech, or hemlock to persist and recruit under deep shade.
Moisture / soils	Best on cool, moist, well-drained, fertile loams and coves; common on north and east slopes, seepage areas, ravines, and decayed woody microsites but intolerant of stagnant flooding.

Dimension	Field-use summary
Geographic affinity	A cool-humid northern and montane species spanning the Great Lakes, southern Canada, Acadian region, Northeast, and Allegheny/Appalachian highlands.
Associations	Characteristic with sugar maple and beech, and with hemlock, red spruce, balsam fir, paper birch, white pine, and northern hardwood associates.
Regeneration	Very small wind-dispersed seed supplies new cohorts; seedlings commonly establish on rotten wood, exposed mineral soil, tip-up mounds, and lightly scarified moist surfaces.
Vegetative reproduction	Young stumps can sprout and low branches may layer, but vegetative reproduction is less reliable than seed for maintaining canopy populations.
Seed / mast	Seed production is variable with periodic good crops; much seed falls near the parent, and viability and first-year survival depend strongly on moisture and seedbed quality.
Establishment	Tiny seedlings are easily smothered by intact litter and drought; moist decayed wood or mineral soil, partial light, and limited competition provide receptive seedbeds.
Succession	A long-lived mid- to late-seral gap species that can enter after disturbance yet persist with northern-hardwood climax associates.
Disturbance response	Windthrow, partial harvest, soil exposure, and group gaps create seedbeds and light; severe opening can dry seedbeds, while fire and bole injury damage thin bark.
Longevity / growth	Long-lived for a birch and capable of large canopy size; juvenile growth is moderate and improves with overhead and side light.
Browse	Deer and moose browse seedlings and saplings; repeated browsing can prevent otherwise suitable regeneration from reaching gap height.
Insects / disease	Bronze birch borer, birch leafminers and defoliators, cankers, decay, and root or bole injuries affect vigor; stressed exposed trees are more vulnerable.
Management	Coordinate seed source, receptive seedbed creation, moisture retention, and timely light; protect residual boles and roots and do not rely on advance regeneration alone where seedlings are scarce.
Silvicultural systems	Group selection, irregular shelterwood, and shelterwood with controlled scarification can recruit yellow birch; single-tree openings may be too small for growth, while clear exposure can make seedbeds too dry.

Personal field cue for Yellow birch

learner-created retrieval hook

Paper birch - *Betula papyrifera*

New this week | Family: Betulaceae | Tier A | Priority 1 - core


Boreal and northern-transition forests across the Great Lakes, ON/QC, northern NY/New England, and NB

Short-lived shade-intolerant pioneer after fire, logging, and severe disturbance




Bark

(c) Quinten Wiegiersma, some rights...
INAT_PHOTO_564773858 | cc-by



Foliage

(c) Dan MacNeal, some rights reserved...
INAT_PHOTO_712523142 | cc-by



Mixed Evidence

(c) Brandon Corder, some rights...
INAT_PHOTO_650441630 | cc-by

Evidence	Diagnostic interpretation
Foliage	Alternate ovate to triangular leaves have a rounded or wedge-shaped base, pointed tip, and doubly serrate margin; lower vein axils may bear hair. Mature heart-shaped blades can overlap heart-leaved paper birch, so foliage must be sampled across the crown.
Bark age series	Young stems are reddish-brown with pale horizontal lenticels and do not look white; pole and mature bark becomes creamy to chalky white with dark lenticels and peels in thin papery sheets; old basal bark becomes dark, thick, and deeply fissured.
Dormant	Slender reddish-brown alternate twigs have pointed appressed buds, conspicuous resin glands or wartlike dots on many current shoots, no true terminal bud, and persistent paired or clustered male catkins.
Fruit / cone / seed	Pendulous male catkins expand in spring; erect cylindrical female strobiles mature then disintegrate into tiny winged nutlets and three-lobed scales. Scale proportions can be diagnostic but require close, complete material.
Whole tree / context	A small to medium, often multi-stemmed, narrow-crowned pioneer forming bright stands after fire or harvest in cool boreal and northern-transition landscapes.
Variation / condition	White bark develops with age and varies from bright white to cream or pinkish. Wetness, lichens, fire scars, old basal furrows, and hybrid or varietal variation can obscure the familiar surface.
Evidence limit	White peeling bark plus appropriate twigs, foliage, or fruit usually supports paper birch, but bark alone does not reliably separate heart-leaved paper birch and can be confused with gray birch. Juvenile brown bark requires other organs.

Confuser control

Alternative	Visible separator	Resolving view
<i>Betula cordifolia</i>	Heart-leaved paper birch tends to have cordate mature leaf bases, more resinous or glandular shoots, and montane or cool northeastern settings, but bark is nearly indistinguishable and character overlap is substantial.	Multiple fully expanded crown leaves, both current and one-year twigs, buds, complete fruiting scales, elevation, substrate, and geography.
<i>Betula populifolia</i>	Gray birch usually has triangular long-pointed leaves, white chalky bark that peels little, and dark triangular marks beneath branch attachments; paper birch more freely exfoliates in sheets.	A representative mature leaf, bark around several branch bases, peeling-layer detail, and fruit scales.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Very shade intolerant; seedlings require abundant light and are rapidly overtopped in closed stands.
Moisture / soils	Occupies cool sites across a broad moisture range, from well-drained glacial tills and sandy soils to moist margins, but grows best where rooting moisture is adequate without prolonged stagnation.
Geographic affinity	Strongly boreal and northern-temperate, widespread through the Great Lakes, Canada, Acadian forests, and cool northeastern uplands.
Associations	Common with aspen, balsam fir, white and black spruce, jack and white pine, tamarack, northern hardwoods, and early-seral mixedwoods.

Dimension	Field-use summary
Regeneration	Produces abundant tiny wind-dispersed seed that colonizes exposed mineral soil, burns, road edges, and disturbed substrates after canopy removal.
Vegetative reproduction	Young stumps can sprout, especially after cutting, but sprout vigor declines with age and seed is the main landscape-scale colonization mechanism.
Seed / mast	Seed production is frequent and can be heavy; small seeds disperse widely over snow or open ground, though most seedlings die unless a moist exposed seedbed and light coincide.
Establishment	Open light, exposed mineral or lightly covered soil, reliable near-surface moisture, and limited early competition are critical; thick litter and deep shade prevent recruitment.
Succession	Classic short-lived pioneer following stand-replacing fire, logging, windthrow, or land clearing, later yielding to more shade-tolerant conifers and hardwoods.
Disturbance response	Fire and severe harvesting create establishment opportunities when seed survives nearby or disperses in; surviving young stems may sprout, but mature thin-barked trees are easily killed or injured.
Longevity / growth	Fast juvenile height growth and early reproduction, commonly declining after several decades and rarely persisting as a dominant for the lifespan of late-seral associates.
Browse	Moose, deer, and snowshoe hare heavily use shoots and seedlings; repeated browse deforms saplings and can alter early-seral mixtures.
Insects / disease	Bronze birch borer, birch leafminers, defoliators, cankers, decay, and drought or heat stress can accelerate decline, especially outside cool favorable sites.
Management	Maintain seed sources and deliberately create adequately large, moist, exposed seedbeds where paper birch is an objective; expect shade-tolerant competitors to replace it without repeated disturbance.
Silvicultural systems	Clearcutting, seed-tree, and large shelterwood or group openings can regenerate it when mineral soil and seed are available; small selection gaps generally provide too little sustained light.

Personal field cue for Paper birch
learner-created retrieval hook

Eastern hemlock - *Tsuga canadensis*

New this week | Family: Pinaceae | Tier A | Priority 1 - core

Great Lakes through southern ON/QC, NY, New England, NB, and the Allegheny region

Extremely shade-tolerant long-lived foundation conifer creating cool moist microclimates



Bark

no rights reserved
INAT_PHOTO_462391786 | cc0



Foliage

(c) Christina S. Baer, some rights reserved
INAT_PHOTO_713743092 | cc-by



Mixed Evidence

(c) Timon T. K., some rights reserved...
INAT_PHOTO_619989178 | cc-by

Evidence	Diagnostic interpretation
Foliage	Short, flat, mostly blunt single needles vary in length along delicate twigs, show two pale stomatal bands beneath, and narrow abruptly to tiny petiole-like bases. Needles often appear two-ranked but small upper-side needles interrupt a perfectly flat spray.
Bark age series	Young bark is smooth gray-brown; intermediate stems develop narrow scaly ridges; mature bark becomes dark gray-brown and deeply furrowed with broad ridges, while cuts expose cinnamon to reddish inner bark. Bark alone overlaps other conifers.
Dormant	Fine flexible yellow-brown twigs carry minute rounded buds, tiny petiolate needles, and often a drooping terminal shoot. Hemlock lacks the woody sterigmata of spruce and the suction-cup-like needle bases of fir.
Fruit / cone / seed	Small ovoid pendent cones, usually only about 1.5 to 2.5 cm long, hang from short stalks and open at maturity; they persist beneath branches. Their size and position sharply contrast with upright fir cones.
Whole tree / context	A graceful conifer with fine feathery sprays, slender horizontal branches, and a drooping leader, often forming dense, deeply shaded groves on cool ravines, slopes, stream terraces, and lake-influenced sites.
Variation / condition	Shade and vigor change shoot density and needle length. Hemlock woolly adelgid, elongate hemlock scale, drought, browsing, and other stress can thin crowns or discolor and shed needles; decline symptoms do not erase attachment characters.
Evidence limit	Needle attachment, lower stomatal bands, twig texture, and a small pendent cone together are decisive. A distant dark crown, deeply furrowed bark, or one cropped needle surface cannot securely separate hemlock from fir, spruce, yew, or planted conifers.

Confuser control

Alternative	Visible separator	Resolving view
<i>Abies balsamea</i>	Balsam fir needles attach on broad round bases that leave smooth circular scars, terminal twigs are stouter, resin blisters occur on young bark, and mature cones stand upright and disintegrate; hemlock needles have tiny petioles and cones hang intact.	Needle bases after removal, lower needle surfaces, terminal buds, young bark, and cone orientation.
<i>Picea</i> species	Spruce needles are usually stiff and four-angled or keeled and sit on persistent woody pegs that make twigs rough; hemlock needles are flat, petiolate, and leave smoother twigs.	A needle cross-section, bare twig after needle removal, underside stomatal bands, and complete cone.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Extremely shade tolerant, among the most tolerant North American trees; seedlings and saplings can remain suppressed for many decades and still respond to release.
Moisture / soils	Favors cool, humid, sheltered sites with reliable moisture and generally well-drained acidic to neutral soils; ravines, north slopes, stream benches, and lake-moderated sites are common.

Dimension	Field-use summary
Geographic affinity	A cool-temperate eastern species central to Great Lakes, Acadian, northeastern, and Allegheny mixedwoods, with southern occurrences concentrated in cool refugia.
Associations	Forms pure groves and mixtures with white pine, yellow birch, sugar maple, beech, red spruce, balsam fir, and northern red oak.
Regeneration	Tiny wind-dispersed seed establishes on continuously moist rotten wood, moss, humus, or mineral soil; abundant advance regeneration can persist under parent stands.
Vegetative reproduction	Natural sprouting is negligible; layering may occur from low branches in far northern or harsh settings, but seed is the normal regeneration pathway.
Seed / mast	Cones are produced periodically, with good crops recurring every few years; seed disperses in autumn and winter but loses viability relatively quickly on the forest floor.
Establishment	Moist, cool microsites protected from drying are essential for germination and early survival; decayed logs and shaded mineral soil often function as nurse substrates.
Succession	Very late-successional and self-replacing where disturbance remains small, creating deep shade, cool microclimates, and slowly cycling forest-floor conditions.
Disturbance response	Small gaps and careful partial cutting release advance regeneration, but abrupt exposure, drought, fire, windthrow, and root or bole injury can cause severe mortality.
Longevity / growth	Very long-lived, often several centuries; exceptionally slow while suppressed and capable of sustained growth after release if roots and crown remain healthy.
Browse	Highly preferred winter browse for white-tailed deer; heavy browsing can eliminate regeneration and alter stand structure even where seed is abundant.
Insects / disease	Hemlock woolly adelgid is the principal invasive threat in much of the region; elongate hemlock scale, hemlock looper, root rots, drought, and secondary agents also matter.
Management	Protect cool moist microclimates, roots, and residual crowns; monitor invasive insects; evaluate deer pressure; and retain structural and riparian functions even where timber objectives are secondary.
Silvicultural systems	Single-tree or small-group selection and gradual shelterwood release can favor established hemlock on suitable sites, while heavy uniform opening risks desiccation, windthrow, competing vegetation, and loss of advance regeneration.

Personal field cue for Eastern hemlock

learner-created retrieval hook

American basswood - *Tilia americana*

New this week | Family: Malvaceae; Tiliaceae in older texts | Tier A | Priority 1 - core

Fertile mesic sites from MN/WI/MI and southern Canada through the Northeast and PA/NJ

Shade-tolerant nutrient-rich northern-hardwood associate; vigorous stump sprouter




Bark

(c) Loïc Mathieu, some rights reserved...
INAT_PHOTO_608142436 | cc-by



Dormant

(c) Loïc Mathieu, some rights reserved...
INAT_PHOTO_607898739 | cc-by



Foliage

(c) sofi v, some rights reserved (CC BY)
INAT_PHOTO_713542533 | cc-by

Evidence	Diagnostic interpretation
Foliage	Alternate, broadly ovate to heart-shaped leaves have an abruptly pointed tip, serrate margin, palmate basal veins, and a conspicuously uneven cordate base. Size varies greatly, especially on stump sprouts, so asymmetry and structure outweigh dimensions.
Bark age series	Young bark is smooth gray to gray-brown; intermediate bark develops narrow vertical fissures; old boles are dark gray-brown with long interlacing ridges and deep furrows. Multi-stemmed sprout clumps are common after cutting.
Dormant	Alternate zigzag twigs carry plump red to reddish-brown ovoid buds with only two or three visible, unequal scales; there is no true terminal bud. Bud scale count and asymmetrical leaf scars are useful winter evidence.
Fruit / cone / seed	Clusters of fragrant pale flowers and later round nutlets hang from a stalk fused partway to a narrow leaf-like bract. The bract-and-fruit construction is one of the most decisive regional reproductive characters.
Whole tree / context	A medium to large, often straight-boled tree of fertile mesic northern hardwoods and rich lower slopes; basal sprouts or multiple trunks commonly reveal vegetative origin.
Variation / condition	Vigorous sprouts bear extremely large leaves; shade leaves are thinner. Decay, fire scars, browse, and old furrowed bark may mask typical form, while planted European and hybrid lindens complicate settled-site identification.
Evidence limit	An asymmetrical cordate leaf plus two-scaled buds or a fruiting bract supports American basswood in forest context. Foliage alone may justify only <i>Tilia</i> where planted lindens occur, and large leaf size alone is not diagnostic.

Confuser control

Alternative	Visible separator	Resolving view
<i>Morus alba</i>	White mulberry leaves are variably lobed, often glossy, and have different buds and fleshy multiple fruits; basswood has plump few-scaled red buds and flowers or nutlets attached to a distinctive narrow bract.	Buds and twig, several leaves from one shoot, reproductive structures, and whether the tree is planted or in forest.
Planted <i>Tilia</i> species and hybrids	Other lindens share the bracted fruit and cordate foliage; lower-surface hair, fruit ribbing, leaf dimensions, buds, provenance, and horticultural context may be needed, and photographs often support only genus.	Lower leaf surface with vein axils, mature fruit with scale, buds, bark, crown form, and planting context.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Shade tolerant, though generally less persistent in deep shade than beech or sugar maple; advance stems and sprouts respond strongly to release.
Moisture / soils	Best on deep, fertile, moist, well-drained loams with high base status, especially calcium-rich tills, lower slopes, and protected coves; poor on droughty or saturated sites.
Geographic affinity	A broad eastern species especially important in the western Great Lakes sugar maple-basswood region and rich northeastern and Allegheny sites.
Associations	Common with sugar and black maples, white ash, elms, bitternut hickory, yellow birch, beech, and other rich-site hardwoods.

Dimension	Field-use summary
Regeneration	Seedlings occur after successful dormancy break and favorable seedbeds, but many post-harvest cohorts arise from vigorous stump sprouts.
Vegetative reproduction	Stump sprouting is exceptionally vigorous, particularly from young and intermediate stems; multiple sprouts can rapidly recapture growing space after cutting.
Seed / mast	Nutlets are produced periodically and dispersed mainly near the parent; deep dormancy and frequent low soundness make natural seedling establishment less dependable than sprouting.
Establishment	Seed requires prolonged stratification and suitable moist forest soil; advance seedlings tolerate shade, while released sprouts need adequate growing space and browse protection.
Succession	A shade-tolerant mid- to late-seral rich-site associate that persists by both seedlings and vegetative replacement rather than acting as a broad pioneer.
Disturbance response	Cutting strongly stimulates stump sprouts; fire and mechanical wounds readily injure thin bark and soft wood, leading to decay, while partial release can improve crown growth.
Longevity / growth	Moderately long-lived and capable of rapid diameter growth on rich sites; soft wood and decay following injury often limit quality before physiological lifespan.
Browse	Seedlings and sprouts are highly palatable to deer and other browsers; repeated browsing can suppress the otherwise vigorous vegetative response.
Insects / disease	Basswood borer, defoliators, leaf spots, cankers, and several decay fungi occur; injury and poor vigor often determine severity.
Management	Use basswood as an indicator and component of fertile sites, protect high-quality residual stems from damage, and account for dense stump sprouting and browse in regeneration plans.
Silvicultural systems	Selection and shelterwood systems can release advance trees, while coppice and even-aged harvests produce vigorous sprout clumps; thinning sprout clumps may improve form where basswood is a crop objective.

Personal field cue for American basswood
learner-created retrieval hook

Black cherry - *Prunus serotina*

New this week | Family: Rosaceae | Tier A | Priority 1 - core

Throughout much of the region; especially important on the Allegheny Plateau

Valuable gap- and disturbance-responsive timber tree; bird-dispersed fruit



Bark

(c) Patrick Jackson, some rights...
INAT_PHOTO_580635573 | cc-by



Foliage

no rights reserved
INAT_PHOTO_713870593 | cc0



Reproductive

(c) Rebecca Turner, some rights...
INAT_PHOTO_651110401 | cc-by

Evidence	Diagnostic interpretation
Foliage	Alternate, simple, elliptic leaves have fine rounded or incurved teeth, one or more small glands near the top of the petiole, and characteristic rusty to pale hairs along the lower midrib near its base. Several characters should be visible before making a foliage-only call.
Bark age series	Young stems are smooth reddish-brown to dark gray with prominent horizontal lenticels; intermediate bark breaks into small dark scaly plates with lifted edges; mature bark forms rough irregular burnt-cornflake plates. Old bases may become deeply furrowed.
Dormant	Slender alternate reddish-brown twigs carry small ovoid scaled buds and semicircular leaf scars; scratched twigs have a bitter-almond odor from cyanogenic compounds. Waft odor only and never taste or ingest plant material.
Fruit / cone / seed	Many small white flowers occur in elongated cylindrical racemes after leaves expand; round drupes mature from red to purple-black on the same raceme. The raceme architecture separates black cherry from clustered chokecherry look-alikes when fully visible.
Whole tree / context	A medium to large canopy tree with a relatively narrow crown, prominent dark scaly bole, and seedlings common in gaps, edges, and Allegheny hardwood stands.
Variation / condition	Shade leaves, stump sprouts, leaf spot, black knot, tent caterpillar defoliation, wet bark, and old furrowed bases alter appearance. Juvenile smooth bark looks unlike the classic mature cornflake surface.
Evidence limit	Mature plated bark is highly suggestive but should be corroborated with leaves, twigs, or fruit. A single generic <i>Prunus</i> leaf, distant black fruits, almond odor, or young lenticled bark alone may support only genus.

Confuser control

Alternative	Visible separator	Resolving view
<i>Prunus pensylvanica</i> and <i>Prunus virginiana</i>	Pin cherry has smaller narrow leaves, bright reddish bark, and fruits in short umbel-like clusters; chokecherry is usually shrub or small tree with broader leaves and shorter dense racemes. Black cherry's mature cornflake bark, petiole glands, lower-midrib hair, and long racemes are decisive in combination.	Whole fruit-cluster attachment, petiole glands, lower midrib, buds, representative bark age, and whole-tree stature.
<i>Betula alleghaniensis</i> and <i>Betula lenta</i>	Birches can have dark lenticled or plated bark and wintergreen odor, but bear catkins, lack fleshy racemes and cherry petiole glands, and have different buds; yellow birch usually shows bronze papery curl at intermediate age.	Intermediate bark, reproductive structures, current twig buds, petiole and lower leaf surface.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Shade intolerant; seedlings may survive briefly under partial canopy, but successful height growth and timber form require substantial light.
Moisture / soils	Best on deep, fertile, moist but well-drained loams, especially Allegheny uplands and protected slopes; poor on saturated, severely droughty, or compacted sites.

Dimension	Field-use summary
Geographic affinity	Broadly eastern, with exceptional commercial and ecological importance on the Allegheny Plateau and good northern Appalachian sites.
Associations	Occurs with red and sugar maples, yellow-poplar, white ash, northern red oak, beech, birches, and other Allegheny or northern-hardwood species.
Regeneration	Birds and mammals disperse drupes; dormant seeds accumulate in the forest floor and germinate after canopy disturbance creates light and favorable temperatures.
Vegetative reproduction	Young stumps sprout readily and damaged seedlings can replace shoots, though high-quality stand recruitment often depends on seed-bank seedlings.
Seed / mast	Fruit crops are frequent but variable; animal dispersal is important, and seeds can remain dormant in soil for multiple years until gap conditions trigger germination.
Establishment	Mineral soil or thin litter, adequate moisture, and strong overhead light favor establishment; seedlings fail under prolonged deep shade or severe herbaceous and fern competition.
Succession	Early- to mid-seral gap and disturbance species that can become a valuable canopy component, then declines in recruitment as shade-tolerant species close the stand.
Disturbance response	Windthrow, harvesting, fire-created openings, and soil disturbance can activate the seed bank and release seedlings; poorly timed or undersized gaps may not provide enough light.
Longevity / growth	Rapid juvenile growth and potentially large size on good sites, with moderate lifespan and strong early allocation to height in competition.
Browse	Deer browse seedlings and sprouts heavily; abundant cohorts can still fail to recruit where browse, ferns, or competing vegetation remain intense.
Insects / disease	Eastern tent caterpillar, black knot, cherry scallop shell moth, borers, leaf spots, cankers, and decay affect form and vigor; damaged foliage and bark should not be mistaken for species characters.
Management	On productive Allegheny sites, coordinate seed-bank potential, competing vegetation control, deer pressure, and sufficiently large openings; retain quality seed sources and avoid residual bole damage.
Silvicultural systems	Shelterwood and even-aged regeneration methods with preparatory vegetation control can produce cohorts; group openings must be large enough for sustained light, while single-tree selection generally favors more tolerant competitors.

Personal field cue for Black cherry

learner-created retrieval hook

Eastern hophornbeam - *Ostrya virginiana*

New this week | Family: Betulaceae | Tier A | Priority 1 - core

Dry-mesic and rocky understories throughout most of the region

Shade-tolerant small tree with very hard wood; common understory associate




Bark

(c) Peter Quakenbush, some rights...
INAT_PHOTO_601318538 | cc-by



Foliage

no rights reserved
INAT_PHOTO_724659862 | cc0



Mixed Evidence

no rights reserved
INAT_PHOTO_713918591 | cc0

Evidence	Diagnostic interpretation
Foliage	Alternate ovate leaves have many straight lateral veins, a sharply pointed tip, and a doubly serrate margin; the surface often looks slightly corrugated between veins. Foliage overlaps American hornbeam, birches, and even beech without bark or fruit.
Bark age series	Young stems are smooth gray-brown with lenticels; older small trunks break into narrow, loose, vertical gray-brown strips that shred at the ends, producing the characteristic cat-scratch or shredded appearance. It does not develop hornbeam's smooth muscular fluting.
Dormant	Slender alternate twigs bear short pointed ovoid buds with many visible scales; several cylindrical male catkins persist through winter near twig tips. Buds are far shorter than beech's cigar buds.
Fruit / cone / seed	Female clusters mature into dangling hoplike structures made of overlapping inflated papery sacs, each enclosing a small nutlet. The complete hoplike cluster is decisive within the regional understory flora.
Whole tree / context	A slow-growing small understory tree, often with a narrow crown and slightly leaning bole, common on rocky dry-mesic slopes, ridges, rich woods, and forest edges.
Variation / condition	Saplings lack shredded bark, while very old stems may be rougher and less neatly striped. Lichens, wetness, browse, suppressed form, and mechanical scars can obscure the narrow-strip pattern.
Evidence limit	Hoplike fruit or shredded bark combined with short scaled buds supports species. A detached doubly serrate leaf, smooth sapling bark, or distant small understory bole alone cannot securely separate <i>Ostrya</i> from <i>Carpinus</i> , <i>Betula</i> , or <i>Fagus</i> .

Confuser control

Alternative	Visible separator	Resolving view
<i>Carpinus caroliniana</i>	American hornbeam has smooth gray bark molded into muscle-like flutes and nutlets attached to flat three-lobed leafy bracts; hophornbeam develops narrow shreddy strips and inflated hoplike fruit sacs.	Representative mature bark, dormant buds and catkins, and the entire fruiting cluster.
<i>Fagus grandifolia</i>	Beech has exceptionally long cigar buds, a simpler tooth-per-vein leaf margin, generally smooth host bark, and nuts in prickly husks; hophornbeam has short buds, doubly serrate leaves, shredded bark, and hoplike fruit.	Full twig with buds, leaf margin and venation, bark away from injuries, and fruit.

Silvics capsule

Dimension	Field-use summary
Shade tolerance	Shade tolerant and able to persist as a slow-growing understory tree beneath hardwood and mixedwood canopies.
Moisture / soils	Common on dry-mesic to mesic, well-drained, often rocky or calcareous soils; avoids prolonged flooding and is not restricted to the richest sites.
Geographic affinity	Widespread across eastern North America and nearly the full course region, from Great Lakes forests through New England and the Allegheny transition.

Dimension	Field-use summary
Associations	Occurs beneath sugar maple, basswood, oaks, hickories, beech, yellow birch, hemlock, and mixed upland hardwoods.
Regeneration	Nutlets disperse mainly near parent trees and establish beneath partial canopy; persistent seedlings and saplings maintain the local understory population.
Vegetative reproduction	Cut or damaged young stems can sprout from the stump or root collar, but the species is not normally a strongly clonal stand former.
Seed / mast	Hoplike clusters mature in late summer or autumn; crops vary, and gravity plus animals provide mostly short-distance dispersal.
Establishment	Well-drained forest soil, some shade, and freedom from severe browse allow establishment; very low growth permits persistence until small gaps provide release.
Succession	Persistent mid- to late-seral understory associate rather than a canopy pioneer or major overstory dominant.
Disturbance response	Small gaps release established stems and top-killed young trees may sprout; severe fire damages the thin bark, while heavy overstory removal may expose suppressed crowns abruptly.
Longevity / growth	Slow-growing, relatively long-lived for a small tree, and structurally limited to understory or subcanopy stature; wood is exceptionally dense and hard.
Browse	Deer and other mammals browse twigs and foliage, but the species is generally less central to browse management than hemlock, maple, or birch regeneration.
Insects / disease	Usually not controlled by a single major pest; leaf spots, cankers, borers, and decay occur, especially after injury or stress.
Management	Retain for structural and taxonomic diversity, wildlife value, and dry-mesic understory function unless it directly conflicts with regeneration objectives; do not mistake small size for a juvenile canopy species.
Silvicultural systems	Selection and small-group openings tend to retain and modestly release hophornbeam; broad even-aged treatments may stimulate sprouts but rarely make it a primary crop-tree objective.

Personal field cue for Eastern hophornbeam
learner-created retrieval hook

Part III - Fillable instructional sessions

Complete sessions in order when possible. Closed-book retrieval means close or hide the teaching atlas before answering.

Session A - Six northern-hardwood anchors

Purpose: diagnostic teaching examples plus field-decision modeling. Taxon scope: listed new taxa. Feedback: during instruction.

Before the session: which species or comparison currently feels least reliable?

Sugar maple: one diagnostic character plus one silvics anchor

Red maple: one diagnostic character plus one silvics anchor

American beech: one diagnostic character plus one silvics anchor

Yellow birch: one diagnostic character plus one silvics anchor

Paper birch: one diagnostic character plus one silvics anchor

Eastern hemlock: one diagnostic character plus one silvics anchor

American basswood: one diagnostic character plus one silvics anchor

Black cherry: one diagnostic character plus one silvics anchor

Eastern hophornbeam: one diagnostic character plus one silvics anchor

Exit retrieval: state the most important correction or strengthened distinction

☐ Session A complete (enter X)

Session B - Anchor retrieval

Purpose: closed-response batches of five to ten unfamiliar practice items. Taxon scope: session a taxa. Feedback: after batch submission.

Before the session: which species or comparison currently feels least reliable?

1. Nomenclature

Give the preferred course common name for *Ostrya virginiana*.

Response

closed-book retrieval

2. Forest Association

Place Sugar maple in the course region and in a characteristic forest association. Explain why the context raises or lowers its plausibility without treating habitat as proof of identity.

Response

closed-book retrieval

3. Family Relationship

To which botanical family does Eastern hophornbeam (*Ostrya virginiana*) belong?

Response

closed-book retrieval

4. Forest Association

A field crew reports Eastern hemlock from an unfamiliar stand. State the regional and community context that would make the report credible, then request morphological evidence needed to confirm it.

Response

closed-book retrieval

5. Nomenclature

Give the preferred course common name for *Betula papyrifera*.

Response

closed-book retrieval

6. Silvics

For Red maple (*Acer rubrum*), describe the moisture, drainage, and soil setting a forester should expect, then place it in its climatic or forest-association context.

Response

closed-book retrieval

7. Silvics

Place Red maple in succession: give its shade tolerance, explain how it recruits or persists, and characterize its longevity or growth strategy.

Response

closed-book retrieval

8. Silvics

A stand prescription lists Eastern hemlock as an objective species. State the site conditions that favor it and one site or climatic constraint that would make that objective questionable.

Response

closed-book retrieval

9. Forest Association

Place Eastern hophornbeam in the course region and in a characteristic forest association. Explain why the context raises or lowers its plausibility without treating habitat as proof of identity.

Response

closed-book retrieval

10. Silvics

Predict the role of Paper birch from suppression through canopy recruitment. Tie the prediction to shade tolerance, successional status, and growth or longevity.

Response

closed-book retrieval

Exit retrieval: state the most important correction or strengthened distinction

☐ Session B complete (enter X)

Session C - Birch, cherry, and hophornbeam contrasts

Purpose: direct comparison and isolated unfamiliar specimens. Taxon scope: listed new taxa plus week confusers. Feedback: after response.

Before the session: which species or comparison currently feels least reliable?

Yellow birch / black cherry bark

Course separator: Golden horizontal curls and wintergreen twig versus burnt-chip plates, petiole glands, and almond twig odor

My visible separator

Best resolving view

Exit retrieval: state the most important correction or strengthened distinction

☐ Session C complete (enter X)


Session D - Anchor multi-organ laboratory

Purpose: single-organ and partial-evidence practice stations. Taxon scope: all module taxa. Feedback: after station batch submission.

Before the session: which species or comparison currently feels least reliable?

Practice 1

Identity: common OR scientific name



Evidence asset INAT_PHOTO_651077467 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 2

Identity: common OR scientific name



Evidence asset INAT_PHOTO_720830462 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 3

Identity: common OR scientific name



Evidence asset INAT_PHOTO_722719712 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 4

Identity: common OR scientific name



Evidence asset INAT_PHOTO_724733783 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high


Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 5

Identity: common OR scientific name



Evidence asset INAT_PHOTO_579273288 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 6

Identity: common OR scientific name



Evidence asset INAT_PHOTO_718762908 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 7

Identity: common OR scientific name



Evidence asset INAT_PHOTO_630118290 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high


Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 8

Identity: common OR scientific name



Evidence asset INAT_PHOTO_649013724 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 9

Identity: common OR scientific name



Evidence asset INAT_PHOTO_602315507 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Practice 10

Identity: common OR scientific name



Evidence asset INAT_PHOTO_713516494 | credit in Sources appendix

Only the identity field is universally required. Instructors may score reasoning fields where the rubric calls for a field decision.
Use genus or 'insufficient evidence' when the photograph cannot defend species.

Required - defensible identity

Confidence: low / moderate / high

Nearest plausible alternative

Visible separator / evidence

Best additional view if unresolved

Exit retrieval: state the most important correction or strengthened distinction

☐ Session D complete (enter X)

Session E - Week 1 cumulative retrieval

Purpose: adaptive retrieval and remediation insertion. Taxon scope: cr 01 eligible taxa prioritized by mastery. Feedback: after batch submission.

Before the session: which species or comparison currently feels least reliable?

1. Silvics

Explain how American basswood normally regenerates, including establishment requirements and any seed, mast, sprouting, layering, or suckering behavior that materially affects a prescription.

Response

closed-book retrieval

2. Forest Association

A field crew reports Yellow birch from an unfamiliar stand. State the regional and community context that would make the report credible, then request morphological evidence needed to confirm it.

Response

closed-book retrieval

3. Family Relationship

To which botanical family does American beech (*Fagus grandifolia*) belong?

Response

closed-book retrieval

4. Silvics

Design the species-specific part of a prescription for Sugar maple: state what to protect or control, and justify the cutting or regeneration approach from its known response.

Response

closed-book retrieval

5. Nomenclature

Give the preferred course common name for *Ostrya virginiana*.

Response

closed-book retrieval

6. Silvics

Design the species-specific part of a prescription for Eastern hophornbeam: state what to protect or control, and justify the cutting or regeneration approach from its known response.

Response

closed-book retrieval

7. Silvics

Predict the role of Black cherry from suppression through canopy recruitment. Tie the prediction to shade tolerance, successional status, and growth or longevity.

Response

closed-book retrieval

8. Family Relationship

To which botanical family does Eastern hemlock (*Tsuga canadensis*) belong?

Response

closed-book retrieval

9. Silvics

Explain how Red maple normally regenerates, including establishment requirements and any seed, mast, sprouting, layering, or suckering behavior that materially affects a prescription.

Response

closed-book retrieval

10. Silvics

Place Yellow birch in succession: give its shade tolerance, explain how it recruits or persists, and characterize its longevity or growth strategy.

Response

closed-book retrieval

Exit retrieval: state the most important correction or strengthened distinction

☐ Session E complete (enter X)

Image credits and instructional references

Image credits apply only to the labeled teaching and formative practice plates in this public learner packet. Secure examination specimens and answer-bearing metadata are not included.

Asset	Creator / license	Source
INAT_PHOTO_458152552	(c) Bryce, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/458152552
INAT_PHOTO_462391786	no rights reserved cc0	https://www.inaturalist.org/photos/462391786
INAT_PHOTO_564773858	(c) Quinten Wiegiersma, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/564773858
INAT_PHOTO_574455120	(c) Amy MacPherson, some rights reserved (CC BY-SA) cc-by-sa	https://www.inaturalist.org/photos/574455120
INAT_PHOTO_579273288	(c) Damien Mullin, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/579273288
INAT_PHOTO_580635573	(c) Patrick Jackson, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/580635573
INAT_PHOTO_601318538	(c) Peter Quakenbush, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/601318538
INAT_PHOTO_602315507	no rights reserved cc0	https://www.inaturalist.org/photos/602315507
INAT_PHOTO_602541471	(c) Matt Pelikan, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/602541471
INAT_PHOTO_607898739	(c) Loïc Mathieu, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/607898739
INAT_PHOTO_608142436	(c) Loïc Mathieu, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/608142436
INAT_PHOTO_611772046	(c) Loïc Mathieu, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/611772046
INAT_PHOTO_612188654	(c) daksu, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/612188654
INAT_PHOTO_619989178	(c) Timon T. K., some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/619989178
INAT_PHOTO_630118290	(c) Alan Prather, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/630118290
INAT_PHOTO_631749583	no rights reserved cc0	https://www.inaturalist.org/photos/631749583
INAT_PHOTO_649013724	no rights reserved cc0	https://www.inaturalist.org/photos/649013724
INAT_PHOTO_650441630	(c) Brandon Corder, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/650441630
INAT_PHOTO_650866298	(c) Nick Kleinschmidt, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/650866298
INAT_PHOTO_651077467	no rights reserved cc0	https://www.inaturalist.org/photos/651077467
INAT_PHOTO_651110401	(c) Rebecca Turner, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/651110401
INAT_PHOTO_669709341	no rights reserved cc0	https://www.inaturalist.org/photos/669709341
INAT_PHOTO_699498715	(c) Steven Lamonde, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/699498715
INAT_PHOTO_702302386	(c) Nash Turley, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/702302386
INAT_PHOTO_705760040	(c) Alan Prather, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/705760040
INAT_PHOTO_712523142	(c) Dan MacNeal, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/712523142
INAT_PHOTO_713516494	(c) Ben Redding, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/713516494
INAT_PHOTO_713542533	(c) sofi v, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/713542533
INAT_PHOTO_713743092	(c) Christina S. Baer, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/713743092
INAT_PHOTO_713870593	no rights reserved cc0	https://www.inaturalist.org/photos/713870593
INAT_PHOTO_713918591	no rights reserved cc0	https://www.inaturalist.org/photos/713918591
INAT_PHOTO_717573410	no rights reserved cc0	https://www.inaturalist.org/photos/717573410
INAT_PHOTO_718762908	no rights reserved cc0	https://www.inaturalist.org/photos/718762908
INAT_PHOTO_720830462	no rights reserved cc0	https://www.inaturalist.org/photos/720830462
INAT_PHOTO_722719712	(c) Nate W., some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/722719712
INAT_PHOTO_724659862	no rights reserved cc0	https://www.inaturalist.org/photos/724659862
INAT_PHOTO_724733783	(c) Alan Prather, some rights reserved (CC BY) cc-by	https://www.inaturalist.org/photos/724733783

Species-profile references

Sugar Maple - Silvics of North America: <https://research.fs.usda.gov/silvics/sugar-maple> - Distribution, sites, associates, seed biology, shade tolerance, growth, damaging agents, and management response.

Acer saccharum - sugar maple: <https://gobotany.nativeplanttrust.org/species/acer/saccharum/> - Regional morphology, reproductive structures, habitat, taxonomy, and comparison characters.

Red Maple - Silvics of North America: <https://research.fs.usda.gov/silvics/red-maple> - Range, ecological breadth, seed and sprout reproduction, succession, growth, damaging agents, and management.

Acer rubrum - red maple: <https://gobotany.nativeplanttrust.org/species/acer/rubrum/> - Regional foliage, buds, flowers, fruit, bark, habitat, and comparative morphology.

American Beech - Silvics of North America: <https://research.fs.usda.gov/silvics/american-beech> - Range, site, shade tolerance, seed and root-sucker reproduction, succession, growth, disease, and management.

Fagus grandifolia - American beech: <https://gobotany.nativeplanttrust.org/species/fagus/grandifolia/> - Regional foliage, buds, bark, fruit, habitat, and comparative identification.

Yellow Birch - Silvics of North America: <https://research.fs.usda.gov/silvics/yellow-birch> - Range, sites, seedbed ecology, tolerance, associations, growth, damaging agents, and silvicultural response.

Betula alleghaniensis - yellow birch: <https://gobotany.nativeplanttrust.org/species/betula/alleghaniensis/> - Regional foliage, bark, buds, catkins, habitat, and comparison with related birches.

Paper Birch - Silvics of North America: <https://research.fs.usda.gov/silvics/paper-birch> - Distribution, sites, seed production, establishment, disturbance response, growth, damaging agents, and management.

Betula papyrifera - paper birch: <https://gobotany.nativeplanttrust.org/species/betula/papyrifera/> - Regional leaves, twigs, bark, catkins, fruit scales, habitat, variation, and birch comparisons.

Eastern Hemlock - Silvics of North America: <https://research.fs.usda.gov/silvics/eastern-hemlock> - Distribution, sites, tolerance, seedbeds, associations, longevity, disturbance, damaging agents, and management.

Tsuga canadensis - eastern hemlock: <https://gobotany.nativeplanttrust.org/species/tsuga/canadensis/> - Regional needle, twig, cone, bark, habit, habitat, and comparative identification.

American Basswood - Silvics of North America: <https://research.fs.usda.gov/silvics/american-basswood> - Range, rich-site relationships, reproduction, sprouting, tolerance, growth, damaging agents, and silvicultural response.

Tilia americana - American linden: <https://gobotany.nativeplanttrust.org/species/tilia/americana/> - Regional foliage, buds, bark, bracted fruit, habitat, taxonomy, and comparison characters.

Black Cherry - Silvics of North America: <https://research.fs.usda.gov/silvics/black-cherry> - Range, productive sites, seed-bank ecology, light requirements, growth, associates, damaging agents, and management.

Prunus serotina - black cherry: <https://gobotany.nativeplanttrust.org/species/prunus/serotina/> - Regional leaves, glands, hairs, bark, flowers, fruit, habitat, and *Prunus* comparisons.

Eastern Hophornbeam - Silvics of North America: <https://research.fs.usda.gov/silvics/eastern-hophornbeam> - Distribution, sites, associates, tolerance, regeneration, growth, damaging agents, and management relevance.

Ostrya virginiana - hop-hornbeam: <https://gobotany.nativeplanttrust.org/species/ostrya/virginiana/> - Regional leaves, buds, bark, catkins, fruit, habitat, and comparison with hornbeam and birches.