
TEAS Math Practice Exam

40 Questions

Mu2 Solutions

Time Limit: 57 minutes

Calculator: Permitted

Questions: 40 multiple choice

Format: A, B, C, D

Content: Numbers & Algebra,
Measurement & Data

Instructions

1. Read each question carefully before selecting your answer.
2. Choose the **BEST** answer from the four options provided.
3. Mark your answers on a separate answer sheet.
4. There is no penalty for incorrect answers — answer every question.
5. Pace yourself: approximately 1 minute 25 seconds per question.
6. A calculator may be used on any question.

Mathematics

40 Questions · 57 Minutes · Calculator Permitted

1. A patient drinks 250 milliliters of water. How many liters is this equivalent to?
 - A) 0.25 L
 - B) 0.025 L
 - C) 25 L
 - D) 2.5 L
2. Solve the inequality: $2x - 5 < 9$. Which of the following represents the solution set?
 - A) $x < 7$
 - B) $x < 2$
 - C) $x > 7$
 - D) $x > 2$
3. The pie chart shows department budgets: Research 40%, Marketing 25%, Operations 35%. If the total budget is \$500,000, how much more money is allocated to Research than Marketing?
 - A) \$25,000
 - B) \$50,000
 - C) \$75,000
 - D) \$100,000
4. A patient is prescribed 1.5 Liters of IV fluid to be infused over 10 hours. How many milliliters per hour should the IV pump be set to?
 - A) 15 mL/hr
 - B) 150 mL/hr
 - C) 1,500 mL/hr
 - D) 15,000 mL/hr
5. A square has a side length of 4 inches. What is the area of the square?
 - A) 8 square inches
 - B) 12 square inches
 - C) 16 square inches
 - D) 16 inches
6. A hospital staffing table shows: Nurse A works three 8-hour shifts, and Nurse B works four 6-hour shifts. What is the total number of hours worked by both nurses?
 - A) 32 hours
 - B) 56 hours
 - C) 64 hours
 - D) 48 hours
7. A nurse is preparing a medication solution by mixing 300 mL of saline with 250 mL of water to create a total volume of 550 mL. What is the concentration of the saline solution in percent?
 - A) 1.0 mL
 - B) 1.5 mL
 - C) 1.25 mL
 - D) 2.5 mL
8. What is the area of a square bandage with each side measuring 6 inches?

- A) 18 square inches
- B) 30 square inches
- C) 36 square inches
- D) 48 square inches

9. A nurse is calculating the total dosage of a medication given over three days. The daily dose is 500 mg. How many milligrams are administered in total?

- A) 1250 mg
- B) 2000 mg
- C) 1000 mg
- D) 1500 mg

10. A table shows daily water intake: Monday 4 cups, Tuesday 6 cups, Wednesday 5 cups. What is the total intake?

- A) 13 cups
- B) 14 cups
- C) 15 cups
- D) 16 cups

11. A nurse needs to calculate the total volume of IV fluid a patient received over 8 hours. The patient received 125 mL of saline in the first hour, 175 mL in the second hour, and 250 mL in the third hour. What is the total volume of fluid administered?

- A) 450 mL
- B) 650 mL
- C) 350 mL
- D) 550 mL

12. A nurse is reviewing a chart that shows the number of patients admitted to the emergency room over a 5-day period: 12, 15, 18, 12, and 20. What is the mean number of daily admissions?

- A) 15.4
- B) 14.8
- C) 16.0
- D) 15.0

13. A nurse is preparing to administer a medication. The prescription calls for 0.75 mL of the solution. Which of the following fractions is equivalent to this decimal measurement?

- A) $\frac{7}{10}$ mL
- B) $\frac{3}{4}$ mL
- C) $\frac{1}{4}$ mL
- D) $\frac{3}{10}$ mL

14. A nurse needs to calculate the total volume of medication administered to a patient who received 0.25 liters of saline in the morning and 375 milliliters in the afternoon. What is the total volume in milliliters?

- A) 325 mL
- B) 600 mL

C) 625 mL

D) 650 mL

15. Solve the inequality: $3x - 7 > 8$. Which of the following represents the solution set?

A) $x < 5$

B) $x > 1$

C) $x > 15$

D) $x > 5$

16. A nurse is calculating the total dosage for a patient receiving two different medications. The first medication is given at a rate of $(3x + 5)$ mg, and the second at $(2x - 1)$ mg. If x represents the patient's weight in kilograms, what is the simplified expression for the total dosage in terms of x ?

A) $5x + 6$

B) $5x + 4$

C) $3x + 4$

D) $5x - 4$

17. A nurse is preparing a medication solution that requires mixing 125 mL of saline with 40 mL of a drug concentrate. What is the total volume of the final solution in milliliters?

A) 1650 mL

B) 165 mL

C) 100 mL

D) 121 mL

18. A nurse is calculating the cost of a medication that costs \$15 per unit. If the nurse needs to purchase x units, which algebraic expression represents the total cost?

A) $x / 15$

B) $15 - x$

C) $15x$

D) $15 + x$

19. In a study of 120 patients: normal cholesterol and BP 42, high cholesterol only 30, high BP only 24, both high 24. What percent have at least one high value?

A) 55%

B) 60%

C) 65%

D) 70%

20. Pain scores frequency: Score 0: 2 patients, 1: 3, 2: 5, 3: 4, 4: 1. What is the mean score?

A) 1.8

B) 2.0

C) 2.1

D) 2.3

21. A pie chart represents monthly expenses: Rent 35%, Food 25%, Utilities 20%, Transportation 15%, Other 5%. If total monthly expenses are \$2,400, how much is spent on Food?

A) \$480

B) \$600

C) \\$840

D) \\$360

22. A patient's doctor orders a medication dose of 250 mg. The available tablets are labeled 0.5 g each. How many tablets should the nurse administer?

A) 1 tablet

B) 0.5 tablets

C) 0.75 tablets

D) 0.25 tablets

23. A patient must drink 1 liter of water. They have already drunk 250 mL. How many milliliters are left to drink?

A) 250 mL

B) 500 mL

C) 750 mL

D) 850 mL

24. Medication errors per 1000 doses by quarter: Q1 4.5, Q2 3.6, Q3 2.7, Q4 1.8. What is the average decrease per quarter?

A) 0.7

B) 0.8

C) 0.9

D) 1.0

25. A nurse needs to calculate the total volume of fluid a patient will receive over the next 4 hours. The IV pump is set to deliver 75 mL/hour. What is the total volume?

A) 250 mL

B) 375 mL

C) 3000 mL

D) 300 mL

26. A nurse needs to administer a medication of 2.5 mL to a patient. The patient's stomach can only hold a maximum of 10 mL at a time. How much additional volume can be added to this single dose without exceeding the stomach's capacity?

A) 5 mL

B) 7.5 mL

C) 8 mL

D) 12.5 mL

27. A nurse is calculating the total amount of medication in a mixture. There are 3 bottles of 50mg pills and 2 bottles of 50mg pills. Which property of operations best justifies that the order in which she adds these quantities does not change the final result?

A) commutative Property

B) Associative Property

C) Distributive Property

D) Identity Property

28. A recipe for a large batch of punch requires 6 liters of water and 2 liters of fruit juice. If you want to make a smaller batch using only 3 liters of water, how many liters of fruit juice will you need?

- A) 0.5 liters of fruit juice
- B) 1 liter of fruit juice
- C) 2 liters of fruit juice
- D) 3 liters of fruit juice

29. What is the area of a triangular wound dressing with a base of 10 cm and a height of 6 cm?

- A) 30 cm²
- B) 60 cm²
- C) 15 cm²
- D) 40 cm²

30. A nurse is calculating the dosage of a medication for a patient. The formula is $D = 500 * (\frac{1}{4}) + 100$.

- A) 225 mg
- B) 150 mg
- C) 125 mg
- D) 250 mg

31. A doctor prescribes a medication dose of 250 mg. If the available formulation is 5 mg/mL, how many milliliters should be administered?

- A) 125 mL
- B) 100 mL
- C) 5 mL
- D) 50 mL

32. A nurse needs to prepare a medication dosage. The order states 150 mg of medication per dose, and the available concentration is 250 mg per 5 mL. How many milliliters should the nurse administer for one dose?

- A) 2.5 mL
- B) 3 mL
- C) 5 mL
- D) 4 mL

33. A triangular bandage has a base of 10 cm and height of 6 cm. What is its area?

- A) 30 cm²
- B) 40 cm²
- C) 60 cm²
- D) 16 cm²

34. Two hundred nurses were surveyed on shift preference: Day 70, Night 50, Rotate 80. There are 100 nurses under age 40, and 42 of them prefer Day. What percent of under-40 nurses prefer Day?

- A) 35%
- B) 42%
- C) 50%
- D) 60%

35. A rectangular tray is 18 inches by 12 inches. What is its perimeter in feet?

- A) 2.5 ft
- B) 5 ft

- C) 6 ft
- D) 7.5 ft

36. A nurse is preparing a solution where the ratio of salt to water is 1 part salt to 4 parts water. If the nurse mixes 20 mL of salt with 80 mL of water, what is the total volume of the solution?

- A) 100 mL
- B) 20 mL
- C) 80 mL
- D) 90 mL

37. A nurse is preparing to administer medication based on a specific ratio. If the required ratio of drug to diluent is 1:5, how many milliliters of diluent are needed to mix with 10 mL of the drug?

- A) 10 mL
- B) 50 mL
- C) 15 mL
- D) 5 mL

38. A histogram displays patient weights in intervals: 100-110 lbs (2 patients), 110-120 lbs (5 patients), 120-130 lbs (8 patients), 130-140 lbs (3 patients). In which weight interval does the median patient fall?

- A) 110-120 lbs
- B) 100-110 lbs
- C) 120-130 lbs
- D) 130-140 lbs

39. A nurse is preparing a medication solution and needs to calculate the total volume required. The order specifies adding 250 mL of saline to 125 mL of another solution. How many milliliters of solution are needed in total?

- A) 50 mL
- B) 37.5 mL
- C) 125 mL
- D) 375 mL

40. What percent of 125 is 25?

- A) 20%
- B) 21%
- C) 25%
- D) 10%

Answer Key

- | | |
|-------|-------|
| 1. B | 21. B |
| 2. A | 22. B |
| 3. C | 23. C |
| 4. B | 24. C |
| 5. C | 25. D |
| 6. C | 26. B |
| 7. C | 27. A |
| 8. C | 28. B |
| 9. D | 29. B |
| 10. C | 30. A |
| 11. D | 31. D |
| 12. A | 32. B |
| 13. B | 33. A |
| 14. C | 34. B |
| 15. D | 35. B |
| 16. B | 36. A |
| 17. B | 37. B |
| 18. C | 38. C |
| 19. C | 39. D |
| 20. B | 40. A |
-

Detailed Explanations

Q1. Answer: B

The question asks to convert 250 milliliters to liters. Since there are 1,000 milliliters in one liter, dividing 250 by 1,000 yields 0.25 liters, which matches option B. Option A is incorrect because it likely suggests 250 liters, ignoring the decimal shift needed for unit conversion. Option C is wrong as it may propose 2.5 liters, mistakenly moving the decimal one place too far. Option D is invalid because it could state 0.025 liters, which represents an extra decimal place error.

Q2. Answer: A

To solve $2x - 5 < 9$, add 5 to both sides getting $2x < 14$, then divide by 2 to find $x < 7$. Option A matches this result exactly while Option B incorrectly subtracts 5 from 9 instead of isolating the term. Option C reverses the inequality sign and Option D also flips the sign and miscalculates the value. Only A represents the true solution set derived from standard algebraic steps.

Q3. Answer: C

The difference between Research and Marketing is 15% (40% minus 25%). Calculating 15% of the \$500,000 total budget yields exactly \$75,000, making option C the right choice. Option A is wrong because it only reflects the 5% gap between Operations and Research. Option B incorrectly assumes a 10% difference, while option D mistakenly calculates a 20% gap.

Q4. Answer: B

To find the hourly rate, convert 1.5 Liters to 1,500 milliliters, then divide by 10 hours to get 150 mL/hr. Option B matches this calculation exactly. Option A is too low because it divides 150 by 10 again. Option C is incorrect as it uses the total volume without dividing by time. Option D is wrong because it multiplies the volume by 10

instead of dividing.

Q5. Answer: C

The area of a square equals side length squared, so 4 times 4 is 16 square inches. Option C correctly states this value with proper units. Option A is wrong because 8 is half of 16, not the area. Option B is incorrect as 12 does not result from squaring 4. Option D matches the number but lacks the required "square inches" unit.

Q6. Answer: C

To find the total hours, calculate Nurse A's 24 hours (3 times 8) plus Nurse B's 24 hours (4 times 6), resulting in 48 total. Option C correctly states this sum. Option A is wrong because it likely ignores one nurse's shifts. Option B is incorrect as it probably adds the shift counts instead of hours. Option D fails by miscalculating the multiplication for one of the nurses.

Q7. Answer: C

The question tests number sense and unit conversion; since 1 teaspoon equals 5 mL, one-fourth of a teaspoon is calculated by dividing 5 by 4, which results in 1.25 mL. The other options represent common errors such as misplacing the decimal point or incorrectly halving the value.

Q8. Answer: C

The area of a square is side length squared, so 6 inches times 6 inches equals 36 square inches. Option C states 36, which matches this calculation. Option A suggests 12, which is the perimeter, not the area. Option B proposes 18, half the correct value, likely a division error. Option D gives 72, double the area, perhaps from multiplying side by perimeter.

Q9. Answer: D

To find the total dosage, multiply the daily amount of 500 mg by the three days, resulting in 1500 mg. Option A is correct because 500 times 3 equals 1500. Option B is wrong as 1250 does not match the calculation. Option C is incorrect since 1000 is too low. Option D is wrong because 2000 exceeds the total. This math problem requires simple multiplication.

Q10. Answer: C

To find the total intake, add Monday's 4 cups, Tuesday's 6 cups, and Wednesday's 5 cups. This sum equals 15 cups, making option C the correct answer. Option A is wrong because 13 is less than the actual total. Option B is incorrect as 14 does not match the calculated sum. Option D is wrong since 16 exceeds the true total of 15 cups.

Q11. Answer: D

To find the total volume, add the amounts from the first three hours: 125 mL plus 175 mL plus 250 mL equals 550 mL. This matches the correct answer because it sums all administered fluid. The 350 mL and 450 mL options are too low, likely missing the third hour's 250 mL. The 650 mL option is too high, perhaps adding an extra 100 mL incorrectly.

Q12. Answer: A

To calculate the mean, first sum the five daily admission counts: $12 + 15 + 18 + 12 + 20$ equals 77. Next, divide this total by the number of days, which is 5. The result is 77 divided by 5, which equals 15.4. Option B (14.8) is incorrect because it likely stems from an improper division or summing error. Option C (16.0) is wrong and may result from a calculation mistake or misreading the data. Option D (15.0) represents the median value, not the mean. The mean is the arithmetic average, which...

Q13. Answer: B

The fraction $\frac{3}{4}$ is equivalent to the decimal 0.75 because dividing 3 by 4 yields exactly 0.75, matching the prescribed volume. To verify, we can convert each option: $\frac{7}{10}$ equals 0.70, which is 0.05 less than required; $\frac{1}{4}$ equals 0.25, which is significantly smaller and represents a quarter of the needed amount; $\frac{3}{10}$ equals 0.30, which is more than double the required volume. Therefore, only $\frac{3}{4}$ mL accurately represents the nurse's dosage. It is crucial for nurses to understand decimal-fraction...

Q14. Answer: C

First, convert 0.25 liters to milliliters by multiplying by 1,000, which equals 250 mL. Then, add the morning dose (250 mL) to the afternoon dose (375 mL) to get a total of 625 mL. The other options result from common errors such as incorrect conversion or arithmetic mistakes.

Q15. Answer: D

To solve $3x - 7 > 8$, add 7 to both sides getting $3x > 15$, then divide by 3 to find $x > 5$. Option A is right because it matches this exact result. Option B is wrong as adding 7 to 8 incorrectly yields 15. Option C fails because dividing 15 by 3 does not give 5. Option D is incorrect since the inequality direction must stay greater than.

Q16. Answer: B

To find the total dosage, add the two expressions $(3x + 5)$ and $(2x - 1)$ by combining like terms. The x terms sum to $5x$ while the constants 5 and -1 result in 4, giving the correct answer $5x + 4$. Option A $(3x + 4)$ fails because it incorrectly adds only the x coefficients. Option B $(5x + 6)$ is wrong as it mistakenly adds 5 and 1 instead of subtracting. Option C $(5x - 4)$ is incorrect because it subtracts the constants rather than adding them. This math problem requires simple algebraic addition ...

Q17. Answer: B

To find the total volume, simply add the 125 mL of saline and the 40 mL of drug concentrate together. The sum is 165 mL, which matches the correct answer. The option 100 mL is wrong because it subtracts the volumes instead of adding them. The choice 1650 mL is incorrect as it mistakenly adds a zero to the true sum. Finally, 121 mL is wrong because it appears to be the result of subtracting 40 from 125.

Q18. Answer: C

The total cost is found by multiplying the price per unit ($\$15$) by the number of units (x), resulting in the expression $15x$. The other options represent addition, division, or subtraction, which do not correctly calculate total cost based on the given scenario.

Q19. Answer: C

To find the percent with at least one high value, add patients with high cholesterol only (30), high BP only (24), and both high (24) for a total of 78 out of 120. Dividing 78 by 120 gives 0.65, which equals 65%, making option C correct. Options A, B, and D are incorrect because they do not match the calculated sum of 78 patients. The question asks for the percentage with at least one high value, not just one specific condition.

Q20. Answer: B

To find the mean, multiply each score by its frequency, sum the results ($0*2 + 1*3 + 2*5 + 3*4 + 4*1 = 27$), and divide by the total patients (15), yielding 2.0. Option B is correct because this calculation matches the data provided. Option A is wrong as 1.8 underestimates the weighted average. Option C is incorrect since 2.1 overestimates the result. Option D is false because 2.3 is too high for this dataset.

Q21. Answer: B

To find the Food expense, multiply the total $\$2,400$ by the 25% Food portion, which equals $\$600$. Option B is correct because it matches this calculation. Option A is wrong as it uses 20% for Utilities instead of 25%. Option C is incorrect because it applies 35% for Rent to the total. Option D is wrong since it divides the total by 6.5 rather than multiplying by the percentage.

Q22. Answer: B

The correct answer is 0.5 because the nurse must convert grams to milligrams (250 mg is half of 500 mg) or recognize that 0.5 g equals 500 mg, meaning the patient needs half of one tablet. The other options result from common calculation errors or misreading the dosage.

Q23. Answer: C

To find the remaining water, subtract the 250 mL already consumed from the total 1 liter (1000 mL), which equals 750 mL. Option C matches this calculation exactly. Option A is incorrect because it represents the amount already drunk, not the remainder. Option B is wrong as it assumes only half a liter was needed initially. Option D is inaccurate because it adds the consumed amount to the total instead of subtracting.

Q24. Answer: C

To find the average decrease, subtract consecutive quarters: Q1 to Q2 is 0.9, Q2 to Q3 is 0.9, and Q3 to Q4 is 0.9. Since the drop is constant at 0.9 per quarter, option C is correct. Options A, B, and D are incorrect because they do not match the calculated difference of 0.9 derived from the data. The consistent pattern confirms C as the only valid answer for the average decrease.

Q25. Answer: D

To find the total volume, multiply the hourly rate (75 mL) by the number of hours (4), which equals 300 mL. The distractors represent common errors such as dividing instead of multiplying ($25 \text{ mL/hour} * 4 = 100$, not

listed, so 250 is likely a mental math error or misreading 75 as $60+90$), multiplying by the wrong time ($75 * 5 = 375$), and a decimal placement error (3000).

Q26. Answer: B

To determine the additional volume that can be safely added, we must first calculate the remaining capacity of the stomach after the initial dose is administered. The stomach's maximum capacity is 10 mL, and the nurse is administering 2.5 mL. By subtracting the administered volume from the total capacity ($10 \text{ mL} - 2.5 \text{ mL}$), we find that there is exactly 7.5 mL of space remaining. Therefore, adding 7.5 mL will fill the stomach to its limit without exceeding it. Any amount greater than 7.5 mL, su...

Q27. Answer: A

The Commutative Property states that changing the order of addends ($a + b = b + a$) does not change the sum. In this scenario, adding the quantity from the first group of bottles to the second group yields the same total as adding them in the reverse order.

Q28. Answer: B

The recipe ratio is 3 liters water to 1 liter juice, so halving the water to 3 liters requires halving the juice to 1 liter. Option A (2 liters) ignores the proportional reduction needed for the smaller batch. Option B (3 liters) assumes equal amounts regardless of the water change. Option C (0.5 liters) incorrectly calculates the ratio instead of applying the direct proportion.

Q29. Answer: B

The area of a triangle equals one-half the base times the height. Substituting the given values, one-half of ten centimeters times six centimeters yields thirty square centimeters. Option B correctly states this result. Options A, C, and D are incorrect because they fail to apply the triangle area formula properly or calculate the wrong value.

Q30. Answer: A

To find the dosage, calculate 500 times one-fourth, which equals 125, then add 100 to get 225 mg. This matches the correct answer of 225 mg. The wrong answer of 125 mg ignores the final addition of 100. The option 150 mg results from incorrectly adding 25 instead of multiplying. The choice 250 mg comes from multiplying 500 by one-fourth and then doubling the result.

Q31. Answer: D

To determine the volume to administer, divide the prescribed dose by the concentration of the medication. The calculation is 250 mg divided by 5 mg/mL, which equals 50 mL. This result is represented by option D. Option A (125 mL) is a common error that occurs when a student mistakenly multiplies the dose by 0.5 instead of dividing by the concentration. Option B (100 mL) might result from dividing the dose by 2.5 instead of the correct concentration of 5. Option C (5 mL) is incorrect because i...

Q32. Answer: B

To find the volume, set up a proportion where 250 mg corresponds to 5 mL and solve for the volume of 150 mg: $(150/250) * 5 = 3 \text{ mL}$. The other options result from calculation errors, such as dividing 150 by 250 and forgetting to multiply by the original volume (2.5 mL) or incorrectly multiplying the numbers (4 mL or 5 mL).

Q33. Answer: A

To find the area of the triangular bandage, multiply the base of 10 cm by the height of 6 cm and divide by 2, resulting in 30 cm². Option A is correct because it matches this calculation. Option B is wrong as it ignores dividing by 2. Option C is incorrect because it multiplies base and height without dividing. Option D is wrong as it adds the dimensions instead of multiplying them.

Q34. Answer: B

The question asks for the percentage of under-40 nurses who prefer the Day shift, not the total surveyed group. Since 42 out of 100 nurses under 40 prefer Day, the calculation is simply 42 divided by 100, which equals 42%. Option A is incorrect because 35% would imply 35 nurses, ignoring the actual count of 42. Option C is wrong as it uses the total number of nurses preferring Day without dividing by the under-40 subgroup. Option D is invalid because 60% does not match the ratio of 42 to 100....

Q35. Answer: B

To find the perimeter, use the formula $P = 2(\text{length} + \text{width})$. Substituting the given values: $P = 2(18 + 12) = 2(30) = 60$ inches. Since the question asks for the perimeter in feet, convert inches to feet by dividing by 12:

$60 \div 12 = 5$ feet, which corresponds to option B. Option A (2.5 ft) is incorrect because it equals 30 inches, representing only one length plus one width rather than the full perimeter. Options C (6 ft) and D (7.5 ft) are incorrect because they result in 72 inches and 90 inch...

Q36. Answer: A

The total volume is 100 mL because adding 20 mL of salt and 80 mL of water equals 100 mL. The 20 mL and 80 mL options are wrong as they represent only individual components, not the sum. The 90 mL choice is incorrect since the volumes add directly without loss in this scenario. This math question requires simple addition of the given quantities to find the final solution volume.

Q37. Answer: B

The ratio 1:5 means 1 part drug requires 5 parts diluent. Since there are 10 mL of drug, multiplying by 5 gives 50 mL of diluent needed. Option B correctly calculates this product. Option A mistakenly uses the drug volume as the diluent amount. Option C incorrectly adds the two volumes together. Option D reverses the ratio logic entirely.

Q38. Answer: C

First, find the total number of patients: $2 + 5 + 8 + 3 = 18$. The median is the average of the 9th and 10th patients when ordered. The first 7 patients fall in the 100-120 lbs range. Patients 8 through 15 fall in the 120-130 lbs range. Since both the 9th and 10th patients are in this group, the median interval is 120-130 lbs.

Q39. Answer: D

To find the total volume, simply add the 250 mL of saline and the 125 mL of the other solution together, which equals 375 mL. The 125 mL option is wrong because it only counts one part of the mixture. The 37.5 mL and 50 mL choices are incorrect as they do not represent the sum of the two specified volumes. Therefore, 375 mL is the only mathematically accurate total for this nurse's order.

Q40. Answer: A

To find the percentage, divide 25 by 125 to get 0.2, then multiply by 100 to obtain 20%. This calculation directly yields option A as the only accurate result. Option B suggests 21%, which is an overestimation resulting from a calculation error. Option C implies 10%, which is half the correct value and fails to reflect the actual ratio. Option D states 25%, incorrectly assuming the parts are equal when they are not. Therefore, only the first choice accurately represents the mathematical relat...