

0.1 Homework 4

0.1.1 Mean-field Solutions for Extended Hubbard Model

The Hamiltonian of the extended Hubbard model can be written as:

$$\hat{H} = -t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow} + V \sum_{\langle i,j \rangle} n_i n_j$$

where:

- $c_{i\sigma}^\dagger$ and $c_{i\sigma}$ are the fermionic creation and annihilation operators for an electron with spin σ at site i .
- $n_{i\sigma} = c_{i\sigma}^\dagger c_{i\sigma}$ is the number operator for electrons with spin σ at site i .
- $n_i = \sum_\sigma c_{i\sigma}^\dagger c_{i\sigma}$ is the number operator for total electrons at site i .
- $U > 0$ is the strength of the on-site interaction between electrons.
- $V > 0$ is the strength of the interaction between electrons at neighboring sites.
- $t > 0$ is the hopping strength of the electrons.

We consider the case of half-filling for two lattice sites ($\langle N \rangle = \langle n_{1\uparrow} + n_{1\downarrow} + n_{2\uparrow} + n_{2\downarrow} \rangle$). In the mean-field approximation, calculate the ground state energy E_{MF} . Please consider initial mean-field values with following four cases.

In the mean-field approximation, the Hamiltonian can be written as

$$\begin{aligned} \hat{H} &= -t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow} + V \sum_{\langle i,j \rangle} n_i n_j \\ &= -t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c.}) + U \sum_i (n_{i\uparrow} \langle n_{i\downarrow} \rangle + n_{i\downarrow} \langle n_{i\uparrow} \rangle - \langle n_{i\uparrow} \rangle \langle n_{i\downarrow} \rangle) \\ &\quad + V \sum_{\langle i,j \rangle} (n_i \langle n_j \rangle + n_j \langle n_i \rangle - \langle n_i \rangle \langle n_j \rangle) \\ &= c^\dagger \begin{bmatrix} U \langle n_{1\downarrow} \rangle + V \langle n_2 \rangle & -t & & \\ -t & U \langle n_{1\uparrow} \rangle + V \langle n_2 \rangle & & \\ & -t & U \langle n_{2\downarrow} \rangle + V \langle n_1 \rangle & -t \\ & & U \langle n_{2\uparrow} \rangle + V \langle n_1 \rangle & \end{bmatrix} c \end{aligned}$$

1. Case 1: Paramagnetic(PM). Initial mean-field value $\langle n_{i\sigma} \rangle = \frac{1}{2}$.

For this case, the interactions are weak, so we expect that the hopping term is dominant. Thus we have

$$\langle n_{i\uparrow} \rangle = \langle n_{i\downarrow} \rangle = \frac{1}{2}, \quad \text{for all } i.$$

$$\begin{bmatrix} U \frac{1}{2} + V & & -t & \\ & U \frac{1}{2} + V & & -t \\ -t & & U \frac{1}{2} + V & \\ & -t & & U \frac{1}{2} + V \end{bmatrix} = U D U^{-1}$$

Except for the different diagonal elements, this matrix is very similar to the case in the lecture. We can get

$$U = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & & -1 \\ 1 & 1 & \\ & 1 & 1 \end{bmatrix}, \quad D = \begin{bmatrix} -t + \frac{U}{2} + V & & & \\ & -t + \frac{U}{2} + V & & \\ & & t + \frac{U}{2} + V & \\ & & & t + \frac{U}{2} + V \end{bmatrix}$$

$$E_{\text{MF}} = -2t + \frac{U}{2} + V$$

2. Case 2: Ferromagnetic(FM). Initial mean-field value $\langle n_{i\uparrow} \rangle = 1$ and $\langle n_{i\downarrow} \rangle = 0$.

When U is large, we expect no double occupancy. For this case, the mean-field values are chosen as

$$\langle n_{1\uparrow} \rangle = \langle n_{2\uparrow} \rangle = 1, \quad \langle n_{1\downarrow} \rangle = \langle n_{2\downarrow} \rangle = 0.$$

$$\begin{bmatrix} V & & -t & \\ & U+V & & -t \\ -t & & V & \\ & -t & & U+V \end{bmatrix} = \begin{bmatrix} & -t & & \\ -t & U & & -t \\ & -t & & U \end{bmatrix} + V\mathbb{I} = UDU^{-1}$$

The effect of V is still just shifting the energy, and we get

$$U = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & & \\ & 1 & 1 & -1 \\ 1 & & 1 & \\ & & 1 & 1 \end{bmatrix}, \quad D = \begin{bmatrix} -t+V & & & \\ & t+V & & \\ & & -t+U+V & \\ & & & t+U+V \end{bmatrix}$$

(a) When $-t+U+V < t+V \iff U < 2t$,

$$\langle n_{1\uparrow} \rangle = \sum_{ij} V_{1i}^* V_{1j} \langle \gamma_i^\dagger \gamma_j \rangle = V_{11}^* V_{11} + V_{13}^* V_{13} = \frac{1}{2}$$

$$\langle n_{1\uparrow} \rangle = \langle n_{2\uparrow} \rangle = \langle n_{1\downarrow} \rangle = \langle n_{2\downarrow} \rangle = \frac{1}{2}$$

which implies the system is still in PM phase and $E_{\text{MF}} = -2t + \frac{U}{2} + V$.

(b) When $U > 2t$,

$$\langle n_{1\uparrow} \rangle = \sum_{ij} V_{1i}^* V_{1j} \langle \gamma_i^\dagger \gamma_j \rangle = V_{11}^* V_{11} + V_{12}^* V_{12} = 1$$

$$\langle n_{1\uparrow} \rangle = \langle n_{2\uparrow} \rangle = 1, \quad \langle n_{1\downarrow} \rangle = \langle n_{2\downarrow} \rangle = 0$$

Now the system is in FM phase and $E_{\text{FM}} = V$.

3. Case 3: Anti-ferromagnetic(AFM). Initial mean-field value $\langle n_{1\uparrow} \rangle = \langle n_{2\downarrow} \rangle = 1 - \alpha$ and $\langle n_{1\downarrow} \rangle = \langle n_{2\uparrow} \rangle = \alpha$.

Another choice when U is large is to give

$$\langle n_{1\uparrow} \rangle = \langle n_{2\downarrow} \rangle = 1 - \alpha, \quad \langle n_{1\downarrow} \rangle = \langle n_{2\uparrow} \rangle = \alpha.$$

$$\begin{bmatrix} \alpha U + V & & -t & \\ & (1-\alpha)U + V & & -t \\ -t & & (1-\alpha)U + V & \\ & -t & & \alpha U + V \end{bmatrix} = \begin{bmatrix} & -t & & \\ -t & (1-2\alpha)U & & -t \\ & -t & (1-2\alpha)U & \\ -t & & -t & \end{bmatrix} + (\alpha U + V)\mathbb{I} = UDU^{-1}$$

The effect of $\bar{V} = \alpha U + V$ is still just shifting the energy. Similar to the contents in the lecture note, mark $\bar{U} = (1-2\alpha)U$ and shift each eigenenergy with \bar{V} , we get

$$\begin{aligned} E_{\text{MF}} &= \bar{U} - \sqrt{4t^2 + \bar{U}^2} + 2\alpha U + 2V - 2\alpha(1-\alpha)U - V \\ &= (1-2\alpha+2\alpha^2)U - \sqrt{4t^2 + \bar{U}^2} + V \end{aligned}$$

and the self-consistent equation is

$$\alpha = \frac{4t^2}{4t^2 + [\sqrt{4t^2 + (1-2\alpha)U^2} + (1-2\alpha)U]^2}$$

- (a) When $U \gg t$, we get $\alpha \approx 0$ and $E_{\text{MF}} \approx -\frac{4t^2}{U} + V$. This corresponds to an AFM solution, which is lower than FM.
- (b) When $U \ll t$, we get $\alpha \approx \frac{1}{2}$ and back to the PM solution.

4. Case 4: Charge density wave(CDW). Initial mean-field value $\langle n_{1\uparrow} \rangle = \langle n_{1\downarrow} \rangle = 1 - \alpha$ and $\langle n_{2\uparrow} \rangle = \langle n_{2\downarrow} \rangle = \alpha$.

When V is much stronger, we expect a double occupancy will occur. Thus the mean-field values are chosen as

$$\langle n_{1\uparrow} \rangle = \langle n_{1\downarrow} \rangle = 1 - \alpha, \quad \langle n_{2\uparrow} \rangle = \langle n_{2\downarrow} \rangle = \alpha.$$

$$\begin{bmatrix} (1-\alpha)U + 2\alpha V & -t & -t \\ -t & (1-\alpha)U + 2\alpha V & -t \\ -t & -t & \alpha U + 2(1-\alpha)V \end{bmatrix} = UDU^{-1}$$

The result is a little complicated and one can solve the matrix by Mathematica easily. Note $\beta = (1 - 2\alpha)(U - 2V)$ and $\gamma = 2t$, we have

$$D = \frac{1}{2} \left((U + 2V)\mathbb{I} + \sqrt{\beta^2 + \gamma^2} \begin{bmatrix} -1 & & & \\ & -1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix} \right)$$

The self-consistent equation is

$$1 - \alpha = \frac{2\beta^2 + \gamma^2 - 2\beta\sqrt{\beta^2 + \gamma^2}}{2\beta^2 + 2\gamma^2 - 2\beta\sqrt{\beta^2 + \gamma^2}}$$

- (a) When $\beta^2 \gg \gamma^2 \iff V \gg \frac{U}{2}$ and $V \gg t$, we have

$$\alpha \approx 0, \quad \langle n_{1\sigma} \rangle = 1, \quad \langle n_{2\sigma} \rangle = 0;$$

$$H_{\text{MF}} \approx U.$$

- (b) When $\beta^2 \ll \gamma^2 \iff V \ll t$ and $U \ll t$, we have $\langle n_{i\sigma} \rangle = \frac{1}{2}$ which corresponds to the PM solution.