
Supplementary Figure S4 Where the ligand sits, pose by pose
4 Å catalytic-placement criterion
6D4O_A
6CXS_A
3K4D_A
P05804_uidA_gus_w214-603
MGYP000574654335
MGYP000751766417
MGYP000497510748
MGYP001319153113
MGYP000016927400
MGYP000968177834
MGYP001050749229
MGYP000051719573
MGYP000729808394
MGYP001178215973
MGYP000245278657 (human faeces)
MGYP000574843407
MGYP001533818318 (human faeces)
P0ABF6_cdd_ecoli
193L_A
1TIM_A
3K4D_A_offsite
93% of 27 poses
89% of 27 poses
67% of 27 poses
80% of 25 poses
56% of 27 poses
52% of 27 poses
42% of 26 poses
59% of 27 poses
52% of 27 poses
19% of 27 poses
30% of 27 poses
33% of 27 poses
35% of 26 poses
0% of 27 poses
0% of 26 poses
0% of 26 poses
0% of 27 poses
22% of 27 poses
15% of 27 poses
4% of 27 poses
0% of 26 poses
0
10
20
30
40
distance from the glucuronide's sugar to the catalytic carboxylates (Å)
positive control (crystal)
positive control (predicted)
shortlisted candidate
unrelated (predicted)
unrelated (crystal)
off-site box
open symbols: sampled from human faeces