



Mercedes College

PHYSICS

YEAR 12

UNIT 3

Name: _____

Teacher: _____

TIME ALLOWED FOR THIS PAPER

Reading time before commencing work: Ten minutes

Working time for the paper: Three hours

MATERIALS REQUIRED/RECOMMENDED FOR THIS PAPER

To be provided by the supervisor:

- This Question/Answer Booklet; ATAR Physics Formulae and Data Booklet

To be provided by the candidate:

- Standard items: pens, pencils, eraser or correction fluid, ruler, highlighter.
- Special items: Calculators satisfying the conditions set by the SCSA for this subject.

IMPORTANT NOTE TO CANDIDATES

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Suggested working time (minutes)	Marks available	Percentage of exam
Section One: Short answer	12	12	50	54	30
Section Two: Extended answer	6	6	90	85	50
Section Three: Comprehension and data analysis	2	2	40	36	20
			Total	175	100

Instructions to candidates

- The rules for the conduct of Western Australian external examinations are detailed in the *Year 12 Information Handbook 2018*. Sitting this examination implies that you agree to abide by these rules.
- Write answers in this Question/Answer Booklet.
- When calculating numerical answers, show your working or reasoning clearly. Give final answers to **three** significant figures and include appropriate units where applicable.

When estimating numerical answers, show your working or reasoning clearly. Give final answers to a maximum of **two** significant figures and include appropriate units where applicable.
- You must be careful to confine your responses to the specific questions asked and follow any instructions that are specific to a particular question.
- Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
 - Planning: If you use the spare pages for planning, indicate this clearly.
 - Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Refer to the question(s) where you are continuing your work.

Section One: Short response**30% (54 marks)**

This section has **twelve (12)** questions. Answer **all** questions. Write your answers in the space provided.

When calculating numerical answers, show your working or reasoning clearly.

Give final answers to three significant figures and include appropriate units where applicable.

When estimating numerical answers, show your working or reasoning clearly. Give final answers to a maximum of two significant figures and include appropriate units where applicable.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page

Suggested working time for this section is 50 minutes.

Question 1

The table below shows some data for two planets orbiting a distant star in another galaxy.

Planets	Mass (kg)	Orbital radius (m)	Radius of planet (m)	Length of one day (s)	Orbital period (s)
Alpha	1.15×10^{25}	4.50×10^{11}	7.9×10^6	9.6×10^4	8.50×10^7
Beta	1.60×10^{24}	9.00×10^{11}	3.8×10^6	4.8×10^4	-


Calculate the value for the orbital period of Beta using the appropriate data in the table.

(5 marks)

Question 2

A student conducted an experiment to illustrate Lenz's Law by dropping identical bar magnets (North at the bottom) through three different hollow tubes. All the tubes were of the **same length (1.00 m) and thickness (0.50 cm)**; however, they were made of three different materials: **aluminium; copper; and PVC plastic**.

The diagram below right illustrates the experiment.



The student knew that **PVC plastic is an excellent insulator**. They also found that the **electrical conductivity (σ)** (measured in Siemens per metre (Sm^{-1})) for **aluminium is $3.766 \times 10^7 \text{ Sm}^{-1}$** ; and for **copper it is $5.977 \times 10^7 \text{ Sm}^{-1}$** . They knew that the **higher ' σ '** for a material, the **better** the conductor.


In terms of the **time** it takes for the magnets to fall through each tube, rank them (aluminium, copper, PVC plastic) **from fastest to slowest**. Place your answers in the table below.

Fastest	
Second fastest	
Slowest	

(3 marks)

Question 3


A uniform concrete block is 1.70 m tall, 0.80 m wide and 0.90 m deep; it has a mass of 900 kg. A minimum horizontal force of 3500 N is required to start sliding the block across the ground.



- (a) Calculate the maximum height 'h' at which the 3500 N horizontal force could be applied without tipping the block over. (3 marks)
- (b) If the ground became more slippery due to rain falling on it, does the block become harder or easier to tip over. Explain briefly. (2 marks)

Question 4

A magnet is free to spin. When the switch is closed, the magnet spins in the direction shown.



- a) Is the end of the **solenoid** marked as 1 a north or south pole? _____ (2 marks)
Give a reason for your answer.


- b) Is the battery terminal marked as 2 positive or negative? _____ (2 marks)
Justify your answer.

- c) Describe the purpose and effect of the soft iron core.

(1 mark)

Question 5

Some students carried out an experiment to investigate the forces required to move an object up a plane. Their equipment was arranged as follows:



The object has a mass of 1.50 kg; the plane's angle of incline (θ) is set at 25° . The students either held the object at rest or moved it along the plane by pulling or releasing the force probe.

When the object is at rest on the inclined plane, the static frictional force between them can be considered to be negligible. However, when the object is in motion the dynamic frictional force can be considered to be 10.0 N.

Calculate the force registered on the force probe in each of the following situations (show working):

- (a) The object is at rest.

(2 marks)

- (b) The object is moving at a constant velocity of 2 ms^{-1} up the plane's surface.

(2 marks)

- (c) The object is accelerating at a rate of 2 ms^{-2} up the plane's surface.

(2 marks)

Question 6

The diagram below shows two horizontal ropes **supporting** a heavy load **above** the ground.




Is it possible to support a heavy load in this way? **Explain your answer.**

(4 marks)

Question 7

The diagram below shows a crane supporting a “heavy object” as shown. The “moving support” can be moved towards the “vertical support” or away from it.



- (a) Explain the role of the “counterweight” and “cable” in this structure.


(2 marks)

- (b) Explain how the tension in the cable changes if the “heavy object” is moved to the right by the “moving support”.

(2 marks)

Question 8

The diagram below shows four positions on a rollercoaster track.



- (a) At which point on the track do the occupants of a rollercoaster on the track experience MAXIMUM normal force? Explain.

(2 marks)


- (b) The occupants of the rollercoaster feel 'weightless' at 'A'. Derive an expression relating the instantaneous speed ' v ' of the rollercoaster and the radius of the track ' r ' at 'A' to cause this sensation.

(3 marks)

Question 9


The diagram below shows a positive point charge near a negatively charged plate. Complete the diagram by drawing the resulting electric field created by these charged objects. Include at least 5 field lines.

(3 marks)



Question 10


A cyclist moves their bicycle by pushing on the pedals of their bicycle. The maximum force the cyclist can exert on one pedal is 150 N. The pedals are both 10 cm from their centre of rotation.



- (a) Describe the location of the pedals in relation to the centre of rotation when maximum torque is being exerted by the cyclist. (1 mark)
- (b) Calculate the maximum torque exerted by the cyclist when the pedals are in this position. (2 marks)
- (c) Describe the positions of the pedals when minimum torque is exerted by the cyclist. Explain your answer. (2 marks)

Question 11

The diagram below shows the trajectory of a projectile as it travels from left to right (ie – from X to Y to Z).




	At 'X'	At 'Y'	At 'Z'
A			
B			
C			
D			
E		0	
F		0	

- (a) Which set of vectors (A – F) best illustrates the acceleration experienced by the ball in flight (ignore air resistance)? (1 mark)
- (b) Which set of vectors (A – F) best illustrates the instantaneous velocity of the ball in flight (ignore air resistance)? (1 mark)
- (c) Which set of vectors best illustrates the vertical component of the ball's velocity in flight (ignore air resistance)? (1 mark)
- (d) If air resistance is taken into account, which set of vectors best illustrates the force due to this air resistance experienced by the ball in flight? (1 mark)

Question 12

Two current carrying wires are situated 50.0 cm apart. Wire A is carrying a current of 0.500 A in an upwards direction; wire B is carrying a current of 0.375 A. The diagram below shows this situation.



The point P is located between the two wires. The net magnetic field at this point due to the current in the two wires is calculated to be zero.

- (a) Indicate the direction of the current in Wire B.

(1 mark)

- (b) Determine the location of point P by calculating distance 'x'. Show all working.

(4 marks)

End of Section One

Section Two: Problem-solving 50% (85 Marks)

This section has **six (6)** questions. You must answer **all** questions. Write your answers in the space provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page.


Suggested working time for this section is 90 minutes.

Question 13**(17 marks)**

In 1996, a joint experiment between NASA and Italy was conducted to try and produce the world's first space tether.

In this experiment, a spherical satellite was deployed by the Space Shuttle. The satellite and the Shuttle were connected by a 20 km long conducting tether (wire). In theory, as the Shuttle dragged the satellite and tether through the Earth's magnetic field, electric power would be produced which could be utilised by the satellite.


The diagram below shows a view of this situation looking down towards the earth. The arrows show the direction of the orbital motion of the satellite, tether and space shuttle. The direction of the earth's magnetic field at this point in its orbit is shown as downwards.



- (a) In your own words, explain the electromagnetism principle that was employed in this experiment to generate the electric power.

(2 marks)

- (b) On the diagram, draw an arrow to show the direction of the conventional electric current that was generated by the motion of the tether. (1 mark)
- (c) One of the design faults that had to be overcome with this setup was that the electric power generated slowed the shuttle and satellite down. (2 marks)
- (i) Explain why the orbital speed of the Shuttle, tether and satellite would have reduced due to the electric power generated. (2 marks)
- (ii) The electric power generated was stored in batteries on the satellite. The stored charge from this battery was used to combat the problem described in part (i); it could even be used to increase the orbital speed of the Shuttle, tether and satellite. Explain how this could occur using electromagnetism concepts. As part of your answer, show on the diagram the direction current would have to flow from the batteries to achieve this. (3 marks)



The Shuttle, tether and satellite followed an orbit around the earth with an average altitude 296 km.

- (d) Using data from your Formulae and Constants Sheet, calculate the orbital speed of the Shuttle, tether and satellite around the earth.

(4 marks)

The maximum EMF generated by the tether was about 3500 V.

- (e) As the tether was unfurled to its full length, scientists predicted that the EMF generated would increase in a highly predictable way. Explain.

(2 marks)

- (f) Use the maximum EMF generated and previous data in this question to calculate the strength of the earth's magnetic field at this altitude. [If you were unable to calculate the orbital speed in part (d), use a value of $7.50 \times 10^3 \text{ ms}^{-1}$]

(3 marks)

Question 14**(16 marks)**

An archer is challenged to hit an apple with an arrow.

The apple is thrown vertically upwards into the air with a velocity of 10ms^{-1} . At the same instant, the archer – who is located a horizontal distance of 17.2m from where the apple is launched – fires the arrow at an angle (θ) and speed (v). The apple and arrow are both launched from the same height.

The archer manages to hit the apple 4.00m above its launch height on its downward path.

- (a) Show through calculations that the maximum height attained by the apple is 5.10m. (3 marks)


- (b) In the space below, draw a diagram showing the paths taken by the arrow and the apple from launch until impact. Also include any relevant distances to indicate scale. (3 marks)

- (c) Calculate the time taken for the arrow to hit the apple. (4 marks)

- (d) Hence, calculate the horizontal component of the arrow's launch velocity. [If you were unable to calculate an answer for part (c), use a time of 1.30s]
- (2 marks)
- (e) Finally, using the information you have calculated earlier in this question, determine the initial launch velocity of the arrow. Show all working. Include a vector diagram in your answer. [If you were unable to calculate an answer for part (d), use a horizontal velocity of 11.0ms^{-1}]
- (4 marks)

Question 15**(15 marks)**

The graph below shows how the torque on one (1) coil in an experimental DC motor varies over time.



(a) On the graph, mark one (1) point where:

(i) the coil would be parallel to the magnetic field. Label this point 'X'.

(1 mark)

(ii) the coil would be perpendicular to the magnetic field. Label this point 'Y'.

(1 mark)

(iii) Briefly explain your answer to part (i). A simple diagram should be included in your explanation.

(2 marks)

- (b) The shape of this graph suggests the presence of a commutator. Explain. (2 marks)

Some specifications for the coil and the DC motor are shown below:

Current in coil (I) = 0.500 A

Dimensions of coil (square shape): 20 cm x 20 cm

Number of turns (n) = 20

- (c) Using appropriate formulae from your Formulae and Constants Sheet, show that the maximum torque (T_{MAX}) generated by the coil can be derived by the following expression:

$$\tau_{\text{max}} = IBAn$$


Where 'B' is equal to the magnetic field strength in the DC motor.

(3 marks)

- (d) Using the expression in part (c), the graph at the beginning of this question, and the specifications of the DC motor, calculate the magnetic field strength B in the DC motor. (3 marks)

Question 16**(16 marks)**

A 60.0 kg person is standing on the end of a uniform 5.00 m long viewing platform with a mass of 500 kg. The platform is horizontal and extends out from the side of a cliff face. It is supported by a steel beam that extends from the side of the cliff face to the end of the platform. The platform is shown below. 'P' is the point of contact between the viewing platform and the cliff face.




- (a) In the space below, draw a free body diagram showing all of the forces acting on the viewing platform with the 60.0 kg person standing in the position stated at the beginning of the question. Label the forces appropriately.

(3 marks)

- (b) Show, with a calculation, that the size of the force exerted by the steel beam is approximately $6.1 \times 10^3 \text{ N}$.
(5 marks)
- (c) Hence, calculate the magnitude of the force exerted by 'P' (ie – the cliff face) on the viewing platform. You may need to alter your answer to part (a) as a result of this calculation.
(5 marks)
- (d) The person now walks towards the cliff face. In your own words, describe how the force exerted by the beam would change as this occurred. Explain your answer.
(3 marks)

Question 17**(12 marks)**

Two particles are injected with the same initial velocity v towards the right and separated by a distance d into a region of uniform, constant magnetic field B as shown in the diagram below. The particles have the same mass m and opposite charges $+q$ and $-q$. Both particles follow the curved radius r . You may ignore gravity in the following.



Show that:

(3 marks)

$$v = \frac{q}{m} B r$$

Consider now the particular case where one particle is an electron with a mass m_e and charge $-e$, while the other particle is a positron with the same mass and the charge $+e$. The distance $d = 1.7\text{mm}$. The magnetic field strength $B = 0.0040\text{ T}$. The radius of each orbit is 2.8 mm .

b) Cross out the incorrect option:

The upper particle is the electron/positron.

Two charges q_1 and q_2 separated by a distance d will experience an electric force of

$$F_e = 9.0 \times 10^9 \frac{q_1 q_2}{d^2}$$

Where F_e is the force in newtons if q_1 and q_2 are in coulombs and d is in metres.

c) Calculate the magnitude of the **electric force** between the particles at injection. (2 marks)


a) Calculate the **magnetic** force on each particle. (3 marks)

b) Explain briefly why the formula derived in Part (a) ignored the existence of the electric force.





(2
marks)

Question 18**(10 marks)**

A small object of mass 50.0 g is being rotated freely in a vertical circle of radius 1.50 m. It is attached to a string of the same length. At the position shown (ie – the top of the vertical circle), the tension in the string is momentarily equal to zero. The string is able to withstand a maximum tension of 2.50 N before it snaps.



- (a) Which of the arrows below best describes the direction of the object's motion at point 'X'? (1 mark)

A	
B	
C	
D	

ANSWER: _____

- (b) Show via calculation that the object is travelling with a speed 3.83 ms^{-1} when it is at the top of the vertical circle. (3 marks)

- (c) Given that the object is rotating freely under the influence of gravity, calculate what its speed would be if it reached the bottom of the circle at point 'Y'.
(3 marks)
- (d) Hence determine whether the string would snap before the object reaches Y'. Support your answer with a calculation.
[If you were unable to calculate an answer for part (c), use a value of 8.60 ms^{-1}]
(3 marks)

End of Section 2

Section Three: Comprehension and Data Analysis**20% (36 Marks)**

This section contains **two (2)** questions. You must answer both questions. Write your answers in the space provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page.

Suggested working time for this section is 40 minutes.

Question 19**(18 marks)**

A group of students conducted an investigation to calculate an experimental value for ' μ_0 ' – **the magnetic constant**. They had the following equation from their Formulae and Constants Sheet at their disposal:

$$B = \frac{\mu_0}{2\pi} \frac{I}{r}$$

Where μ_0 = the magnetic constant; I = current in the conductor; r = distance from the conductor.

They also had the following equipment at their disposal:

1 x power pack; 1 x ammeter; 1 x 50cm length of straight conductor; 1 x rheostat; 1 x switch; 2 x wooden stands; 4 x wooden blocks; 1 x orienteering compass; 1 x ruler; enough connecting wires and alligator clips as required.

The horizontal straight conductor is connected into a circuit that allows the current flowing through it to be controlled. **This current is set at a constant value of 5.00 A.**


The horizontal straight conductor is aligned in a North-South direction using the compass. It is elevated 40 cm above the bench by the wooden stands.

The compass is set on wooden blocks directly beneath the straight conductor. The distance the compass is below the straight conductor ' r ' can be varied.


The earth's magnetic field strength (B_E) is known to be equal to 5.00×10^{-5} T and can be considered to be horizontal in this part of the laboratory.

The diagrams below offer some more information about this set up.


SIDE VIEW



TOP VIEW



When the current in the straight conductor is turned on, the compass needle deviates through an angle ' θ '. This deviation is due to the magnetic field produced by the current-carrying conductor (B_I). With this orientation of the straight conductor, ' B_E ' and ' B_I ' are both perpendicular to each other. Their relationship with ' θ ' is shown below:



$$\frac{B_I}{B_E} = \frac{\mu_0 I}{2\pi r}$$


$$\therefore \alpha_I = \alpha_E \frac{\mu_0 I}{2\pi r} \quad \text{where } \alpha_E = \frac{B_E}{\mu_0 I}$$

Using the above concept, the students were able to calculate experimental values for ' B_I ' at different distances ' r ' from the wire. They collected two values for ' θ ' at each distance ' r ' by reversing the current for each trial. The values they collected are displayed in the table below:


r (m)	θ_{left}	θ_{right}	θ_{average}	$\tan \theta$	B_I ($\times 10^{-5}$ T)	$1/r$ (m^{-1})
0.050	21.5°	22.1°	21.8°	0.400		2.0×10^1
0.10	10.7°	11.9°	11.3°	0.200	1.00	1.0×10^1
0.15	6.5°	8.7°	7.6°	0.134	0.67	
0.20		6.2°	5.7°	0.100	0.50	5.0
0.25	4.3°	4.7°	4.5°	0.080	0.40	4.0
0.30	3.4°	4.0°	3.7°	0.066	0.33	3.3

- (a) On the diagram below, draw an arrow on the straight conductor to show the direction of conventional current that would have produced the deflection shown on the compass. (1 mark)

TOP VIEW



- (b) Explain why the students collected two different values of ' θ ' (θ_{left} and θ_{right}). As part of your answer, also explain why wooden stands are used to elevate the straight conductor instead of steel retort stands. (3 marks)
- (c) The table is incomplete. Calculate values for these missing quantities and write them in the table. Show your working in the space below. Round to the correct number of significant figures. (3 marks)
- (d) On the grid on the next page, plot a graph of ' B_l ' against ' $1/r$ ' (place ' B_l ' values on the vertical axis). Draw a line of best fit for your data. (4 marks)
- (e) Calculate the gradient of the line of best fit. Show working below. Include units. (4 marks)
- (f) Use your answer from part (e) to calculate an experimental value for the magnetic constant ' μ_0 '. Show your working. (3 marks)



Question 20**(18 marks)**

“Cassini Mission to Saturn”

From 10 Cassini fact sheet

(Source: https://saturn.jpl.nasa.gov/system/downloadable_items/10_cassini_fact_sheet.pdf)

The Cassini mission to Saturn is the most ambitious effort in planetary space exploration ever mounted.

Launched in 1997, Cassini will reach Saturn in 2004 after an interplanetary cruise spanning nearly seven years. Along the way, it has flown past Venus, Earth and Jupiter in “gravity assist” manoeuvres to increase the speed of the spacecraft.

The Mission

In manoeuvres called gravity-assist flyby's, Cassini flew twice past Venus, then once each past Earth and Jupiter. The spacecraft's speed relative to the Sun increased as it approached and swung around each planet, giving Cassini the cumulative boost it needs to reach Saturn.

Cassini executed its first Venus flyby on April 26, 1998, at an altitude of 287.2 kilometres. The second Venus flyby took it within 600 kilometres of the planet on June 24, 1999. Two months later, on August 18, 1999, Cassini swung past Earth at an altitude of 1,171 kilometres. It flew by Jupiter at an altitude of 9.7 million kilometres on December 30, 2000.

Upon reaching Saturn on July 1, 2004, Cassini will fire its main engine for about 96 minutes to brake the spacecraft's speed and allow it to be captured as a satellite of Saturn. Passing through a gap between two of Saturn's rings, called F and G rings, Cassini will swing in close to the planet – to an altitude only one-sixth the diameter of Saturn itself – to begin the first of 75 orbits during the rest of its four-year mission.

Saturn is colder than Jupiter, but the colours of Saturn's cloud layers are due to the same basic cloud chemistry as on Jupiter. Near the top of the atmosphere, the ammonia becomes cold enough to crystallize into ice particle clouds, much like high cirrus clouds in Earth's skies. These ammonia clouds are the visible part of Saturn. Gravity at the top of Saturn's clouds is similar to the gravity near the surface of Earth.

Saturn is $9\frac{1}{2}$ times farther from the Sun than Earth is, so it receives only about 1 percent as much sunlight per square metre as does Earth.

The Rings

Although the best telescopes on Earth show three nested main rings about Saturn, we now know that the ring system is collection of thousands of ringlets. They are not solid but rather are made up of countless unconnected particles, ranging in size from nearly invisible dust to icebergs the size of a house. The spacing and width of the ringlets are orchestrated by gravitational tugs from a retinue of orbiting moons and moonlets, some near ring edges but most far beyond the outermost main rings.

And what is the origin of the rings themselves? One theory is that they are the shattered debris of moons broken apart by repeated meteorite impacts. Scientists believe that Saturn's ring system may even serve as a partial model for the disk of gas and dust from which all the planets formed about the early Sun. The Cassini mission will undoubtedly provide important clues to help determine the answers.

The Cassini Spacecraft

The Cassini spacecraft, including the orbiter and the Huygens probe, is one of the largest, heaviest and most complex interplanetary spacecraft ever built. The orbiter alone weighs 2,125 kilograms. When the 320-kilogram Huygens probe and a launch vehicle adapter were attached and 3,132 kilograms of propellants were loaded, the spacecraft at launch weighed 5,712 kilograms. Of all interplanetary spacecraft, only the two Phobos spacecraft sent to Mars by the former Soviet Union were heavier.

- (a) Describe how Cassini was able to increase its speed relative to the Sun by using "gravity-assist swing-by's" around planets like Venus and the Earth. Use a diagram as part of your answer.

(3 marks)

- (b) The rings of Saturn consist of “countless unconnected particles, ranging in size from nearly invisible dust to icebergs the size of a house.” In a particularly thin ring, the orbital speed of each these vastly different masses would be the same. Using relevant formulae, explain.
(3 marks)
- (c) Cassini’s orbit around Saturn is at an altitude of only one-sixth the diameter of Saturn itself. Saturn’s diameter is 116 464km; and its mass is 5.68×10^{26} kg.
- (i) Calculate the radius of Cassini’s orbit around Saturn (in metres).
(2 marks)
- (ii) Hence, calculate the orbital speed of Cassini at this altitude. Show working.
(3 marks)

- (d) A quote from paragraph 6 states that: "Gravity at the top of Saturn's clouds is similar to the gravity near the surface of Earth."
- (i) The data in part (c) illustrates how much bigger in volume Saturn is compared to the earth. Hence, explain how gravity at the top of Saturn's clouds could be similar to the gravity near the surface of Earth.
- (3 marks)
- (ii) Hence, using data provided in part (c), calculate an estimate for the height of the clouds above Saturn's surface.
- (4 marks)

End of Section 3

Additional working space

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Additional working space

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Additional working space

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Additional working space

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Additional graph if required.



End of examination