

Discrete Convolution and Correlation

Convolution is a mathematical operation equivalent to finite impulse response (FIR) filtering. It is a method of finding the zero-state response of relaxed linear time-invariant systems.

Correlation is a measure of similarity between two signals and is found using a process similar to convolution. There are two types of correlation:

- i) Cross correlation.
- ii) Auto-correlation.

Convolution can be represented as \rightarrow convolution operator

$$y(n) = x(n) * h(n) = h(n) * x(n)$$

$$= \sum_{k=-\infty}^{\infty} x(k) h(n-k) = \sum_{k=-\infty}^{\infty} h(k) x(n-k)$$

Properties of convolution:

1. Commutative property: $x(n) * h(n) = h(n) * x(n)$
2. Associative property: $[x(n) * h_1(n)] * h_2(n) = x(n) * [h_1(n) * h_2(n)]$
3. Distributive property: $x(n) * [h_1(n) + h_2(n)] = x(n) * h_1(n) + x(n) * h_2(n)$
4. Shifting property:

If $x(n) * h(n) = y(n)$, then $x(n-k) * h(n-m) = y(n-k-m)$

5. Convolution with an impulse: $x(n) * \delta(n) = x(n)$

Ex-2.2 find $y(n)$ if $x(n) = n+3$ for $0 \leq n \leq 2$

$$h(n) = \alpha^n u(n) \text{ for all } n$$

\Rightarrow we have $y(n) = \sum_{k=-\infty}^{\infty} x(k) h(n-k)$

Given $x(n) = n+3$ for $0 \leq n \leq 2$

$$h(n) = \alpha^n u(n) \text{ for all } n$$

$h(n) = 0$ for $n < 0$, so the system is causal. $x(n)$ is a causal finite duration sequence whose value is zero for $n > 2$. Therefore,

$$\begin{aligned} y(n) &= \sum_{k=0}^2 x(k) h(n-k) \\ &= \sum_{k=0}^2 (k+3) \alpha^{n-k} u(n-k) \\ &= 3\alpha^n u(n) + 4\alpha^{n-1} u(n-1) + 5\alpha^{n-2} u(n-2) \end{aligned}$$

Ex-2.3

Given, $x(n) = 3^n u(n)$ and $h(n) = \left(\frac{1}{3}\right)^n u(n)$

A causal signal is applied to a causal system

$$\begin{aligned} \therefore y(n) &= \sum_{k=0}^n x(k) h(n-k) \\ &= \sum_{k=0}^n 3^k \left(\frac{1}{3}\right)^{n-k} \\ &= \left(\frac{1}{3}\right)^n \sum_{k=0}^n 3^k \times 3^k \end{aligned}$$

SUBJECT:

$$\sum_{k=0}^n r^k = \frac{r^{n+1} - 1}{r - 1}$$

NO

DATE

/ /

SAT	SUN	MON	TUE	WED	THU	FRI
<input type="checkbox"/>						

$$= \left(\frac{1}{3}\right)^n \sum_{k=0}^n (3^2)^k$$

$$= \left(\frac{1}{3}\right)^n \left[\frac{1 - (3^2)^{n+1}}{1 - 3^2} \right]$$

$$= \left(\frac{1}{3}\right)^n \left[\frac{9^{n+1} - 1}{9 - 1} \right] = \left(\frac{1}{3}\right)^n \left[\frac{9^{n+1} - 1}{8} \right]$$

Ex-2.4

Given, $x(n) = 3^n u(n)$ and $h(n) = 2^n u(n)$

Since both $x(n)$ and $h(n)$ are causal, we have

$$y(n) = x(n) * h(n) = \sum_{k=-\infty}^{\infty} x(k)h(n-k) = \sum_{k=0}^n x(k)h(n-k)$$

~~$\sum_{k=0}^n 3^k \cdot 2^{n-k}$~~

$$= \sum_{k=0}^n 3^k \cdot 2^{n-k} = 2^n \sum_{k=0}^n 3^k \cdot 2^{-k} = 2^n \sum_{k=0}^n \left(\frac{3}{2}\right)^k$$

$$= 2^n \left[\frac{1 - \left(\frac{3}{2}\right)^{n+1}}{1 - \frac{3}{2}} \right]$$

$$= 2^n \left[\frac{\left(\frac{3}{2}\right)^{n+1} - 1}{\frac{1}{2}} \right]$$

$$= 2^{n+1} \left[\left(\frac{3}{2}\right)^{n+1} - 1 \right]$$

EX-2.5

Given $x(n) = \text{constant } u(n)$ and $h(n) = \left(\frac{1}{2}\right)^n u(n)$
 $= (-1)^n u(n)$

$$y(n) = x(n) * h(n) = \sum_{k=-\infty}^{\infty} x(k) h(n-k)$$

$$= \sum_{k=-\infty}^{\infty} (-1)^k u(k) \left(\frac{1}{2}\right)^{(n-k)} u(n-k)$$

$$= \sum_{k=0}^n (-1)^k \left(\frac{1}{2}\right)^{n-k}$$

$$= \left(\frac{1}{2}\right)^n \sum_{k=0}^n (-1)^k (2)^k$$

$$= \left(\frac{1}{2}\right)^n \sum_{k=0}^n (-2)^k$$

$$= \left(\frac{1}{2}\right)^n \left[\frac{1 - (-2)^{n+1}}{1 - (-2)} \right]$$

$$= \left(\frac{1}{2}\right)^n \left[\frac{1 + 2(-2)^n}{3} \right]$$

$$= \frac{1}{3} \left(\frac{1}{2}\right)^n + \left(\frac{1}{2}\right)^n \cdot \frac{2(-2)^n}{3}$$

$$= \frac{1}{3} \left(\frac{1}{2}\right)^n + \frac{2(-1)^n}{3} \quad \text{for } n > 0$$

$$= \frac{1}{3} \left(\frac{1}{2}\right)^n u(n) + \frac{2}{3} (-1)^n u(n)$$

Ex 2.6

Given $x(n) = u(n)$, $h(n) = u(n-3)$


$$\begin{aligned}
 y(n) &= x(n) * h(n) = \sum_{k=-\infty}^{\infty} u(k) u(n-3-k) \\
 &\quad \left| \begin{array}{l} \text{number of terms} = \text{last value} \\ -1 \\ 0 \end{array} \right. \\
 &\quad \left| \begin{array}{l} \text{first value} + 1 \\ -1 \\ 0 \end{array} \right. \\
 &\quad \left| \begin{array}{l} (n-3-0)+1 \\ n-3+1 \\ n-2 \end{array} \right. \\
 &\quad \left| \begin{array}{l} 0 \leq k \leq n-3 \\ n-3 \end{array} \right. \\
 &\therefore y(n) = \sum_{k=0}^{n-3} u(k) u(n-3-k) \\
 &= \sum_{k=0}^{n-3} (1) (1) = n-2
 \end{aligned}$$

Ex 2.7

We know that the output $y(n)$ is given by

$$y(n) = \sum_{k=-\infty}^{\infty} x(k) h(n-k)$$

From this we can observe that to obtain the n th value of the output sequence we must form the product $x(k) h(n-k)$ and sum the values of the resulting sequence. The two component sequences are shown in figure as a function of k , with $h(n-k)$ shown to reversed values of n .







Fig: Component sequences in evaluating the convolution sum with $h(-n-k)$ shown for several values of n .

As we see in figure, for $n < 0$, $h(-n-k)$ and $x(k)$ have no non-zero samples that overlap, and consequently $y(n) = 0$, $n < 0$. For n greater than or equal to zero but less than N , $h(-n-k)$ and $x(k)$ have ~~2 or~~ nonzero samples that overlap from $k=0$ to $k=n$; thus for $0 \leq n < N$,

$$y(n) = \sum_{k=0}^n a^{n-k} = a^n \frac{1 - a^{(n+1)}}{1 - a^{-1}} = \frac{1 - a^{n+1}}{1 - a}, \quad 0 \leq n < N$$

For, $N-1 \leq n$, the nonzero samples that overlap extend from $k=0$ to $k=N-1$ and thus

$$y(n) = \sum_{k=0}^{N-1} a^{n-k} = a^n \frac{1 - a^N}{1 - a^{-1}} = a^{n-(N-1)} \left[\frac{1 - a^N}{1 - a} \right], \quad n \in \mathbb{Z}$$



Example 2.8

Given $x(n) = \left(\frac{1}{2}\right)^n u(n)$, $h(n) = u(n) - u(n-10)$

For $n < 10$, i.e., for $n \leq 9$,

$$y(n) = x(n) * h(n) = \sum_{k=-\infty}^{\infty} \left(\frac{1}{2}\right)^k u(k) (1) \quad \text{for } 0 \leq n \leq 9$$

$$= \sum_{k=0}^n \left(\frac{1}{2}\right)^k \quad 0 \leq n \leq 9$$

$$= \frac{1 - \left(\frac{1}{2}\right)^{n+1}}{1 - \frac{1}{2}} = \left(\frac{1}{2}\right)^n \left[\frac{1 - \left(\frac{1}{2}\right)^{-n+1}}{1 - \left(\frac{1}{2}\right)^{-1}} \right]$$

$$= \left(\frac{1}{2}\right)^n [2^{n+1} - 1] \quad 0 \leq n \leq 9$$

For $n \geq 10$, i.e., for $n \geq 10$,

$$y(n) = \sum_{k=n-9}^n \left(\frac{1}{2}\right)^k$$

$$= \left(\frac{1}{2}\right)^{n-9} + \left(\frac{1}{2}\right)^{n-9+1} + \left(\frac{1}{2}\right)^{n-9+2} + \dots + \left(\frac{1}{2}\right)^n$$

$$= \left(\frac{1}{2}\right)^{n-9} \left[1 + \frac{1}{2} + \left(\frac{1}{2}\right)^1 + \dots + \left(\frac{1}{2}\right)^9 \right]$$

$$= \left(\frac{1}{2}\right)^{n-9} \left[\frac{1 - \left(\frac{1}{2}\right)^{10}}{1 - \frac{1}{2}} \right] \quad \text{for } n \geq 10$$

$$= 2 \left(\frac{1}{2}\right)^{n-9} - 2 \left(\frac{1}{2}\right)^{n+1}$$

$$= \left(\frac{1}{2}\right)^n [2^{10} - 1]$$

SUBJECT:

NO

DATE

/ /

SAT	SUN	MON	TUE	WED	THU	FRI
<input type="checkbox"/>						

Exer 2.9

$$\text{Let } y(n) = x(n) * h(n) = \sum_{k=-\infty}^{\infty} x(k) h(n-k)$$

$$x(k) = \left(\frac{1}{3}\right)^{-k} u(-k-1) \text{ and } h(k) = u(k-2) \text{ or}$$

Here, $x(k) = 0$ for $k > -1$ and $h(n-k) = 0$ for $k > n-2$.

For $n-1 < -1$, i.e., for $n < 0$, the interval of summation is from $K = -\infty$ to $n-1$.

$$\begin{aligned} \therefore y(n) &= \sum_{k=-\infty}^{n-1} \left(\frac{1}{3}\right)^{-k} \text{ for } n-1 \leq -1 \text{ or for } n \leq 0 \\ &= \left(\frac{1}{3}\right)^{-(n-1)} + \left(\frac{1}{3}\right)^{-(n-2)} + \dots \\ &= \left(\frac{1}{3}\right)^{-(n-1)} \left[1 + \frac{1}{3} + \left(\frac{1}{3}\right)^2 + \dots \right] \\ &= 3^{n-1} \left[\frac{1}{1 - 1/3} \right] = 0.5(3)^n \end{aligned}$$

For $n-1 > -1$, i.e., for $n > 0$, the interval of summation is from $K = -\infty$ to -1 .

$$\begin{aligned} \therefore y(n) &= \sum_{K=-\infty}^{-1} \left(\frac{1}{3}\right)^{-k} \\ &= \sum_{K=1}^0 \left(\frac{1}{3}\right)^k \\ &= \frac{1/3}{1 - 1/3} \\ &= 0.5 \end{aligned}$$

Liberty

Example 2.30

a) Given, $h(n) = \delta(n) - \delta(n-2)$

The step response $r(n) = h(n) * u(n)$

$$= [\delta(n) - \delta(n-2)] * u(n)$$

$$= \delta(n) * u(n) - \delta(n-2) * u(n)$$

$$= u(n) - u(n-2)$$

b) Given, $h(n) = \left(\frac{1}{q}\right)^n u(n)$

$$\therefore r(n) = \left(\frac{1}{q}\right)^n u(n) * u(n)$$

$$= \sum_{k=-\infty}^{\infty} u(k) \left(\frac{1}{q}\right)^{n-k} u(n-k)$$

$$= \sum_{k=0}^n \left(\frac{1}{q}\right)^{n-k}$$

$$= \left(\frac{1}{q}\right)^n \sum_{k=0}^n \left(\frac{1}{q}\right)^{-k}$$

$$= \left(\frac{1}{q}\right)^n \sum_{k=0}^n q^k$$

$$= \left(\frac{1}{q}\right)^n \left[\frac{q - q^{n+1}}{1 - q} \right]$$

c) Given, $h(n) = n u(n)$

$$\begin{aligned}\therefore r(n) &= h(n) * u(n) \\ &= n u(n) * u(n) \\ &= \sum_{k=0}^n k u(k) u(n-k) \\ &= \sum_{k=0}^n k\end{aligned}$$

\approx

d) Given, $h(n) = u(n)$

$$\begin{aligned}\therefore r(n) &= h(n) * u(n) \\ &= u(n) * u(n) \\ &= \sum_{k=-\infty}^{\infty} u(k) u(n-k) \\ &= \sum_{k=0}^{n-1} 1 \\ &= n + 1\end{aligned}$$

\approx

Show that \approx an impulse response and convolution.

Methods to Compute the Convolution sum of two sequences $x(n)$ and $h(n)$.

Folding Method

Method 1: Linear Convolution Using Graphical Method

Method 2: Linear Convolution using Tabular Array

Method 3: Linear Convolution using Tabular Method

Method 4: Linear Convolution using Matrices.

Method 5: Linear Convolution using the Sum-by-Column Method

Method 6: Linear Convolution using the Flip, Shift, Multiply, and Sum Method.

Example 2.31 Determine the Convolution sum of two sequences.

$$x(n) = \{1, 2, 1, 3\}, \quad h(n) = \{1, 2, 2, 0\}$$

$\Rightarrow x(n)$ starts at $n_1=0$ and $h(n)$ starts at $n_2=-1$. Therefore, the starting sample of $y(n)$ is at

$$n = n_1 + n_2 = 0 - 1 = -1$$

$x(n)$ has 4 samples, $h(n)$ has 4 samples. Therefore, $y(n)$ will have $N = 4 + 4 - 1 = 7$ samples, i.e., from

$$n = -1 \text{ to } n = 5$$

Method 1 Graphical method:

We know that

$$y(n) = \sum_{k=-\infty}^{\infty} x(k)h(n-k)$$

$$\text{For } n = -1, \quad y(-1) = \sum_{k=-\infty}^{\infty} x(k)h(-1-k)$$

$$= 1 \cdot 1 = 1$$

SUBJECT:	NO	DATE	/ /	SAT	SUN	MON	TUE	WED	THU	FRI
----------	----	------	-----	-----	-----	-----	-----	-----	-----	-----

for $n=0$, $y(0) = \sum_{k=-\infty}^{\infty} x(k) h(-k) = 4 \cdot 2 + 2 \cdot 1 = 10$

for $n=1$, $y(1) = \sum_{k=-\infty}^{\infty} x(k) h(1-k) = 4 \cdot 2 + 2 \cdot 2 + 1 \cdot 1 = 13$

for $n=2$, $y(2) = \sum_{k=-\infty}^{\infty} x(k) h(2-k) = 4 \cdot 1 + 2 \cdot 2 + 1 \cdot 2 + 3 \cdot 1 = 13$

for $n=3$, $y(3) = \sum_{k=-\infty}^{\infty} x(k) h(3-k) = 2 \cdot 1 + 1 \cdot 2 + 3 \cdot 1 = 10$

for $n=4$, $y(4) = \sum_{k=-\infty}^{\infty} x(k) h(4-k) = 1 \cdot 1 + 3 \cdot 2 = 7$

for $n=5$, $y(5) = \sum_{k=-\infty}^{\infty} x(k) h(5-k) = 3 \cdot 1 = 3$

$\therefore y(n) = \{ 9, 10, 13, 13, 10, 7, 3 \}$

Method 2 Tabular Array:

k	-4	-3	-2	-1	0	1	2	3	4	5	6	7
$x(k)$	-	-	-	-	4	2	1	3	-	-	-	-
$h(-k)$	-	-	-	1	2	2	1	3	-	-	-	-
$n = -1$	$h(-1-k)$	-	1	2	2	1	-	-	-	-	-	-
$n = 0$	$h(-k)$	-	-	1	2	2	1	-	-	-	-	-
$n = 1$	$h(-1-k)$	-	-	-	1	2	2	1	-	-	-	-
$n = 2$	$h(2-k)$	-	-	-	-	1	2	2	1	-	-	-
$n = 3$	$h(3-k)$	-	-	-	-	-	1	2	2	1	-	-
$n = 4$	$h(4-k)$	-	-	-	-	-	-	1	2	2	1	-
$n = 5$	$h(5-k)$	-	-	-	-	-	-	-	1	2	2	1

The starting value at $n = -1$. From the table, we can see that

$$\text{For } n = -1, y(-1) = \sum_{k=-\infty}^{\infty} x(k) h(-1-k) = 4 \cdot 1 = 4$$

$$\text{For } n = 0, y(0) = \sum_{k=-\infty}^{\infty} x(k) h(-k) = 4 \cdot 2 + 2 \cdot 1 = 10$$

$$\text{For } n = 1, y(1) = \sum_{k=-\infty}^{\infty} x(k) h(1-k) = 4 \cdot 2 + 2 \cdot 2 + 1 \cdot 1 = 13$$

$$\text{For } n = 2, y(2) = \sum_{k=-\infty}^{\infty} x(k) h(2-k) = 4 \cdot 1 + 2 \cdot 2 + 3 \cdot 2 + 1 \cdot 1 = 13$$

$$\text{For } n = 3, y(3) = \sum_{k=-\infty}^{\infty} x(k) h(3-k) = 2 \cdot 1 + 1 \cdot 2 + 3 \cdot 2 = 10$$

$$\text{For } n = 4, y(4) = \sum_{k=-\infty}^{\infty} x(k) h(4-k) = 1 \cdot 1 + 3 \cdot 2 = 7$$


$$\text{For } n = 5, y(5) = \sum_{k=-\infty}^{\infty} x(k) h(5-k) = 3 \cdot 1 = 3$$

$$\therefore y(n) = \{4, 10, 13, 13, 10, 7, 3\}$$

Liberty

Method 3 + Tabular Method:

$$\text{Given, } x(n) = \{4, 2, 1, 3\}, h(n) = \{1, 2, 2, 1\}$$



$$\therefore y(n) = 4, 10, 13, 13, 10, 7, 3$$

The starting value of n is equal to -1 , mark the symbol \uparrow at time origin ($n=0$).

$$\therefore y(n) = \{4, 10, 13, 13, 10, 7, 3\}$$

Method Matrices Method:

The given sequences are: $x(n) = \{x(0), x(1), x(2), x(3)\}$
 $= \{4, 2, 1, 3\}$

and $h(n) = \{h(0), h(1), h(2), h(3)\} = \{1, 2, 2, 0\}$

The sequence $x(n)$ is starting at $n=0$ and the sequence $h(n)$ is starting at $n=-1$. So, the sequence $y(n)$ corresponding to the linear convolution of $x(n)$ and $h(n)$ will start at $n = 0 + (-1) = -1$. $x(n)$ is of length 4 and $h(n)$ is also of length 4. So length of $y(n) = 4+4-1 = 7$.

Substituting the sequence values in matrix form and multiplying as shown below, we get,

$$\begin{array}{c}
 \begin{array}{|cccc|} \hline & 1 & 0 & 0 & 0 \\ & 2 & 1 & 0 & 0 \\ & 2 & 2 & 1 & 0 \\ & 1 & 2 & 2 & 1 \\ & 0 & 1 & 2 & 2 \\ & 0 & 0 & 1 & 2 \\ & 0 & 0 & 0 & 1 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 4 \\ 2 \\ 1 \\ 3 \\ \hline \end{array} = \quad \begin{array}{|c|} \hline 4+0+0+0 \\ 4 \cdot 2 + 2 \cdot 1 + 0 + 0 \\ 4 \cdot 2 + 3 \cdot 2 + 1 \cdot 1 + 0 \\ 4 \cdot 1 + 2 \cdot 2 + 2 \cdot 1 + 3 \cdot 2 \\ 40 + 2 \cdot 1 + 1 \cdot 2 + 3 \cdot 2 \\ 0 + 0 + 1 \cdot 1 + 3 \cdot 2 \\ 0 + 0 + 0 + 3 \cdot 1 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 4 \\ 10 \\ 13 \\ 13 \\ 10 \\ 7 \\ 3 \\ \hline \end{array}
 \end{array}$$

$$\therefore y(n) = x(n) * h(n) = \{4, 10, 13, 13, 10, 7, 3\}$$

Note:

To check my convolution is correct or not:

$$\text{sum of } x(n) \times \text{num of } *h(n) = \text{sum of } y(n)$$

Example 2.12 Find the convolution of the signals

$$x(n) = \begin{cases} 2 & n = -2, 0, 1 \\ 3 & n = -1 \\ 0 & \text{elsewhere} \end{cases}$$

$$h(n) = \delta(n) - 2\delta(n-1) + 3\delta(n-2) - \delta(n-3)$$

$$\Rightarrow \text{Given } x(n) = \{2, 3, 2, 2\}; h(n) \{1, -2, 3, -1\}$$

$x(n)$ starts at $n_1 = -2$ and $h(n)$ starts at $n_2 = 0$. The starting sample of $y(n)$ is at $n = n_1 + n_2 = -2 + 0 = -2$.

Since number of samples in $x(n)$ is 4, and the number of samples in $h(n)$ is 4, the number of samples in $y(n)$ will be $4+4-1 = 7$. So $y(n)$ exists from -2 to 4 .

Method 1 Graphical method:

$$\text{We know that } y(n) = \sum_{k=-\infty}^{\infty} x(k) h(n-k)$$

$$\text{For } n = -2, y(-2) = \sum_{k=-\infty}^{\infty} x(k) h(-2-k) = 2 \cdot 1 = 2$$

$$\text{For } n = -1, y(-1) = \sum_{k=-\infty}^{\infty} x(k) h(-1-k) = 2 \cdot (-2) + 3 \cdot 1 = -1$$

$$\text{For } n = 0, y(0) = \sum_{k=-\infty}^{\infty} x(k) h(-k) = 2 \cdot 3 + 3 \cdot (-2) + 2 \cdot 1 = 2$$

$$\text{For } n = 1, y(1) = \sum_{k=-\infty}^{\infty} x(k) h(1-k) = 2 \cdot (-1) + 3 \cdot 3 + 2 \cdot (-3) + 2 \cdot 1 = 5$$

$$\text{For } n = 2, y(2) = \sum_{k=-\infty}^{\infty} x(k) h(2-k) = 3 \cdot (-1) + 2 \cdot 3 + 2 \cdot (-3) + 2 \cdot 1 = -1$$

$$\text{for } n=3, y(3) = \sum_{k=-\infty}^{\infty} x(k) h(3-k) = 2 \cdot (-1) + 2 \cdot 3 = 4$$

$$\text{for } n=4, y(4) = \sum_{k=-\infty}^{\infty} x(k) h(4-k) = 2 \cdot (-1) = -2$$

$$\therefore y(n) = \{2, -1, 2, 5, -1, 4, -2\}$$

Method 2 Tabular array:

K	-5	-4	-3	-2	-1	0	1	2	3	4	5
x(n)	-	-	-	-2	3	2	-	-	-	-	-
h(-k)	-	-	-1	3	-2	1	-	-	-	-	-
n=-2	h(-2-k)	-1	3	-2	1	-	-	-	-	-	-
n=-1	h(-1-k)	-	-1	3	-2	1	-	-	-	-	-
n=0	h(-k)	-	-1	3	-2	1	-	-	-	-	-
n=1	h(1-k)	-	-1	3	-2	1	-	-	-	-	-
n=2	h(2-k)	-	-	-	-1	3	-2	1	-	-	-
n=3	h(3-k)	-	-	-	-	-1	3	-2	1	-	-
n=4	h(4-k)	-	-	-	-	-	-1	3	-2	1	-

The starting sample of $y(n)$ is at $n=-2$. $y(n)$ is calculated as shown below:

From the table, we can see that

$$\text{For } n=-2, y(-2) = \sum_{k=-\infty}^{\infty} x(k) h(-2-k) = 2 \cdot 1 = 2$$

$$\text{For } n=-1, y(-1) = \sum_{k=-\infty}^{\infty} x(k) h(-1-k) = 2 \cdot (-1) + 3 \cdot 1 = -1$$

$$\text{For } n=0, y(0) = \sum_{k=-\infty}^{\infty} x(k) h(-k) = 2 \cdot 3 + 3 \cdot (-2) + 2 \cdot 1 = 2$$

$$\text{for } n=1, y(1) = \sum_{k=-\infty}^{\infty} x(k) h(1-k) = 2 \cdot (-1) + 3 \cdot 3 + 2 \cdot (-2) + 2 \cdot 1 = 5$$


$$\text{For } n=2, y(2) = \sum_{k=-\infty}^{\infty} x(k) h(2-k) = 3 \cdot (-1) + 2 \cdot 3 + 2 \cdot (-3) = -1$$

$$\text{For } n=3, y(3) = \sum_{k=-\infty}^{\infty} x(k) h(3-k) = 2 \cdot (-1) + 2 \cdot 3 = 4$$

$$\text{For } n=4, y(4) = \sum_{k=-\infty}^{\infty} x(k) h(4-k) = 2 \cdot (-1) = -2$$

$$\therefore y(n) = \{2, -1, 2, 5, -1, 4, -2\}$$

Method 3 Tabular Method:



$$\therefore y(n) = \{2, -1, 2, 5, -1, 4, -2\}$$

Method 4 Matrices Method:

The given sequences are: $x(n) = \{x(0), x(1), x(2), x(3)\}$

$$= \{2, 3, 2, 2\}$$

$$\text{and } h(n) = \{h(0), h(1), h(2), h(3)\} = \{1, -2, 3, -1\}$$

The sequence $x(n)$ is starting at $n=-2$ and the sequence $h(n)$ is starting at $n=0$. So the sequence $y(n)$ corresponding to the linear convolution of $x(n)$ and $h(n)$ will start at $n=-2$.

$n = -2 + 0 = -2$. $x(n)$ is of length 4 and $h(n)$ is also of length 4. So length of $y(n) = 4 + 4 - 1 = 7$.

Substituting the sequence values in matrix form and multiplying, we get the convolution of $x(n)$ and $h(n)$.

$$\begin{bmatrix}
 1 & 0 & 0 & 0 \\
 -2 & 1 & 0 & 0 \\
 3 & -2 & 1 & 0 \\
 -1 & 3 & -2 & 1 \\
 0 & -1 & 3 & -2 \\
 0 & 0 & -1 & 3 \\
 0 & 0 & 0 & -1
 \end{bmatrix}
 \begin{bmatrix}
 2 \\
 3 \\
 2 \\
 2
 \end{bmatrix}
 =
 \begin{bmatrix}
 2 \cdot 1 + 0 + 0 + 0 \\
 2 \cdot (-2) + 3 \cdot 1 + 0 + 0 \\
 2 \cdot 3 + 3 \cdot (-2) + 2 \cdot 1 + 2 \cdot 0 \\
 2 \cdot (-1) + 3 \cdot 3 + 2 \cdot (-2) + 2 \cdot 1 \\
 0 + 3 \cdot (-2) + 2 \cdot 3 + 2 \cdot (-2) \\
 0 + 0 + 2 \cdot (-2) + 2 \cdot 1 \\
 0 + 0 + 0 + 2 \cdot (-1)
 \end{bmatrix}
 =
 \begin{bmatrix}
 2 \\
 -1 \\
 2 \\
 5 \\
 -1 \\
 4 \\
 -2
 \end{bmatrix}$$

$$\therefore y(n) = x(n) * h(n) = \{2, -1, 2, 5, -1, 4, -2\}$$

Deconvolution:

Deconvolution is the process of finding the input $x(n)$ to [or impulse response $h(n)$] applied to the system once the output $y(n)$ and the impulse response $h(n)$ [or the input $x(n)$] of the system are known.

The Z-transform also can be used for deconvolution operation.

$$Y(z) = X(z)/H(z) \quad \text{or} \quad X(z) = \frac{Y(z)}{H(z)}$$

To solve this, we have to substitute $z = e^{j\omega}$

SUBJECT:

NO

DATE

/ /

SAT SUN MON TUE WED THU FRI

Ex-2.2

$$\text{Given, } h(n) = \{2, 1, 0, -1, 3\}$$

$$\therefore H(z) = \sum_{n=-\infty}^{\infty} h(n) z^{-n} = \sum_{n=0}^{\infty} h(n) z^{-n}$$

$$= 2 + z^{-1} + 0 - z^{-3} + 3z^{-4} = 2 + z^{-1} - z^{-3} + 3z^{-4}$$

$$\text{and } g(n) = \{2, -5, 1, 1, 6, -11, 6\}$$

$$\therefore Y(z) = \sum_{n=-\infty}^{\infty} g(n) z^{-n} = \sum_{n=0}^{6} g(n) z^{-n}$$

$$= 2 - 5z^{-1} + z^{-2} + z^{-3} + 6z^{-4} - 11z^{-5} + 6z^{-6}$$

$$\therefore X(z) = \frac{Y(z)}{H(z)} = \frac{2 - 5z^{-1} + z^{-2} + z^{-3} + 6z^{-4} - 11z^{-5} + 6z^{-6}}{2 + z^{-1} - z^{-3} + 3z^{-4}}$$

$$1 - 3z^{-1} + 2z^{-2}$$

$$2 + z^{-1} - z^{-3} + 3z^{-4} \left[\begin{array}{r} 2 - 5z^{-1} + z^{-2} + z^{-3} + 6z^{-4} - 11z^{-5} + 6z^{-6} \\ 2 + z^{-1} - z^{-3} + 3z^{-4} \end{array} \right]$$

$$-6z^{-1} + z^{-2} + 2z^{-3} + 3z^{-4} - 11z^{-5} + 6z^{-6}$$

$$-6z^{-1} - 3z^{-2} + 3z^{-3} - 9z^{-5}$$

$$9z^{-2} + 2z^{-3} - 2z^{-5} + 6z^{-6}$$

$$9z^{-2} + 2z^{-3} - 2z^{-5} + 6z^{-6}$$

6

$$\therefore X(z) = 1 - 3z^{-1} + 2z^{-2}$$

Taking inverse z-transform, we have input

$$x(n) = 8(n) - 38(n+1) + 28(n+2)$$

$$\therefore x(n) = \{1, -3, 2\}$$

Deconvolution by Recursion:

Assuming $y(n)$ and $h(n)$ are one sided sequences,

$$y(n) = \sum_{k=0}^n x(k) h(n-k)$$

$$x(n) = \frac{y(n) - \sum_{k=0}^{n-1} x(k) h(n-k)}{h(0)}$$

Example 2.23

Given, $y(n) = \{1, 2, 2, 0, 2, 3\}$ and $h(n) = \{1, -1, 1\}$

Let N_1 be the number of samples in $x(n)$, and N_2 be the number of samples in $h(n)$. The number of samples in $y(n) = N_1 + N_2 - 1 = 6$, therefore, $N_1 = 6 - N_2 + 1 = 6 - 3 + 1 = 4$.

Let, $x(n) = \{x(0), x(1), x(2), x(3)\}$

$$\text{we have, } x(n) = \frac{y(n) - \sum_{k=0}^{n-1} x(k) h(n-k)}{h(0)}$$

$$\text{for, } n=0, x(0) = \frac{y(0)}{h(0)} = \frac{1}{1} = 1$$

$$\text{for } n=1, x(1) = \frac{y(1) - x(0) h(1)}{h(0)} = \frac{1 - 1 \cdot (-1)}{1} = 2$$

$$\text{for } n=2, x(2) = \frac{y(2) - x(0) h(2) - x(1) h(3)}{h(0)} = \frac{2 - 1 \cdot 1 - 2 \cdot (-1)}{1} = 3$$

$h(3)$ exist | NO
as $\exists n$, so, $h(3) = 0$

$$\text{For } n=3, x(3) = \frac{y(3) - x(0)h(3) - x(1)h(2) - x(2)h(1)}{h(0)}$$


$$= \frac{0 - 1 \cdot 0 - 2 \cdot 1 - 3(-1)}{1} = \frac{-2 + 3}{1} = 1$$

$$\therefore x(n) = \{1, 2, 3, 1\}$$

Linear Time Invariant

Q) Interconnection of LTI System:

Parallel Connection of Systems:



Prove that, If two systems are connected in parallel, the overall impulse response is equal to the sum of two impulse responses.

Proof:

Let the output of system 1 is, $y_1(n) = x(n) * h_1(n)$
 and the output of system 2 is, $y_2(n) = x(n) * h_2(n)$
 \therefore The

Consider two LTI systems connected in parallel having impulse responses $h_1(n)$ and $h_2(n)$. Let the input be $x(n)$.

Output of system 1:

$$y_1(n) = x(n) * h_1(n)$$

Output of system 2:

$$y_2(n) = x(n) * h_2(n)$$

The ^{Overall} output of the system $y(n)$ is,

$$y(n) = y_1(n) + y_2(n)$$

$$= x(n) * h_1(n) + x(n) * h_2(n)$$

$$= \sum_{k=-\infty}^{\infty} x(k) h_1(n-k) + \sum_{k=-\infty}^{\infty} x(k) h_2(n-k)$$

$$= \sum_{k=-\infty}^{\infty} x(k) [h_1(n-k) + h_2(n-k)]$$


$$= \sum_{k=-\infty}^{\infty} x(k) h(n-k)$$

$$= x(n) * h(n)$$

where $h(n) = h_1(n) + h_2(n)$

[proved]

Cascade Connection of Systems:



Prove that, When two LTI systems are connected in cascade, the overall impulse response is the convolution of individual impulse responses.

proof:

Consider two LTI systems connected in cascade having impulse responses $h_1(n)$ and $h_2(n)$. Let the input be $x(n)$, output of the system 1:

$$y_1(n) = x(n) * h_1(n)$$

The output becomes the input of the second system. So, output of the system 2:

$$y_2(n) = y_1(n) * h_2(n)$$

$$= [x(n) * h_1(n)] * h_2(n)$$

$$= x(n) * [h_1(n) * h_2(n)]$$

$$= x(n) * h(n)$$

$$\text{where } h(n) = h_1(n) * h_2(n)$$

[proved]

Circular Shift: Normal shift - (signal to left/right) shift by ≥ 0 outside value 0. For circular shift - (signal to left) shift by ≥ 0 first \rightarrow last element

④ Periodic or Circular Convolution:

Methods of performing periodic or Circular Convolution:

Method 1 : Graphical Method (Concentric Circle Method)

Method 2 : Circular convolution using Tabular Array


Method 3 : Circular convolution using Matrices.

Example 2.2.8

Let $x_3(n)$ be the circular convolution of $x_1(n)$ and $x_2(n)$. To find the circular convolution, both sequences must be of same length. Here $x_1(n)$ is of length 6 and $x_2(n)$ is of length 3. Therefore, we append three zeroes to the sequence $x_2(n)$ and use concentric circle method to find circular convolution. So we have

$$x_1(n) = \{1, 2, -1, -2, 3, 3\}, \quad x_2(n) = \{3, 2, 1, 0, 0, 0\}$$


Graph all points of $x_1(n)$ on the outer circle in the anticlockwise direction. Starting at the same point as $x_1(n)$, graph all points of $x_2(n)$ on the inner circle in clockwise direction.



Multiply corresponding samples on the circles and add to obtain


$$\begin{aligned}x_3(0) &= 1 \cdot 3 + 2 \cdot 0 + (-1) \cdot 0 + (-2) \cdot 0 + 3 \cdot 3 + 1 \cdot 2 \\&= 8\end{aligned}$$

Rotate the inner circle in anticlockwise direction by one sample and multiply the corresponding samples and add the products to obtain




$$\begin{aligned}x_3(1) &= 1 \cdot 2 + 2 \cdot 3 + (-1) \cdot 0 + (-2) \cdot 0 + 3 \cdot 0 + 1 \cdot 1 \\&= 9\end{aligned}$$


Continue it to obtain $x_3(2)$, $x_3(3)$, $x_3(4)$, and $x_3(5)$




$$\begin{aligned}x_3(2) &= 1 \cdot 1 + 2 \cdot 2 + (-1) \cdot 3 + (-2) \cdot 0 + 3 \cdot 0 + 1 \cdot 0 \\&= 2\end{aligned}$$



$$\begin{aligned}x_3(3) &= 1 \cdot 0 + 2 \cdot 1 + (-1) \cdot 2 + (-2) \cdot 3 + 3 \cdot 0 + 1 \cdot 0 \\&= -6\end{aligned}$$

$x_1(n)$ 

$$x_3(4) = 1 \cdot 0 + 2 \cdot 0 + (-1) \cdot 1 + (2) \cdot 2 + 3 \cdot 3 + 1 \cdot 0 \\ = 4$$

 $x_2(n)$ 

$$x_3(5) = 1 \cdot 0 + 2 \cdot 0 + (-1) \cdot 0 + (2) \cdot 1 + 3 \cdot 2 + 1 \cdot 3 \\ = 8$$

Therefore, the circular convolution of $x_1(n)$ and $x_2(n)$ is:

$$x_3(n) = x_1(n) \oplus x_2(n) = \{8, 9, 2, -6, 7, 2\}$$


Example 2.29

Let $x_3(n)$ be the circular convolution of $x_1(n)$ and $x_2(n)$.

Here $x_1(n)$ and $x_2(n)$ are of the same length. So no padding of zeroes is required. The computation of the circular convolution $x_3(n)$ of the given sequences

$$x_1(n) = \{1, 2, 1, 2\} \text{ and } x_2(n) = \{4, 3, 2, 1\}$$

by the concentric circles method is done.



$$x_3(0) = 1 \cdot 4 + 2 \cdot 1 + 1 \cdot 2 + 2 \cdot 3 = 14$$

$$x_3(1) = 1 \cdot 3 + 2 \cdot 4 + 1 \cdot 1 + 2 \cdot 2 = 16$$

$$x_3(2) = 1 \cdot 2 + 2 \cdot 3 + 1 \cdot 4 + 2 \cdot 1 = 14$$

$$x_3(3) = 1 \cdot 1 + 2 \cdot 2 + 1 \cdot 3 + 4 \cdot 2 = 16$$

Therefore, the circular convolution of $x_1(n)$ and $x_2(n)$ is:

$$x_3(n) = x_1(n) \oplus x_2(n) = \{14, 16, 14, 16\}$$

~~Q1~~ Given, $x(n) = \{2, 1, 0, 5\}$ and $h(n) = \{2, 0, 1, 1\}$

- Compute the linear convolution of the following signals.
 $x(n) = \{2, 1, 0, 5\} \quad h(n) = \{2, 0, 1, 1\}$
- Compute the circular convolution of the following signals.
 $x(n) = \{1, 0, 2, 0\} \quad h(n) = \{2, -1, -1, 2\}$

10

10

before,

will

Tabular Method:

		$x(n)$			
		2	1	0	5
$h(n)$	2	4	2	0	10
	0	0	0	0	0
	1	2	1	0	5
	4	2	1	0	5

$$\therefore y(n) = \{4, 2, 2, 13, 2, 5, 5\}$$

2) Circular Convolution (Matrix Method)

Given, $x(n) = \{1, 0, 2, 0\}$ $h(n) = \{2, -1, -1, 2\}$

Both $x(n)$ and $h(n)$ are ~~length~~ of length 4.

Let, $y(n)$ be the Circular convolution of $x(n)$ and $h(n)$.
The circular convolution of the given sequences:

$$x(n) = \{x(0), x(1), x(2), x(3)\}$$

$$= \{1, 0, 2, 0\}$$

$$\text{and } h(n) = \{h(0), h(1), h(2), h(3)\}$$

$$= \{2, -1, -1, 2\}$$

Can be determined using matrices as follows:

$$\begin{bmatrix} h(0) & h(3) & h(2) & h(1) \\ h(1) & h(0) & h(3) & h(2) \\ h(2) & h(1) & h(0) & h(3) \\ h(3) & h(2) & h(1) & h(0) \end{bmatrix} \begin{bmatrix} x(0) \\ x(1) \\ x(2) \\ x(3) \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 2 & -1 & -1 \\ -1 & 2 & 2 & -1 \\ -1 & -1 & 2 & 2 \\ 2 & -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 2+0-2+0 \\ -1+0+4+0 \\ -2+0+4+0 \\ 2+0-2+0 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 3 \\ 0 \end{bmatrix}$$

$$\therefore y(n) = \{0, 3, 3, 0\}$$

SUBJECT:	NO	DATE	/ /	SAT	SUN	MON	TUE	WED	THU	FRI
----------	----	------	-----	-----	-----	-----	-----	-----	-----	-----

Discrete Correlation

Correlation measures the similarity between two discrete signals as one signal is shifted over another. There are two types :

- 1) Cross - Correlation.
- 2) Auto - Correlation.

Applications :

- 1) used in radar and sonar system to find the location of a target by comparing the transmitted and reflected signals.
- 2) image processing.
- 3) Control engineering.

Cross Correlation :

Cross-Correlation is the measure of similarity between two discrete time signals as one signal is shifted with respect to the other.

The Cross Correlation between a pair of sequences $x(n)$ and $y(n)$,

$$R_{xy}(n) = \sum_{k=-\infty}^{\infty} x(k) y(k-n) \quad (i)$$

$$= \sum_{k=-\infty}^{\infty} x(k+n) y(k) ; n = 0, \pm 1, \pm 2, \dots$$

If we wish to fix the sequence $y(n)$ and do shift the sequence $x(n)$, then the correlation can be written as:

$$\begin{aligned} R_{yx}(n) &= \sum_{k=-\infty}^{\infty} y(k) x(k-n) \quad \text{(i)} \\ &= \sum_{k=-\infty}^{\infty} y(k+n) x(k), \quad n = 0, \pm 1, \pm 2, \dots \end{aligned}$$

$$R_{xy}(n) \neq R_{yx}(n)$$

But if the time shift $n=0$, then we get

$$R_{xy}(0) = R_{yx}(0) = \sum_{k=-\infty}^{\infty} x(k) y(k)$$

Comparing the (i) and (ii),

$$R_{xy}(n) = R_{yx}(-n)$$

Prove that, the Cross-correlation is the convolution of one signal with a flipped version of the other, that is: $R_{xy}(n) = x(n) * y(-n)$

where, the symbols have their usual meanings.

State Proof:

The Cross-Correlation between two discrete-time signals $x(n)$ and $y(n)$ is defined as,

$$R_{xy}(n) = \sum_{k=-\infty}^{\infty} x(k) y(k-n)$$

The convolution of two signals $x(n)$ and $h(n)$ is,

$$x(n) * h(n) = \sum_{k=-\infty}^{\infty} x(k) h(n-k)$$

Let $h(n) = y(-n)$, which is the time-reversed or flipped version of $y(n)$, then,

$$\begin{aligned} x(n) * y(-n) &= \sum_{k=-\infty}^{\infty} x(k) y(-(n-k)) \\ &= \sum_{k=-\infty}^{\infty} x(k) y(k-n) = R_{xy}(n) \end{aligned}$$

$$\therefore R_{xy}(n) = x(n) * y(-n) \quad [\text{proved}]$$

Auto-correlation:

Auto-correlation is the correlation of a discrete-time signal with itself to measure similarity at different time shifts.

The auto-correlation of a sequence $x(n)$ is defined as:

$$R_{xx}(n) = \sum_{k=-\infty}^{\infty} x(k) x(k-n) = \sum_{k=-\infty}^{\infty} x(k+n) x(k)$$

If the time shift $n=0$, then we have

$$R_{xx}(0) = \sum_{k=-\infty}^{\infty} x^2(k) \quad \left| \begin{array}{l} R_{xx}(n) \text{ is given by} \\ R_{xx}(n) = x(k) * x(-k) \end{array} \right.$$

P

Example 2.51 Find the auto correlation of $x(n) = a^n u(n)$

→ To compute $R_{xx}(n)$ which is even symmetric, we need to compute the result only for $n \geq 0$ and create its even extension.

$$\begin{aligned} \text{For } n \geq 0, R_{xx}(n) &= \sum_{k=-\infty}^{\infty} x(k) x(k-n) = \sum_{k=n}^{\infty} a^k a^{k-n} \\ &= \sum_{m=0}^{\infty} a^{m+n} a^m = a^n \sum_{m=0}^{\infty} a^{2m} \\ &= \frac{a^n}{1-a^2} u(n) \end{aligned}$$

$$\left. \begin{array}{l} m = k-n \\ \therefore k = m+n \\ k = nh, m = o \end{array} \right|$$

The even extension of this result gives

$$R_{xx}(n) = \frac{a^{|n|}}{1-a^2}$$

=

$$(a^{|n|})^2 = a^{2|n|}$$

(Evening session is over)

With thanks to Prof. Dr. A. S. Salim

With thanks to Prof. Dr. A. S. Salim

(Evening session is over)

With thanks to Prof. Dr. A. S. Salim

(Evening session is over)