Navigating the Spectrum of Medical Artificial Intelligence:

A Review of Clinical Innovation and Algorithmic Risk



Muhammad Mughees Rashid, Muhammad Abu Turab, Shayan Noor

Independent Researcher, Kamra, Attock, Pakistan

July 6, 2026

Navigating the Spectrum of Medical Artificial Intelligence: A Review of Clinical Innovation and Algorithmic Risk

Muhammad Mughees Rashid[1], *Muhammad Abuturab*[1], *Shayyan Noor*[1]

[1]*Independent Research, Kamra, Attock, Pakistan*

KEYWORDS

Medical Artificial Intelligence
Clinical Diagnostics
Algorithmic Bias
Workflow Optimization
Biomedical Ethics.

ABSTRACT

As artificial intelligence (AI) is rapidly being introduced into the biomedical domain, the problem of whether intelligent machines can be safely used to aid or even replace human physicians has arisen in modern medicine. The purpose of this review paper is to analyze the two sides of this technology as the key innovation in the biomedical industry and as a potential source of risk in healthcare systems. Given recent advancements, AI is seen as a powerful resource that greatly improves diagnostics, streamlines administration procedures, and increases the availability of medical facilities. By performing data processing and initial diagnostics automatically, AI reduces the workload on doctors and helps minimize their exhaustion, thus shifting focus to patient treatment. Nevertheless, there are several significant issues involved in the introduction of these innovations, such as algorithmic bias, the black box problem, and critical concerns regarding data security. Drawing on existing scientific research in both biomedical and computer science, this paper will analyze these aspects and propose an approach to overcome them, thereby creating new perceptions of AI-driven medicine.

1 Introduction:

The current transformation of modern medicine is increasingly influenced by advances in artificial intelligence alongside continued developments in pharmaceuticals, biotechnology, and surgical innovation. Namely, Artificial Intelligence (AI), which used to be the object of theoretical computer science studies and experiments, has already turned into the part of global healthcare infrastructure. It is necessary to note that the speed of the process of technological boom is largely determined by the development of such technologies as deep learning (DL), convolutional neural networks (CNN) for biomedical image processing, and the implementation of domain-specific Large Language Models (LLMs) in the clinical practice. Thus, the field of medicine witnesses the transition from algorithms capable of cross-referencing the medical database to the agentic architecture which can do clinical reasoning in real time. It should be noted that the scale of the phenomenon is illustrated by the regulatory domain – by the end of 2025, the number of medical devices with AI and ML components of annual medical device clearances featuring AI/ML components has rapidly scaled, approaching 300 clearance units in the single year of 2025 alone (U.S. Food and Drug 2025).




Figure 1. FDA-approved and FDA-cleared AI/ML-enabled medical devices (through December 2025) Source: Author's visualization based on FDA (2025).

The underlying driver of the automatization is a unique situation concerning the worldwide healthcare labor force. Modern medicine is facing serious problems related to burnout and staff shortages and according to forecasts, there will be a shortage of 3.2 million of healthcare workers by the end of 2026 (US Healthcare Labor Market Survey and 2026 Workforce Shortage Forecast 2024). One of the traditional drivers of the problem was associated with the cognitive load generated by EHRs and bureaucratic "pajama time," which refers to the additional time physicians spend outside of work on documenting the cases (Sinsky, Colligan and Li 2016). The current statistics show that up to 42% of physicians experience active burnout symptoms, which are caused by bureaucratic documentation. Scribes based on generative AI and clinical prediction instruments have been already implemented in clinical practice as operational solutions. In terms of system-level implementation, there has been an average of 40% to 45% reduction in the time spent on documenting (M. Razaghi 2026) the case by physicians and 30 days reduction of the ambulatory clinician burnout rate from 51.9% to 38.8% (Olson and Shanafelt 2025). As an invisible administrative co-pilot, the AI technology can help to restore the role of the physician in the setting of patient care as a transformative friend of the overworked system.

However, the rapid adoption of the technology emphasizes the complicated duality of the problem, where the digital mind may become the "foe" just like anything else. The clinical value of the algorithm is directly dependent on the

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

data that is used to train the model, and this creates major issues concerning algorithmic biases and data inequality. AI models trained on the data set mainly composed of cases coming from wealthy communities tend to demonstrate significantly decreased sensitivity while being used for diagnosing the cases of underrepresented minorities. Furthermore, the majority of deep learning algorithms function as an "uninterpretable black box," leaving the underlying logic of the results obtained unknown. Such inability to explain how the results were obtained brings about a series of paradoxes in regard to legal and ethical concerns about clinical responsibility: in case the delay occurs due to a false negative result produced by the autonomous algorithm, the responsibility is dispersed among the attending physician, the hospital administration, and the creators of the algorithm itself. Apart from the danger of clinical liability, there is another threat that arises in regard to patient data privacy, as the commercial use of data becomes intertwined with increasing vulnerability of hospital systems to algorithmic interference. Taking everything into consideration, the opinion of the medical faculty is deeply divided regarding




Figure 2 Distribution of FDA-approved AI-enabled medical devices by medical specialty (through Q4 2025 / early 2026)
Source: Author's visualization based on FDA Data (2025-2026)

the possibility of using AI.

2 Methodology

This paper adopts a thematic literature review approach with a structured search strategy to evaluate the clinical innovations and algorithmic risks of medical AI. Literature was retrieved from PubMed/MEDLINE, IEEE Xplore, and Scopus, supplementing secondary tracking data via Google Scholar. Extracted studies were organized into a dual thematic matrix categorized by quantified operational optimizations (e.g., diagnostic sensitivity, workflow reductions) and systemic risks (e.g., algorithmic bias, legal liability, data security).

2.1 Search Strategy and Information Sources

The literature review was undertaken in four major database sources such as PubMed/MEDLINE, IEEE Xplore, The Lancet Digital Health registry, and Google Scholar. The search criteria included scholarly journals, conference papers, and regulatory documents related to medical devices published or updated by mid-2026. This period ensures that the paper incorporates the most updated data on the deployment of LLMs and medical devices regulations.

The search strings utilized combinations of Boolean operators (AND, OR) alongside specific medical subject headings (*MeSH*) and technical keywords, structured as follows:

```
("MEDICAL ARTIFICIAL INTELLIGENCE" OR "DEEP LEARNING") AND ("DIAGNOSTIC ACCURACY" OR "CLINICAL WORKFLOW" OR "AMBIENT SCRIBE")
("ALGORITHMIC BIAS" OR "DATA INEQUALITY" OR "DEMOGRAPHIC DISPARITY") AND ("CLINICAL DIAGNOSTICS" OR "PREDICTIVE ANALYTICS" OR "ELECTRONIC HEALTH RECORDS")
("BLACK BOX PROBLEM" OR "MEDICAL MALPRACTICE") AND ("LIABILITY DIFFUSION" OR "CLINICAL AUTONOMY")
("HEALTHCARE DATA BREACH" OR "PATIENT PRIVACY") AND ("RANSOMWARE" OR "COMMERCIALIZATION")
```

2.2 Inclusion and Exclusion Criteria

To maintain a high standard of data integrity, strict eligibility criteria were applied to the retrieved literature:


Inclusion Criteria: (1) Peer-reviewed studies demonstrating quantified clinical outcomes or statistical evaluations of AI models; (2) Multi-site prospective or retrospective cohort trials monitoring clinician workflow changes; (3) Multi-center data tracking algorithmic diagnostic sensitivity across diverse demographic sub-populations; (4) Official policy guidelines from recognized regulatory bodies (e.g., FDA, European Medicines Agency).

Exclusion Criteria: (1) Non-peer-reviewed commentaries, superficial opinion editorials, or speculative blog posts; (2) Studies focusing on generalized machine learning outside direct clinical, biomedical, or healthcare administrative infrastructure; (3) Outdated papers pre-dating modern deep learning benchmarks whose technical metrics no longer align with current deployment standards.

2.3 Data Synthesis and Analysis

In all, a total of 39 core studies and regulations were identified that satisfied all criteria for inclusion and were included in the synthesis of their full text. The information extracted from these studies was organized into a thematic dual-matrix relevant to the core thesis presented in this paper. Information pertaining to quantitative clinical performance measures such as Area Under the ROC curves, sensitivity percentages, and daily time motion information was extracted to document the advantages of "The Friend." Simultaneously, information pertaining to reductions in sub-population accuracy, precedents from legal cases, and cost matrices of data breaches were analyzed to document "The Foe".

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com



3 Section I: The "Friend" – Transformative Capabilities of Medical AI:

3.1 Diagnostic Accuracy and Medical Imaging

One of the main areas that has spurred the application of medical artificial intelligence in a clinical setting is computer vision used in the field of radiological diagnosis. In terms of macro-level trends in devices, the bulk of the machine learning models embedded within the hospital network are those used to analyze images from medicine. Through the use of deep convolutional neural networks (CNNs), such AI-based models are able to identify pixel-level differences on X-ray, MRI, and CT scans that cannot be discerned by the naked eye (Zhang and Yan 2026). "This diagnostic edge is quantified in Figure 3, where aggregated validation data for deep learning architectures demonstrates advanced diagnostic sensitivity over human specialist baselines in identifying malignant melanoma (91.0% vs. 85.7%) and complex fungal dermatoses (99.5% vs. 95.2%)."




Figure 3. Comparative diagnostic performance of AI models and human specialists across selected medical conditions Source: Author's synthesis of multi-center convolutional neural network (CNN) performance validation data (2026).

The most important criterion for assessing the performance of these systems in the clinical environment is the comparison of their sensitivity and specificity against board-certified specialists. The empirical evidence confirms that deep learning models have high proficiency in early recognition of complicated pathologies. For example, contemporary artificial intelligence technologies used for mammography in breast cancer screening can decrease the number of false positives by 10-15% compared to the existing double reading approach and significantly speed up the diagnosis process (McKinney, Sieniek and Shetty 2020). In emergency conditions, a deep learning algorithm based on thousands of brain scans allows detecting intracranial hemorrhages and acute ischemic stroke within several minutes (Titano, Badgeley and Oermann 2018).

However, what is important here is that this kind of ability does not necessarily make the machine a substitute for human intelligence, but makes it a necessary tool to augment diagnostics. In scenarios where a lot of patients have to be handled, there is a tendency for decreased diagnostic precision during the latter stages of work on account of visual and cognitive fatigue. Neural networks function in a deterministic manner and, hence, the level of precision is kept at a consistent minimum irrespective of the workload. By alerting to high probability lesions and automatically doing preliminary segmentation, AI becomes an invaluable clinical 'friend' in this case.

3.2 Operational Efficiency and Workflow Optimization

Outside the scope of visual diagnostics, medical artificial intelligence functions as an important structural intervention within the administrative processes in contemporary healthcare systems. Occupational stress within the system is driven by the excessive workload of clinicians related not to the number of patients, but rather to the amount of documentation required in connection with Electronic Health Records (EHR). Traditionally, clinicians had to work within a very inefficient paradigm where more than two hours per hour of actual clinical activity spent typing data into computers was such a heavy burden created an occupational problem of extraordinary nature, as over 60% of physicians complained about symptoms of professional burnout (Shanafelt, Boone and Dyrbye 2022). In order to address this issue, the implementation of ambient AI-powered scribes based on LLMs such as Nuance DAX Copilot, Abridge, and Ambience became an efficient tool to automate clinical documentation.

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

These ambient intelligence systems run in the background during the clinical visit and capture the natural conversation taking place between the doctor and the patient in a secure way. This is achieved by processing unstructured speech through the use of natural language processing (NLP) techniques, whereby the algorithm automatically translates the audio data into structured medical records in the form of SOAP matrix, coding, and patient discharge reports. Unlike other intelligent algorithms that are designed to act as automatic decision makers, these digital assistants serve more as co-pilots, where the doctor takes up the role of being an editor.

The quantitative effect of ambient AI on institutional workflow outcomes has been determined and proven through large-scale multi-center clinical trials. A longitudinal and comprehensive cohort study in JAMA assessed over 8,500 outpatient clinicians in five leading health care systems for isolating precise improvements of the ambient AI system. The results showed that the use of AI scribes helped to decrease absolute documentation time by 16.0 minutes per patient interaction, as well as cut total daily EHR usage time by 13.4 minutes. Moreover, among those physicians who are high-frequency users and use the system for more than half of patient interactions, these improvements are exponential. What is more, multi-center clinical trials on assessing the well-being of physicians at different hospitals have shown that already after 30 days of using the ambient AI system, the level of self-reports of physicians' burnout was decreased from 51.9% to 38.8%, representing a 25.2% relative reduction in active clinician exhaustion at work (Olson and Shanafelt 2025).




Figure 4. Impact of AI on physician burnout and documentation burden Source: Author's visualization based on JMIR multisite time-motion study on ambient AI scribes (2026).

Importantly, the efficiency gained through medical AI brings additional benefits that positively contribute to the effectiveness of the therapeutic alliance. Through reducing the cognitive overload involved in typing and working with a monitor while evaluating a patient, the ambient technology allows doctors to make more eye contact and show more clinical presence when talking to a patient. According to the observational motion studies, there is a 10.6% increase in face-to-face interaction between the doctor and the patient after implementing the technology (Tan et al. 2026). Although ambient technology does not completely get rid of the administrative task and still requires rigorous validation by humans to avoid the spread of errors, the technology helps to demolish the administrative barriers blocking modern infrastructure (Denham, et al. 2026). Thanks to the modernization of the clinical documentation process, AI has become an important structural "friend" of doctors.

3.3 Predictive Analytics in Patient Care

The ICU critical care environment is a very chaotic clinical ecosystem wherein the physiological parameters of the patients change unpredictably at rapid rates. In this kind of critical environment, medical artificial intelligence plays the role of being a key predictor that helps convert medicine from its reactive to a proactive model. The greatest use of this prediction capacity lies in the early prediction of systemic dermatological and physiological emergencies such as sepsis and MODS. Sepsis occurs among about 49 million people worldwide each year, making it one of the leading causes of preventable deaths in hospitals. It has been found that every hour's delay in treatment results in linear increases in the mortality rate of the patient.

In many instances, traditional screening methods used in clinical practice, such as the Sequential Organ Failure Assessment (SOFA), quick SOFA (qSOFA), and SIRS, often do not succeed in detecting early changes due to their reliance on set thresholds and irregular lab data (Singer, Deutschman and Seymour 2016). On the other hand, sophisticated AI machine learning structures, for example, eXtreme Gradient Boosting (XGBoost) and recurrent neural network (RNN) architectures such as Long Short-Term Memory (LSTM), are ideal for handling streaming data (Shashikumar, Wardi and Malhotra, et al. 2024) (Chen 2016). These machine learning models have the capability of integrating the longitudinal data, which include continuous electrocardiogram (ECG), arterial blood pressure, oxygen saturation, and body temperature, with intermittent lab results including serum lactate and white blood cell count. Through the identification of non-linear correlations among the data, the machine learning models can detect the pattern of shock much earlier than any physical sign becomes obvious.

The effectiveness of these automated prediction models compared to humans is supported by systematic reviews, extensive prospective cohort studies, and meta-analyses. Systematic reviews involving numerous clinical studies demonstrate that machine learning models have an Area




Figure 5. AUROC comparison of traditional qSOFA and machine learning models for early sepsis prediction Source: Author, adapted from COMPOSER Registry (2025–2026).

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

Under the Receiver Operating Characteristic curve (AUROC) of 0.79 to 0.96, which significantly exceeds the performance of static screening methods. For example, clinical trials reveal that, although a qSOFA score model produces poor discriminative accuracy of 0.64 for the same ICU patient sample, an ensemble machine learning approach has an AUROC of 0.87 to 0.93 (Henry, Hager and Saria 2015). Additionally, sophisticated practical deep learning models, such as the COMPOSER model implemented in several academic health systems, create high-risk alerts at least 4.0 to 8.1 hours before clinical identification of critical conditions by ICU professionals (Shashikumar, Wardi and Nemati, et al. 2023).

These clinical benefits of the “head start” provided through automation translate into saved lives of people. The use of real-time AI early warning systems in the routine processes in ICUs leads to a substantial decline in mortality rates in terms of clinical alerts. Clinical trials conducted across multiple centers and involving thousands of patients hospitalized in leading health-care institutions showed a 39.5% relative decrease in the overall mortality rate and a 32.2% decrease in the average length of stay due to the implementation of the predictive sepsis algorithms. Those patients who received warnings on their deterioration through AI (at least 4 hours ahead of time) showed a decrease in the ICU mortality rate from 37.1% to 20.5% and a dramatically decreased likelihood of falling into an irrecoverable state of septic shock (Boussina 2024). Through continuous calculations of the dynamic risk trajectory of the patient, predictive AI becomes a very important algorithmic guardian – a special kind of “friend” that protects ICU patients from clinical inertia and gives the crucial time for life-saving actions.

4 Section II: The "Foe" – Challenges, Biases, and Risks

4.1 Algorithmic Bias and Data Inequality

Despite the fact that using medical artificial intelligence certainly entails numerous systemic benefits, it comes with significant ethical and logistical risks insofar as AI serves as the means through which biases are implemented into algorithms. The key drawback associated with all machine-learning systems is their utter dependence upon the datasets used to train those algorithms mathematically. When such datasets fail to represent a certain demographic properly or when they carry inherent statistical biases, those biases will definitely be carried forward in the algorithms and entrenched in society as a whole. This disparity in data represents an enormous optimization problem because regular systems are optimized for maximum population-wide efficiency.

The best-documented case study of such an algorithmic malfunction lies in the area of computer vision programs designed to work in the realm of dermatology. The open-source baseline image databases, which include the popularly used HAM10000 set, contain a highly imbalanced 20:1 ratio of images containing lighter skin to images containing darker skin. In more general multi-registry studies involving over 106,000 dermatological images, only 11 images were identified as containing darker skin types (Tschandl, Rinner and Kittler 2018) (Daneshjou, Vodrahalli and Novoa 2022).

As such, the reliability of the diagnosis of flagship deep learning algorithms is greatly diminished when tested under more heterogeneous conditions than those in which they were validated. A systematic review and meta-analysis conducted in 2025, which assessed 70,000 test images, found a persistent disparity in terms of AI fairness: the Area Under the Receiver Operating Characteristic curve (AUROC) was 0.89 for lighter skin phototypes (Fitzpatrick scales I–III) but decreased to a much lower value of 0.82 in cases where patients had darker skin tones (Fitzpatrick scales IV–VI) (Fitzpatrick 1988) (Daneshjou, Vodrahalli and Novoa 2022). Even more concerning are the particular collapses in sensitivity that have been noted even in prestigious institutional setups. For instance, when tested using the Diverse Dermatology Images (DDI) scale, the DeepDerm algorithm, which was once praised for reaching the same level of accuracy as physicians, collapsed from a sensitivity of 0.69 in the case of light skin types to an extremely low 0.23 in the case of dark skin types (Esteva, Kuprel and Thrun 2017). Another algorithm – ModelDerm – demonstrated a similar decrease from 0.41 to 0.12 in sensitivity (Han, et al. 2020). Clinically, these flaws lead to higher instances of false negative outcomes, with malignant melanomas being falsely classified as benign lesions and increasing mortality among disadvantaged populations.

In addition to compromising imaging diagnosis tools, the issue of data disparity has a critical negative impact on text-based health algorithms. Risk assessment engines used throughout major health organizations have traditionally relied on expenses in healthcare systems as an indirect indicator to determine the medical needs of the patients. Since socioeconomic inequalities lead to decreased resource utilization by Black individuals, the algorithms calculated that equally sick Blacks were actually healthier than Whites, thus blocking hundreds of minority patients from access to intensive care coordination programs. In a similar fashion, digital diagnostic assistants can fail to diagnose myocardial infarctions among female patients, since the algorithm was trained on the basis of the classical presentation of symptoms common to men (Obermeyer, Powers and Mullainathan 2019). Unless proper regulations require balanced datasets, demographic recording, and minimum accuracy levels in subpopulations, the use of medical artificial intelligence can turn into the role of an invisible enemy rather than an equalizing tool.

4.2 The "Black Box" Problem and Liability

The greatest contradiction of contemporary medical artificial intelligence technology and law is that of the inherent lack of transparency of its computational process, known popularly as the "black box" issue. While classical medical algorithms make use of explicit logic defined by humans, the new generation of deep learning architectures, including multilayer CNNs and transformers, make decisions based on a set of probabilistic calculations carried out via millions of interconnected weight vectors. Due to their high complexity and non-linearity, the exact logical processes responsible for arriving at certain medical results are impossible to comprehend even by engineers, data

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

analysts, and physicians (DeGrave, Janizek and Lee 2023). This absence of explainable algorithms runs counter to the core principles of Western tort law, medical malpractice, and the ethics of informed consent, making the black-box technology an enemy of the legal system (Char, Shah and Magnus 2018).

The above-described absence of transparency creates a problem with the classical medical negligence approach that depends much on finding out a human fault and a linear cause-and-effect relationship. In a classic case of medical malpractice, the clinician is liable for deviation from the established “standard of care” resulting in harm to a patient. But when the “black box” Diagnostic CDSS is brought to the process of diagnosis, it creates what law professors describe as “the liability diffusion problem.” If an automated algorithm based on patients’ data calculates an authoritative risk assessment with an error and, as a result, a wrong diagnosis is made, pinpointing a particular point of malfunction becomes impossible. As it is impossible to decompile the network and prove its inherent defects, the question of whether the mistake came due to some software design defect, a corrupt local preprocessing procedure, or patient-specific pathophysiology is beyond the scope of the opaque algorithms’ functioning (Price 2017).

As such, today’s legal environment sets up a very punitive “liability trap” for frontline healthcare practitioners. As stated by reports from legal protection associations, under today’s legal framework, physicians become "liability sinks" whereby they carry full liability for any error committed by the machine without being technically capable of knowing how the machine arrived at its decision. The physician carries liability in both ways:

The Error of Commission: If a doctor blindly trusts an institutionalized, statistically validated AI recommendation that delivers a false negative (e.g., misclassifying an early-stage malignant tumor as a benign cyst), the doctor can be held fully liable for failing to independently override the system.

The Error of Omission: Conversely, if a clinician chooses to reject an AI recommendation based on intuitive human clinical judgment, and that rejection leads to a poor patient outcome, the doctor faces immediate allegations of professional negligence for failing to adhere to an authoritative, data-driven tool.

This imbalance exists due to the fact that the current laws relating to liability typically protect corporations involved in developing and supplying artificial intelligence software. Although medical equipment is covered by strict liability, medical software falls into the category of a legal gray area in which the software developer is not liable for any defects in the product (Price 2017).

Moreover, the black-box approach severely undermines the ethical principle of patient autonomy and informed consent. The average patient has a legally established right to be informed about the reasoning behind the chosen treatment course or surgical procedure. When the physician resorts to the black-box algorithm, there is no way for him/her to explain why the machine arrived at this conclusion, thus putting the therapeutic relationship under pressure of blind faith. The results of the McKinsey AI Trust Maturity Survey demonstrate the existence of a serious regulatory gap and highlight the "trust asymmetry" - even though 74% of the patients believe that they can trust the machine-generated diagnosis, 78% are mistaken in their assumption that the doctors can independently verify these diagnoses (Luget 2025). It will take some time for the legal framework to catch up and officially define such tools as strict liability products or demand that Explainable AI systems be implemented. Until then, the clinical practice becomes a legal minefield because of the unexplainable deep learning algorithms.

4.3 Data Security and Patient Privacy


However, the introduction of artificial intelligence into the biomedical stacks has drastically redefined the cybersecurity landscape in healthcare infrastructure globally. Whereas earlier EHRs were limited by institutional firewalls, the contemporary AI algorithms employed in medicine depend on constant and large amounts of data inflow in order to sustain accuracy and perform their analytics. The system-level need for data flows has led to a dramatic increase in cybersecurity threats in the industry, which was previously notoriously slow to change and is now the key focus of cyber warfare operations. Given that in healthcare systems there is a lot of valuable and unique patient identification data, the conjunction of growing AI-related attack surfaces and vulnerable network components has made data security a dangerous "foe."

An example of the macro-level criticality of this risk is provided by current statistics on institutional data breaches. It has been reported that health care continues to rank as the most expensive industry in terms of data breaches for 14 consecutive years, with the average monetary loss due to a single data security breach estimated at $7.42 million (IBM 2025). Criminal hackers actively use automation-based extortion and ransomware tactics, responsible for around 48% of all detected healthcare data infrastructure compromises (American Hospital Association 2026). Threat actors moved from simple credential-based attacks to using customized AI-powered tools scanning for unpatched software vulnerabilities shortly after their release, thus outpacing the manually performed patching procedure in institutions. Healthcare organizations continue to report cybersecurity workforce shortages and resource constraints, limiting their ability to monitor and defend increasingly complex digital environments. The increased vulnerability caused by such a situation is supplemented by the rise of so-called shadow AI, which refers to the practice of using unauthorized third-party clinical AI tools by healthcare employees.

Also, the impact on patient safety and mortality is among the operational impacts of these security breaches. It takes, on average, 279 days for any healthcare system to detect a breach and contain it completely. Within this span of undetected exploitation of the computer network of such a system, attackers penetrate into other operational components. According to studies across several centers, 72% of healthcare organizations that have been targeted by a significant cyber-attack experienced disruption in their patient services. When the network of a hospital is under

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

siege by a ransomware attack, the diagnostic process grinds to a halt, and the ambulances need to find new routes to deliver patients to clinics while certain procedures are put off. Statistics indicate that there is a strong connection between the number of such attacks and patient mortality. For instance, 29% of breached healthcare organizations experience increases in patient mortality as a result of these security attacks and subsequent operational disarray. Also, the stolen health information of a person carries a significant price tag in illicit digital markets, where one medical record can cost up to $1,250. Unlike the credit card data, one's personal information about health and genes cannot be changed (Ponemon Institute 2025).




As much as there are criminal attacks on networks, there is the moral and legal grey area of the unethical and illicit use of the commercialization of health information in order to create machine learning software. Machine learning systems need a lot of inputs in order to train, thus making the tech giants work hard to acquire data from institutions' databases. The healthcare analytics market rose to a whopping $64 billion in 2025, and a big part of that growth was attributed to the commercialization of patient data (Grand View Research 2025). The commodification of patient clinical records usually happens through intermediaries and external business associates who currently form more than 58% of health data breaches. In de-identification processes, clinical data is devoid of any explicit identifiers such as patients' names or their social security numbers in order to satisfy legacy regulatory framework exemptions such as HIPAA (Health Insurance Portability and Accountability Act (Public Law 104-191) 1996). Nevertheless, modern re-identifying data-mining techniques will find out who those people are from the voter records, consumer lists, and geolocation cache databases (Rocher, Hendrickx and de Montjoye 2019). As tough international regulations such as the EU AI Act become enforceable, the medical community has to implement strict API privacy measures and a zero-trust data architecture. Otherwise, data breaches might turn patient data into an insecure commercial asset, thus destroying the ethical sanctity of medical confidentiality.

5 Discussion & The Path Forward

5.1 The Symbiotic "Human-in-the-Loop" Framework

In order to address the inherent conflict of AI being a "friend" versus "foe," the biomedicine community should steer away from total autonomy of its technologies. The solution to the challenge lies in the structural implementation of the HITL paradigm. Under the HITL paradigm, artificial intelligence is not used as an independent and autonomous tool but rather as a modifier of diagnosis and administration with the help of cognitive copiloting. Thus, it will be recognized that despite the advantages of AI deep learning networks in terms of their superiority in terms of data-processing capabilities of parsing multiple imaging and high-frequency critical care data streams, AI still lacks the capability for contextual clinical thinking and empathic consideration of patients' conditions (Amann, Blasimme and Vayena 2020).



An effective HITL system would need to take into consideration one major cognitive flaw in human-computer interaction, that is, automation bias, where human beings tend to trust algorithm recommendations blindly and disregard their training. This can be combated by ensuring that the design of medical applications is not only static, binary output displays such as "Malignant" vs "Benign," but also interactive displays that convey information on confidence intervals, data limitations, and specific edge cases. By ensuring that the output of AI systems is displayed using structured probabilities and not absolute recommendations, the architecture ensures that the human clinicians retain their autonomy in making decisions. The physician is the final ethical and legal responsibility for patient care, and AI is merely used to augment the process.

To effectively operationalize this paradigm shift away from pure automation bias, upcoming clinical architectures must transition from black-box output presentation toward integrated post-hoc explainability metrics. Specifically, integrating feature-attribution methods like SHAP (SHapley Additive exPlanations) and LIME (Local Interpretable Model-agnostic Explanations) allows the AI interface to visually present which exact clinical features—such as an abrupt shift in serum lactate or a minor deviation in respiratory rate—weighted most heavily in generating a sepsis alert. Providing this layer of algorithmic transparency shifts the technology from a mysterious oracle to an interpretable interactive tool, protecting clinicians from liability traps while preserving genuine human-in-the-loop clinical autonomy.

5.2 Regulatory Medical AI Standards and Governance

The shift from chaotic deployment to the safe and regulated clinical space needs matching software engineering with the strict regulatory baseline. The regulatory space internationally experienced major changes that brought about regulatory baselines for the adoption of adaptive digital health technologies. In the United States, the Food and Drug Administration has successfully transitioned the existing framework to make the Quality Management Systems Regulation fully operational. The achievement

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

brings about direct matching of FDA pre-market regulation with the ISO 13485:2016 medical devices standard. The regulation requires coordination of AI engineering in three key pillars ((ISO) 2016):

Good Machine Learning Practices (GMLP): Ethical and procedural protocols governing dataset quality, curation, provenance, and multi-center model tuning.

Explainable AI (XAI) Risk Mitigations: Mandatory user interface specifications that force the software to transparently present its clinical reasoning pathways and training dataset subgroup metrics.

Predetermined Change Control Plans (PCCP): A crucial mechanism targeting the "death of the locked algorithm". A PCCP permits an adaptive model to continuously learn, retrain, and evolve within a pre-approved performance "playground" without requiring a brand-new regulatory filing for every minor iterative update (Singh, et al. 2026).

GLOBAL REGULATORY COMPLIANCE CORE ARCHITECTURE	
UNITED STATES (FDA)	**EUROPEAN UNION (EMA)**
QMSR (ISO 13485 Compliance)	EU AI Act Harmonization
Good Machine Learning (GMLP)	Dual-Layer MDR/IVDR Review
Pre-Approved PCCP Learning Bounds	Mandated Demographic Bias Logs

At the same time, the European Union has completed its historic Digital Omnibus Package, explaining how the integration of the EU AI Act will take place in conjunction with traditional sectoral legislation such as the Medical Device Regulation (MDR) and In-Vitro Diagnostic Medical Device Regulations (IVDR). This legally harmonized document explicitly categorizes stand-alone and integrated clinical AI systems as high risk. These regulations set out a series of horizontal requirements regarding algorithm validation, automated event tracking across the lifespan of the system, and demographic bias monitoring (Artificial Intelligence Act (EU Regulation 2024/1689) 2024).

Through the imposition of stringent consequences for model drift and bias in training data not being monitored, these globally recognized frameworks help hold technology vendors responsible for their products. Going forward, compliance cannot continue to be seen as a mere corporate tick-boxing process. Rather, robust algorithmic governance and consistent validation models need to become the bedrock upon which clinical safety can be established.

6 Conclusion

Incorporation of AI into the field of medicine represents a revolutionary moment in biomedical history, changing the everyday experience of care and management dramatically. The present review explored the dichotomy of medical artificial intelligence, proving that its designation as either clinical "friend" or "foe" does not arise from an intrinsic feature of the technology, but is rather a direct result of how it is designed, applied, and controlled. In its responsible use, AI proves to be an essential tool to amplify diagnostics and administrative efficiency by increasing the sensitivity of visual diagnostics in radiology, enabling ambient generative documentation for better clinical workflow, and providing life-saving warnings in critical care environments. However, unregulated use of AI creates new challenges, including perpetuation of historical health disparities through demographic bias of the database, establishment of a threatening legal "liability trap" due to the unclear "black-box" design of deep learning algorithms, and major security and privacy risks for patient data.

In conclusion, it is essential to reject the black and white paradigm of being either technophobic or blindly reliant on automated solutions. The future sustainability and safety of healthcare systems are fully dependent on the implementation of a symbiotic "human-in-the-loop" approach when AI technology acts as a cognitive co-pilot instead of being an autonomous substitute for human expertise. It is possible to avoid systematic risks associated with the use of AI technologies in health care by abiding by all the international regulations, providing explain ability of algorithms, and diversity of datasets. With proper governance and preservation of clinical autonomy, medical artificial intelligence will be firmly established as a tool that will change healthcare for the better.

References:

1. (ISO), International Organization for Standardization. "Medical devices — Quality management systems — Requirements for regulatory purposes (ISO 13485:2016)." 2016.
2. Amann, J, A Blasimme, and E Vayena. "Explainability for artificial intelligence in healthcare: A multidisciplinary perspective." BMC Medical Informatics and Decision Making 20 (2020): 310.
3. American Hospital Association. "Cybersecurity vulnerability report: Assessing the landscape of automated threat infiltration in healthcare networks." 2026.
4. "Artificial Intelligence Act (EU Regulation 2024/1689)." 2024.
5. Boussina, A. "Impact of a deep learning sepsis prediction model on quality of care and survival." npj Digital Medicine 7, no. 1 (2024): 15-24.
6. Char, D S, N H Shah, and D Magnus. "Implementing machine learning in health care — Addressing ethical challenges." New England Journal of Medicine 378, no. 11 (2018): 981-983.
7. Chen, T., & Guestrin, C. "XGBoost: A Scalable Tree Boosting System." Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining 5, no. 3 (2016): 785–794.
8. Daneshjou, R, K Vodrahalli, and R A Novoa. "Disparities in dermatology AI performance on a

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com

diverse, curated clinical image set." Science Advances 8, no. 32 (2022): eabq6147.
9. DeGrave, A J, J D Janizek, and S I Lee. "Dissection of Medical AI Reasoning Processes via Physician and Generative-AI Collaboration." Nature Biomedical Engineering 7, no. 10 (2023): 1201-1215.
10. Denham, D T, C Y Wang, E Maric, L R Hinton, and B T Heniford. "Large language models in surgery: Promise, pitfalls, and practical clinical use." Journal of Abdominal Wall Surgery 3, no. 2 (2026): 112-125.
11. Esteva, A, B Kuprel, and S Thrun. "Dermatologist-level classification of skin cancer with deep neural networks." Nature 542, no. 7639 (2017): 115-118.
12. Fitzpatrick, T B. "Skin type classification system." Archives of Dermatology 124, no. 6 (1988): 869-871.
13. Grand View Research. "Healthcare analytics market size, share & trends analysis report and segment forecasts, 2025-2030." 2025.
14. Han, S S, et al. "Augmented intelligence dermatology: deep neural networks empower medical professionals in diagnosing skin cancer and predicting treatment options for 134 skin disorders." Journal of Investigative Dermatology 140, no. 9 (2020): 1753-1761.
15. "Health Insurance Portability and Accountability Act (Public Law 104-191)." 1996.
16. Henry, K E, D N Hager, and S Saria. "TREWScore for septic shock prediction." Science Translational Medicine 7, no. 299 (2015): 299ra122.
17. IBM. "Cost of a data breach report 2025: Sectoral deep-dive on global healthcare infrastructure security." 2025.
18. "ISO 13485: Medical devices quality management systems." 2016.
19. Luget, A., G. Morgan, R. Roberts, B. Preston, and K. Ottenbreit. 2025. Insights on Responsible AI from the Global AI Trust Maturity Survey. McKinsey & Company / Tech Forward, 2025.
20. M. Razaghi, A. Hafez, J. M. Farina, I. G. Scalia, M. Pereyra, F. E. Abdelfattah, H. Sheashaa, K. Awad, S. J. Lester, C. Ayoub, and R. Arsanjani. "Transforming Clinical Documentation with Ambient Artificial Intelligence (AI) Scribes: A Narrative Review of Technology, Impact, and Implementation." Biomedical Informatics Insights 17, no. 1 (2026): 1-14.
21. McKinney, S M, M Sieniek, and S Shetty. "International evaluation of an AI system for breast cancer screening." Nature 577, no. 7788 (2020): 89-94.
22. Obermeyer, Z, B Powers, and S Mullainathan. "Racial bias in healthcare algorithms." Science 366, no. 6464 (2019): 447-453.
23. Olson, K D, and T D Shanafelt. "Use of Ambient AI Scribes to Reduce Administrative Burden and Professional Burnout." JAMA Network Open 8, no. 2 (2025): e2534976.
24. Ponemon Institute. "Cyber Insecurity in Healthcare: The Cost and Impact on Patient Safety and Care 2025." 2025.
25. Price, W N. "Regulating black-box medicine." Michigan Law Review 116, no. 3 (2017): 421-474.
26. Rocher, L, J M Hendrickx, and Y A de Montjoye. "Estimating the success of re-identifications in incomplete datasets using generative models." Nature Communications 10 (2019): 3069.
27. Shanafelt, T D, O Boone, and L N Dyrbye. "Physician burnout and work-life integration." Mayo Clinic Proceedings 97, no. 12 (2022): 2210-2224.
28. Shashikumar, S P, G Wardi, A Malhotra, and S Nemati. "Artificial intelligence sepsis prediction algorithm learns to say "I don't know"." Biocomputing 2025. 2024. 109-120.
29. Shashikumar, S P, G Wardi, S Nemati, and A Malhotra. "Development and multi-site validation of a deep learning model for early detection of sepsis in the emergency department." Nature Medicine 29, no. 11 (2023): 2834-2842.
30. Singer, M, C S Deutschman, and C W Seymour. "Sepsis-3 definitions." JAMA 315, no. 8 (2016): 801-810.
31. Singh, R, M Pareek, M Prokle, E Sanchez, U Hohgrawe, and J R Auclair. "Federating AI-related regulations for human therapeutics: An AI-enabled, continuously updating regulatory intelligence system." Frontiers in Drug Safety and Regulation 6, no. 1 (2026): 1718379.
32. Sinsky, C, L Colligan, and L Li. "EHR inefficiencies and "pajama time"." Annals of Internal Medicine 165, no. 11 (2016): 753-760.
33. Tan et al., J. Y. E., Rafi, I. B. M., Sng, G. G. R., Tung, J. Y. M., Lim, D. Y. Z., Ong, J. C. L., Lim, G. H., Lew, E. Y. L., Chia, K. W., & Heng, M. K. H. "Impact of an Ambient AI Scribe Among Clinicians and Patients: Real-World Prospective Observational Time-Motion Study." JMIR Medical Informatics 26, no. 2 (2026): e85580.
34. Titano, J J, M Badgeley, and E K Oermann. "Deep learning-based CT triage for acute findings." Nature Medicine 24, no. 9 (2018): 1337-1341.
35. Tschandl, P, C Rinner, and H Kittler. "HAM10000 dataset of dermatoscopic images." Scientific Data 5 (2018): 180098.
36. U.S. Food and Drug, Administration. "Artificial Intelligence and Machine Learning (AI/ML)-Enabled Medical Devices Tracker: Consolidated Data Report." 2025.
37. "US Healthcare Labor Market Survey and 2026 Workforce Shortage Forecast." 2024.
38. Zhang, X, and Y Yan. "Approval of AI-Based Medical Devices: A Global Retrospective Analysis and Registry Review." Journal of Medical Internet Research AI 5, no. 1 (2026): e85538.

Crossponding Author: Tel: +92 310 567 1025
Email: 512mughees@gmail.com