


Roles of long noncoding RNAs in gastric cancer and their clinical applications

Weiliang Sun¹ · Yunben Yang² · Chunjing Xu² · Yi Xie² · Junming Guo² 

Received: 22 November 2015 / Accepted: 19 May 2016 / Published online: 31 May 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract

Purpose Gastric cancer ranks as the most common cancer in the world. However, the progresses of its diagnosis and treatment are still not satisfactory. The purpose of this study is to summarize the roles of lncRNAs associated with gastric cancer.

Methods We searched lncRNAs associated with gastric cancer in PubMed.

Results Long noncoding RNAs (lncRNAs), transcripts larger than 200 nucleotides, regulate gene expression at various levels. They are playing important roles in the occurrence and development of gastric cancer. They are involved in signaling pathways, crosstalk with microRNAs, and affecting metastasis by regulating epithelial-to-mesenchymal transition. By acting as oncogenes or tumor suppressors, lncRNAs contribute to gastric cancer occurrence and development. Several lncRNAs including HOTAIR, HULC, LINC00152, MALAT2, H19, GHET1, and GACAT3 have been demonstrated having oncogene activities, while other lncRNAs including LEIGC, GAS5, and FERIL4 have been thought as tumor suppressors.

Conclusions Several lncRNAs from tissue, blood, and gastric juice have shown potential values in gastric cancer diagnosis or prognosis evaluation.

Keywords Long noncoding RNA · Biological function · Regulatory mechanism · Gene expression · Tumorigenesis · Clinical relevance

Introduction

Gastric cancer ranks the fifth most common malignant neoplasm in the world, with approximately 951,600 new cases diagnosed in 2012 (Ferlay et al. 2015). Despite the declined mortality rate of gastric cancer these years, it is still the third most common cause of cancer death (Li et al. 2015a; 2016). However, as low sensitivity and specificity, the commonly used tumor markers, such as carcinoembryonic antigen (CEA) and carbohydrate antigen 19-9 (CA 19-9), are not satisfied for the diagnosis of gastric cancer (Cui et al. 2013; Fang et al. 2015). As a result, it is urgently required to further explore the molecular mechanisms underlying gastric cancer occurrence and development.

Long noncoding RNAs (lncRNAs) are defined as transcripts larger than 200 nucleotides (Mercer et al. 2009; Wang and Chang 2011). LncRNAs regulate gene expression at various levels and play important roles in carcinogenesis (Gupta et al. 2010; Kung et al. 2013; Mercer et al. 2009). As signals, some lncRNAs first combine with proteins (such as hormone receptors and transcriptional factors); then the complex binds to the promoter site and regulates downstream gene expression (Fig. 1a). Moreover, to regulate gene expression, lncRNAs may serve as guides to lead regulatory proteins to the promoter site of targeted genes, as decoys to prevent regulatory proteins from binding with the promoter site, or as scaffolds to recruit protein complexes (Fig. 1b–d).

Increasing researchers have identified numbers of dysregulated expression of lncRNAs in gastric cancer and

✉ Junming Guo
guojunming@nbu.edu.cn

¹ Ningbo Yinzhou People's Hospital and the Affiliated Yinzhou Hospital, Ningbo University School of Medicine, 315040 Ningbo, China

² Zhejiang Key Laboratory of Pathophysiology, Department of Biochemistry and Molecular Biology, Ningbo University School of Medicine, 315211 Ningbo, China




Fig. 1 Roles of lncRNAs on gene expression by interacting with proteins. **a** To response the stimulus, an lncRNA serves as a signal molecule and combines with a protein (hormone receptor and transcriptional factor); then the lncRNA–protein complex regulates downstream gene expression. **b** An lncRNA guides a protein to the promoter site of the targeted gene. **c** An lncRNA serves as a decoy to prevent a protein from binding with the promoter site of the targeted gene. **d** An lncRNA acts as a scaffold to recruit protein complexes. lncRNA, long noncoding RNA

uncovered their potential roles in gastric cancer (Li et al. 2014). Functioned as oncogenes or tumor suppressors, lncRNAs participate in several vital steps of gastric cancer occurrence and development. They are involved in signaling pathways, crossing talk with microRNAs (miRNAs), and affecting metastasis by regulating epithelial-to-mesenchymal transition (EMT) process. In this article, we review the roles of lncRNAs in gastric cancer and their potential clinical applications.

Roles of lncRNAs in gastric cancer

lncRNAs regulate signaling pathways in gastric cancer

Studying the aberrantly expressed lncRNAs involved in signaling pathways may provide us new insights into the occurrence and development of gastric cancer.

H19, a maternal allele-specific expressed lncRNA, affects the proliferation of gastric cancer cells (Song et al. 2013). This property is associated with oncogene c-Myc and tumor suppressor gene P53 (Yang et al. 2012; Zhang et al. 2014). C-Myc is one member of Myc family, which shows its oncogenicity via adjusting the activities of histone acetyltransferase (HAT) or RNA polymerase II (RNAPII) and then regulates the expression of targeted genes (Amente et al. 2011). The up-regulated H19 levels in gastric cancer cell lines and gastric cancer tissues were first found by Yang et al. (2012). Then, to investigate the relationship between H19 and c-Myc, a plasmid expressing c-Myc was constructed by Zhang et al. (2014). The results demonstrated that H19 expression was induced by c-Myc. Through this mechanism, H19 promotes cell proliferation (Fig. 2a). The induced H19 by c-Myc demonstrates its oncogene property. Beside, H19's oncogene property can also be showed through inhibiting the activity of tumor suppressor gene P53. Using flow cytometry and RNA immunoprecipitation (RIP) assay, Yang et al. found that overexpression of H19 partially inactivated P53 and decreased the protein level of Bax, one of the P53's targets and a key




Fig. 2 lncRNAs involved in signaling pathways in gastric cancer. **a** lncRNAs associate with c-Myc. H19 is induced by c-Myc. The E-box element in the CCAT1 promoter can be activated by c-Myc binding, resulting in CCAT1 expression. GHET1 shows its tumorigenic function via interacting with both IGF2BP1 and c-Myc mRNA. **b** H19 reduces P53 activity and decreases the protein level of Bax. **c**

GAS5 suppresses protein levels of E2F1 and Cyclin D1 and increases P21 protein levels. CCAT1, colon cancer-associated transcript 1; E2F1, E2F transcription factor 1; IGF2BP1, IGF2 mRNA-binding protein 1; GAS5, growth arrest-specific 5; GHET1, gastric carcinoma high expressed transcript 1

apoptosis regulator (2012). By regulating P53, H19 contributes gastric cancer cells to resist apoptosis (Fig. 2b).

Similar to H19, colon cancer-associated transcript 1 (CCAT1) is another lncRNA that can be induced by c-Myc. A study showed that CCAT1 activated by c-Myc promoted progression of gastric cancer (Yang et al. 2013). The binding of c-Myc with E-box element in the CCAT1 gene promoter will activate CCAT1 expression (Fig. 2a).

In addition, gastric carcinoma high expressed transcript 1 (GHET1) is also associated with c-Myc. This lncRNA was found overexpressed in gastric cancer tissues comparing with normal tissues, while its deletion by small interference RNA (siRNA) blocked gastric cancer cell proliferation (Yang et al. 2014). Through gain-of-function and loss-of-function assays, GHET1 was identified as an upstream c-Myc controller (Yang et al. 2014). GHET1 plays its tumorigenic role via interacting with both IGF2 mRNA-binding protein 1 (IGF2BP1) and c-Myc mRNA. These processes maintain c-Myc mRNA stability (Fig. 2a).

Beside to regulate P53 (Fig. 2b), the tumor suppressor activity of lncRNAs in gastric cancer can be further showed by the regulation of growth arrest-specific 5 (GAS5) on retinoblastoma (RB) pathway (Fig. 2c). GAS5 is one of the down-regulated lncRNAs in gastric cancer (Sun et al. 2014). It inhibits cell proliferation and induces apoptosis through decreasing the protein levels of E2F transcription factor 1 (E2F1) and cyclin D1, two key participators in RB pathway. GAS5 also increases the activity of P21, a vital player of cell cycle arrest (Fig. 2c).

LncRNAs cross talk with miRNAs in gastric cancer

The activities of lncRNAs can be associated with miRNAs. On one hand, lncRNAs may be the precursor of miRNAs. For instance, miR-675, a product from H19 cleavage, plays a crucial role in tumorigenesis of gastric cancer (Zhuang

et al. 2014). It has been revealed that H19 promotes gastric cancer cell proliferation via modulating the expression of runt-related transcription factor 1 (RUNX1) and calneuron 1 (CALN1), two targets of miR-675 (Zhuang et al. 2014). On the other hand, lncRNAs may be the targets of miRNAs. Our group found that gastric cancer-associated transcript 3 (GACAT3) was one of the targets of miR-129-5p in gastric cancer cells (Xu et al. 2014). This lncRNA is up-regulated in gastric cancer and associates with poor prognosis of patients with gastric cancer (Xu et al. 2014).

Another important form of the relationships between lncRNAs and miRNAs is the competing endogenous RNA (ceRNA) network.

Salmena et al. (2011) proposed that miRNA response elements (MREs) built a bridge for the communication among various types of RNA, that is, the competing endogenous RNA (ceRNA) hypothesis. Acting as ceRNAs, lncRNAs may regulate the level of miRNAs' targets in post-transcriptional level. Notably, these cross talks between lncRNAs and miRNAs may play a significant role in tumorigenesis (Cheng et al. 2015; Karreth and Pandolfi 2013).

Based on our previous gastric cancer data of lncRNA microarray (Song et al. 2013), we firstly established an lncRNA-miRNA-mRNA network in gastric cancer (Xia et al. 2014). Lately, we found that both fer-1-like family member 4 (FER1L4, an lncRNA) and phosphatase and tensin homolog deleted on chromosome ten (PTEN) mRNA are targets of miR-106a-5p (Xia et al. 2015). We demonstrated that FER1L4 functions as a ceRNA to regulate the expression of PTEN by sequestering miR-106a-5p in gastric cancer. Since the fact that PTEN is a tumor suppressor gene, the suppressed expression of FER1L4 in gastric cancer will release more miR-106a-5p to inhibit the expression of PTEN, and thus affect cell cycle distributions (Fig. 3a).

Human epidermal growth factor receptor-2 (HER2), known as an oncogene, plays a role in proliferation and apoptosis.




Fig. 3 Two representative examples of lncRNAs acting as competing endogenous RNA (ceRNAs) in gastric cancer. **a** Acting as a ceRNA, FER1L4 regulates PTEN expression by shared miR-106a-5p. **b** Functioning as a ceRNA, HOTAIR regulates HER2 expression by

sequestering miR-331-3p. HER2, human epidermal growth factor receptor-2; PTEN, phosphatase and tensin homolog deleted on chromosome ten

It encodes a 185-kDa transmembrane protein to trigger the activation of cell signaling networks, impacting on various malignant cell functions such as proliferation, motility, angiogenesis, and apoptosis (Lemoine et al. 1991). Liu et al. found that there was a cross talk between the HOX antisense intergenic RNA (HOTAIR) and HER2 mRNA via competition for miR-331-3p (2014a). This correlation was confirmed by luciferase and RIP assays (Liu et al. 2014a). Liu et al. found that miR-331-3p can directly bind to both HOTAIR and HER2 mRNA through respective miRNA recognition sites (2014a). Furthermore, the effect of HOTAIR expression on endogenous HER2 protein in combination with modulation of miRNA or lncRNA levels was confirmed (Liu et al. 2014a). Western blot analysis also showed that forced expression of miR-331-3p or knockdown of HOTAIR triggered a significant silencing effect on endogenous HER2 protein expression (Liu et al. 2014a). HOTAIR acts as an endogenous ‘sponge’ by binding miR-331-3p, thus abolishing the miRNA-induced repressing activity on the HER2 mRNA (Fig. 3b). Overexpression of HOTAIR up-regulated the level of HER2 and contributed to gastric cancer occurrence (Fig. 3b).

Recently, more lncRNAs including maternally expressed 3 (MEG3) and BC032469 have been shown their characteristics as ceRNAs to promote proliferation in gastric cancer cells (Lü et al. 2015; Peng et al. 2015).

LncRNAs affect metastasis by regulating EMT in gastric cancer

EMT, a critical process in cancer metastasis, is known as the appearance of migratory potential in epithelial cells and adding the mesenchymal characteristics to the cells (Bonnomet et al. 2010; Creighton et al. 2013; Pasquier et al. 2015).

E-cadherin is regarded as a suppressor of tumor metastasis. Its low expression is one of markers of EMT. E-cadherin expression is inhibited by Snail. Several lncRNAs

regulate EMT by targeting E-cadherin and Snail (Fig. 4). Xu et al. (2013) found that HOTAIR facilitated EMT by regulating Snail in gastric cancer. They first found that overexpression of HOTAIR was closely related to metastasis. To further explore the role of HOTAIR in metastasis, they performed real-time reverse transcription-polymerase chain reaction (RT-PCR) and Western blotting. They found that the levels of matrix metalloproteinase-1 (MMP-1) and MMP-9 were suppressed after inhibition of HOTAIR via transfecting siRNA. Finally, morphological changes in gastric cancer cells indicated the reversion of EMT process when the deletion of HOTAIR. There was a decrease in the expression of mesenchymal markers (e.g., vimentin, N-cadherin), while there was an increase in the level in epithelial markers containing E-cadherin and ZO-1 (Xu et al. 2013). They also demonstrated that through ectopic expression of Snail, the invasion restrained by inhibition of HOTAIR in gastric cancer cells could be restored (Xu et al. 2013). Up-regulated HOTAIR contributed to the gastric cancer cell migration, invasion, and metastasis with decreasing expression of E-cadherin and ZO-1, yet increasing expression of N-cadherin and vimentin (Fig. 4).

Snail not only regulates the expression of EMT markers but also correlates with miRNAs. MiR-34a is one of downstream targets of HOTAIR (Liu et al. 2015). To determine whether HOTAIR regulates miR-34a expression levels by binding with polycomb repressive complex 2 (PRC2), Liu et al. (2015) used ENCODE histone modification tracks embedded in UCSC Genome Browser (UC Santa Cruz, CA, USA) and found H3K27me3 enrichment peaks in the miR-34a promoter region. Furthermore, RIP assays showed that HOTAIR could bind to PRC2, and the results of chromatin immunoprecipitation (ChIP) assays showed that enhancer of zeste homolog 2 (EZH2) could directly bind to the promoter region of miR-34a and mediate H3K27me3 modification (Liu et al. 2015). As a result, by recruiting




Fig. 4 Involvement of lncRNAs in EMT process. By recruiting the PRC2 complex, HOTAIR regulates the expressions of Snail and miR-34 and affects MMP-1 and MMP-9 levels. Other lncRNAs such as MALAT2, Linc00152, and HULC promote EMT; however, LEIGC suppresses EMT. EMT, epithelial-to-mesenchymal transition; HOTAIR, HOX antisense intergenic RNA; HULC, hepatocellular

carcinoma up-regulated long noncoding RNA; LEIGC, lower expression in gastric cancer; Linc00152, long intergenic non-protein coding RNA 152; MALAT2, metastasis-associated lung adenocarcinoma transcript 2; MMP-1, matrix metalloproteinase-1; MMP-9, matrix metalloproteinase-9; PRC2, polycomb repressive complex 2

the PRC2, HOTAIR down-regulates miR-34a expression (Fig. 4). Actually, Snail is one of the confirmed targets of miR-34a (Siemens et al. 2011). Therefore, the overexpression of HOTAIR silences miR-34a and then leads to a liberation of Snail that promotes EMT in gastric cancer (Fig. 4).

Another lncRNA named lower expression in gastric cancer (LEIGC) was identified for its down-regulation in gastric cancer tissues (Han et al. 2014). *In vivo* and *in vitro* experiments showed that LEIGC overexpression suppressed tumor growth and migration of gastric cancer cells, yet knockdown accumulated tumor progression (Han et al. 2014). The morphological change in LEIGC knockdown cells suggested its regulating EMT processes. By analyzing the gene expression profiles and performing Western blotting, Han et al. (2014) proved the mesenchymal cell-related genes (e.g., *zeb*, *twist*, *snail*, and *slug*) and their corresponding proteins increased when LEIGC was knockdown. These indicate that LEIGC is a key controller in preventing EMT procedure.

Other lncRNAs such as hepatocellular carcinoma up-regulated long noncoding RNA (HULC), long intergenic nonprotein coding RNA 152 (LINC00152), and metastasis-associated lung adenocarcinoma transcript 2 (MALAT2) have also been shown their features in contributing EMT in gastric cancer (Chen et al. 2015; Zhao et al. 2014, 2015).

The clinical relevance of lncRNAs in gastric cancer

Although the underlying mechanisms of most lncRNAs need to be further clarified, increasing evidences show that their expression levels are related to the clinicopathological features such as TNM stage, tumor size, metastasis, differentiation, and invasion of gastric cancer. These indicate the possible clinical applications of lncRNAs in the diagnosis of gastric cancer.

By using lncRNA microarray, our group identified 135 lncRNAs, whose differential expression levels between tumor and non-tumorous tissues were more than twofold (GEO No. GSE47850, <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE47850>). The most down-regulated lncRNAs include FER1L4, BG491697, AF131784, uc009ycs, uc001lsz, BG981369, AF147447, HMLincRNA1600, and AK054588, and most up-regulated lncRNAs contain BM709340, BQ213083, AK054978, DB077273, and H19 (Song et al. 2013). For their potential diagnostic values, we found that the positive detection rates of H19 and uc001lsz were higher than traditional biomarkers, CEA and CA19-9. The areas under receiver operating characteristic curve (AUC) of H19 and uc001lsz were up to 0.613 and 0.751, respectively (Song et al. 2013).

It was demonstrated that LINC00152, gastric cancer-associated transcript 2 (GACAT2, AC130710), small ubiquitin-like modifier (SUMO) 1 pseudogene 3 (SUMO1P3),

and ABHD11 antisense RNA 1 (ABHD11-AS1) were markedly up-regulated in gastric cancer tissues (Lin et al. 2014; Mei et al. 2013; Pang et al. 2014; Xu et al. 2014), while GACAT1 (AC096655.1-002), GACAT2 (HMLincRNA717), AA174084, FER1L4, and AI364715 were significantly down-regulated in gastric cancer tissues (Liu et al. 2014b; Shao et al. 2014a, b; Sun et al. 2013; Zhu et al. 2015). More important, as AC096655.1-002, HMLincRNA717, and AC130710 were firstly found to be closely related to gastric cancer by us, they have been officially nominated as gastric cancer-associated transcript 1 (GACAT1), GACAT2, and GACAT3, respectively (Chen et al. 2014; Xiao and Guo 2013).

GACAT1 was thought as one of the gastric-associated lncRNAs (Sun et al. 2013). Its level was significantly associated with lymph node metastasis, distant metastasis, TNM stages, and differentiation, suggesting lower GACAT1 expression along with poor differentiation and later stage indicating poor prognosis of patients with gastric cancer (Sun et al. 2013). Its AUC was up to 0.731, with the sensitivity and specificity being 0.513 and 0.872, respectively (Sun et al. 2013).

GACAT2's expression level was negatively associated with gastric cancer invasion and metastasis (Shao et al. 2014a). It was down-regulated in 62.6 % gastric cancer tissues compared to the paired adjacent normal tissues (Shao et al. 2014a). Contrary to GACAT2, the expression level of GACAT3 was significantly up-regulated in gastric cancer tissues (Xu et al. 2014). Higher level of GACAT3 indicates poor prognosis of patients with gastric cancer. Its level was significantly correlated with tumor size, distant metastasis, and TNM stages (Xu et al. 2014).

Based on above findings and other reports, in which the altered expression of lncRNAs in gastric cancer tissues was found correlated with clinicopathological features of patients with gastric cancer, many groups considered using circulating lncRNAs to screen gastric cancer. It may also be a novel way to assess the prognostic values by comparing circulating lncRNAs levels between pre- and postoperative patients with gastric cancer. For instance, two weeks after surgery, plasma FER1L4 level in gastric cancer patients was found sharply decreased (Liu et al. 2014b). By investigating AA174084 level in 335 plasma samples, we found that it was markedly dropped on 15 days after surgery and that the change was associated with lymphatic metastasis and invasion (Shao et al. 2014b). Furthermore, by comparing LINC00152 levels between plasma and exosomes from blood and observed by transmission electron microscopy, we found that one possible mechanism underlying circulating lncRNAs' stably existing in peripheral blood was their protection by exosomes (Li et al. 2015b). Recently, Dong et al. (2015) reported that the combinative use of three circulating lncRNAs may provide a superior diagnostic value

with an AUC reaching 0.92 and sensitivity and specificity being up to 74.1 and 100 %, respectively.

Due to its specificity, that is, only existing in stomach, we and other researchers take an interest in the use of gastric juice lncRNAs to screen gastric cancer. We first reported that LINC00152 level in gastric juice from patients with gastric cancer was significantly higher than those from healthy people (2014). Our further study showed that gastric juice AA174084, with an AUC of 0.848, provided a power diagnostic value than circulating lncRNAs (Shao et al. 2014b). Importantly, our group showed that using gastric juice ABHD11-AS1 as a biomarker increased the positive detection rate of early gastric cancer to 71.4 % (Yang et al. 2015). Other studies found that gastric juice lncRNAs including LINC00982 and urothelial cancer-associated 1 (UCA1) also have the potential clinical values for the diagnosis of gastric cancer (Fei et al. 2015; Zheng et al. 2015).

Conclusion

With the development of microarray and high-throughput screening as well as RT-PCR, thousands of cancer-associated lncRNAs have been found. In this review, we emphasized those gastric cancer-associated lncRNAs involved in signaling pathways, interacting with miRNAs, and regulating EMT. Some of them such as HOTAIR, HULC, LINC00152, MALAT2, H19, GHET1, and GACAT3 act as oncogenes. Others such as LEIGC, GAS5, and FER1L4 have the property of tumor suppressor genes. Many of them might be used in gastric cancer diagnosis or prognosis evaluation.

Funding This study was funded by the Zhejiang Provincial Natural Science Foundation of China (No. LY14C060003), the Applied Research Project on Nonprofit Technology of Zhejiang Province (No. 2016C33177), the National Natural Science Foundation of China (No. 81171660), and the K. C. Wong Magna Fund in Ningbo University.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

Amente S, Lania L, Majello B (2011) Epigenetic reprogramming of Myc target genes. *Am J Cancer Res* 1(3):413–418

Bonnomet A, Brysse A, Tachsidis A, Waltham M, Thompson EW, Polette M et al (2010) Epithelial-to-mesenchymal transitions and circulating tumor cells. *J Mammary Gland Biol Neoplasia* 15(2):261–273

Chen S, Li P, Xiao B, Guo J (2014) Long noncoding RNA HMLincRNA717 and AC130710 have been officially named as gastric cancer associated transcript 2 (GACAT2) and GACAT3, respectively. *Tumour Biol* 35(9):8351–8352

Chen F, Tian Y, Pang EJ, Wang Y, Li L (2015) MALAT2-activated long noncoding RNA indicates a biomarker of poor prognosis in gastric cancer. *Cancer Gene Ther*. doi:10.1038/cgt.2015.6

Cheng DL, Xiang YY, Ji LJ, Lu XJ (2015) Competing endogenous RNA interplay in cancer: mechanism, methodology, and perspectives. *Tumour Biol* 36(2):479–488

Creighton CJ, Gibbons DL, Kurie JM (2013) The role of epithelial-mesenchymal transition programming in invasion and metastasis: a clinical perspective. *Cancer Manag Res* 5:187–195

Cui L, Zhang X, Ye G, Zheng T, Song H, Deng H et al (2013) Gastric juice MicroRNAs as potential biomarkers for the screening of gastric cancer. *Cancer* 119(9):1618–1626

Dong L, Qi P, Xu MD, Ni SJ, Huang D, Xu QH et al (2015) Circulating CUDR, LSINCT-5 and PTENP1 long noncoding RNAs in sera distinguish patients with gastric cancer from healthy controls. *Int J Cancer* 137(5):1128–1135

Fang XY, Pan HF, Leng RX, Ye DQ (2015) Long noncoding RNAs: novel insights into gastric cancer. *Cancer Lett* 356(2 Pt B):357–366

Fei ZH, Yu XJ, Zhou M, Su HF, Zheng Z, Xie CY (2015) Upregulated expression of long non-coding RNA LINC00982 regulates cell proliferation and its clinical relevance in patients with gastric cancer. *Tumour Biol*. doi:10.1007/s13277-015-3979-9

Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M et al (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136(5):E359–E386

Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ et al (2010) Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature* 464(7291):1071–1076

Han Y, Ye J, Wu D, Wu P, Chen Z, Chen J et al (2014) LEIGC long non-coding RNA acts as a tumor suppressor in gastric carcinoma by inhibiting the epithelial-to-mesenchymal transition. *BMC Cancer* 14:932

Karreth FA, Pandolfi PP (2013) ceRNA cross-talk in cancer: when cebling rivalries go awry. *Cancer Discov* 3(10):1113–1121

Kung JT, Colognori D, Lee JT (2013) Long noncoding RNAs: past, present, and future. *Genetics* 193(3):651–669

Lemoine NR, Jain S, Silvestre F, Lopes C, Hughes CM et al (1991) Amplification and overexpression of the EGF receptor and c-erbB-2 proto-oncogenes in human stomach cancer. *Br J Cancer* 64(1):79–83

Li PF, Chen SC, Xia T, Jiang XM, Shao YF, Xiao BX, Guo JM (2014) Non-coding RNAs and gastric cancer. *World J Gastroenterol* 20(18):5411–5419

Li P, Chen S, Chen H, Mo X, Li T, Shao Y et al (2015a) Using circular RNA as a novel type of biomarker in the screening of gastric cancer. *Clin Chim Acta* 444:132–136

Li Q, Shao Y, Zhang X, Zheng T, Miao M, Qin L et al (2015b) Plasma long noncoding RNA protected by exosomes as a potential stable biomarker for gastric cancer. *Tumour Biol* 36(3):2007–2012

Li T, Mo X, Fu L, Xiao B, Guo J (2016) Molecular mechanisms of long noncoding RNAs on gastric cancer. *Oncotarget* 7(8):8601–8612

Lin X, Yang M, Xia T, Guo J (2014) Increased expression of long noncoding RNA ABHD11-AS1 in gastric cancer and its clinical significance. *Med Oncol* 31(7):42

Liu XH, Sun M, Nie FQ, Ge YB, Zhang EB, Yin DD et al (2014a) Lnc RNA HOTAIR functions as a competing endogenous RNA to regulate HER2 expression by sponging miR-331-3p in gastric cancer. *Mol Cancer* 13:92

- Liu Z, Shao Y, Tan L, Shi H, Chen S, Guo J (2014b) Clinical significance of the low expression of FER1L4 in gastric cancer patients. *Tumour Biol* 35(10):9613–9617
- Liu YW, Sun M, Xia R, Zhang EB, Liu XH, Zhang ZH et al (2015) LincHOTAIR epigenetically silences miR34a by binding to PRC2 to promote the epithelial-to-mesenchymal transition in human gastric cancer. *Cell Death Dis* 6:e1802
- Lü MH, Tang B, Zeng S, Hu CJ, Xie R, Wu YY et al (2015) Long noncoding RNA BC032469, a novel competing endogenous RNA, upregulates hTERT expression by sponging miR-1207-5p and promotes proliferation in gastric cancer. *Oncogene*. doi:10.1038/ncr.2015.413
- Mei D, Song H, Wang K, Lou Y, Sun W, Liu Z et al (2013) Up-regulation of SUMO1 pseudogene 3 (SUMO1P3) in gastric cancer and its clinical association. *Med Oncol* 30(4):709
- Mercer TR, Dinger ME, Mattick JS (2009) Long non-coding RNAs: insights into functions. *Nat Rev Genet* 10(3):155–159
- Pang Q, Ge J, Shao Y, Sun W, Song H, Xia T et al (2014) Increased expression of long intergenic non-coding RNA LINC00152 in gastric cancer and its clinical significance. *Tumour Biol* 35(6):5441–5447
- Pasquier J, Abu-Kaoud N, Al Thani H, Rafii A (2015) Epithelial to mesenchymal transition in a clinical perspective. *J Oncol* 2015:792182
- Peng W, Si S, Zhang Q, Li C, Zhao F, Wang F et al (2015) Long non-coding RNA MEG3 functions as a competing endogenous RNA to regulate gastric cancer progression. *J Exp Clin Cancer Res* 34:79
- Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP (2011) A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell* 146(3):353–358
- Shao Y, Chen H, Jiang X, Chen S, Li P, Ye M et al (2014a) Low expression of lncRNA-HMlincRNA717 in human gastric cancer and its clinical significances. *Tumour Biol* 35(10):9591–9595
- Shao Y, Ye M, Jiang X, Sun W, Ding X, Liu Z et al (2014b) Gastric juice long noncoding RNA used as a tumor marker for screening gastric cancer. *Cancer* 120(21):3320–3328
- Siemens H, Jackstadt R, Hunten S, Kaller M, Menssen A, Gotz U et al (2011) miR-34 and SNAIL form a double-negative feedback loop to regulate epithelial-mesenchymal transitions. *Cell Cycle* 10(24):4256–4271
- Song H, Sun W, Ye G, Ding X, Liu Z, Zhang S et al (2013) Long non-coding RNA expression profile in human gastric cancer and its clinical significances. *J Transl Med* 11:225
- Sun W, Wu Y, Yu X, Liu Y, Song H, Xia T et al (2013) Decreased expression of long noncoding RNA AC096655.1-002 in gastric cancer and its clinical significance. *Tumour Biol* 34(5):2697–2701
- Sun M, Jin FY, Xia R, Kong R, Li JH, Xu TP et al (2014) Decreased expression of long noncoding RNA GAS5 indicates a poor prognosis and promotes cell proliferation in gastric cancer. *BMC Cancer* 14:319
- Wang KC, Chang HY (2011) Molecular mechanisms of long noncoding RNAs. *Mol Cell* 43(6):904–914
- Xia T, Liao Q, Jiang X, Shao Y, Xiao B, Xi Y et al (2014) Long non-coding RNA associated-competing endogenous RNAs in gastric cancer. *Sci Rep* 4:6088
- Xia T, Chen S, Jiang Z, Shao Y, Jiang X, Li P et al (2015) Long non-coding RNA FER1L4 suppresses cancer cell growth by acting as a competing endogenous RNA and regulating PTEN expression. *Sci Rep* 5:13445
- Xiao B, Guo J (2013) Long noncoding RNA AC096655.1-002 has been officially named as gastric cancer-associated transcript 1, GACAT1. *Tumour Biol* 34(5):3271
- Xu ZY, Yu QM, Du YA, Yang LT, Dong RZ, Huang L et al (2013) Knockdown of long non-coding RNA HOTAIR suppresses tumor invasion and reverses epithelial-mesenchymal transition in gastric cancer. *Int J Biol Sci* 9(6):587–597
- Xu C, Shao Y, Xia T, Yang Y, Dai J, Luo L et al (2014) lncRNA-AC130710 targeting by miR-129-5p is upregulated in gastric cancer and associates with poor prognosis. *Tumour Biol* 35(10):9701–9706
- Yang F, Bi J, Xue X, Zheng L, Zhi K, Hua J et al (2012) Up-regulated long non-coding RNA H19 contributes to proliferation of gastric cancer cells. *FEBS J* 279(17):3159–3165
- Yang F, Xue X, Bi J, Zheng L, Zhi K, Gu Y et al (2013) Long noncoding RNA CCAT1, which could be activated by c-Myc, promotes the progression of gastric carcinoma. *J Cancer Res Clin Oncol* 139(3):437–445
- Yang F, Xue X, Zheng L, Bi J, Zhou Y, Zhi K et al (2014) Long non-coding RNA GHET1 promotes gastric carcinoma cell proliferation by increasing c-Myc mRNA stability. *FEBS J* 281(3):802–813
- Yang Y, Shao Y, Zhu M, Li Q, Yang F, Lu X et al (2015) Using gastric juice lncRNA-ABHD11-AS1 as a novel type of biomarker in the screening of gastric cancer. *Tumour Biol*. doi:10.1007/s13277-015-3903-3
- Zhang EB, Han L, Yin DD, Kong R, De W, Chen J (2014) c-Myc-induced, long, noncoding H19 affects cell proliferation and predicts a poor prognosis in patients with gastric cancer. *Med Oncol* 31(5):914
- Zhao Y, Guo Q, Chen J, Hu J, Wang S, Sun Y (2014) Role of long non-coding RNA HULC in cell proliferation, apoptosis and tumor metastasis of gastric cancer: a clinical and in vitro investigation. *Oncol Rep* 31(1):358–364
- Zhao J, Liu Y, Zhang W, Zhou Z, Wu J, Cui P et al (2015) Long non-coding RNA Linc00152 is involved in cell cycle arrest, apoptosis, epithelial to mesenchymal transition, cell migration and invasion in gastric cancer. *Cell Cycle* 14(19):3112–3123
- Zheng Q, Wu F, Dai WY, Zheng DC, Zheng C, Ye H et al (2015) Aberrant expression of UCA1 in gastric cancer and its clinical significance. *Clin Transl Oncol* 17(8):640–646
- Zhu S, Mao J, Shao Y, Chen F, Zhu X, Xu D et al (2015) Reduced expression of the long non-coding RNA AI364715 in gastric cancer and its clinical significance. *Tumour Biol* 36(10):8041–8045
- Zhuang M, Gao W, Xu J, Wang P, Shu Y (2014) The long non-coding RNA H19-derived miR-675 modulates human gastric cancer cell proliferation by targeting tumor suppressor RUNX1. *Biochem Biophys Res Commun* 448(3):315–322

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com