

## Review

## Crosstalk between kinases, phosphatases and miRNAs in cancer



Júlia L.F. Abrantes, Thaís F. Tornatore, Karin J. Pelizzaro-Rocha, Marcelo B. de Jesus, Rodrigo T. Cartaxo, Renato Milani, Carmen V. Ferreira-Halder\*

Department of Biochemistry, Institute of Biology, UNICAMP, 13083-970 Campinas, Brazil

## ARTICLE INFO

## Article history:

Received 18 February 2014

Accepted 4 September 2014

Available online 16 September 2014

## Keywords:

Protein tyrosine kinases

Protein tyrosine phosphatases

miRNA

Cell signaling

Cancer aggressiveness

Chemotherapy resistance

## ABSTRACT

Reversible phosphorylation of proteins, performed by kinases and phosphatases, is the major post translational protein modification in eukaryotic cells. This intracellular event represents a critical regulatory mechanism of several signaling pathways and can be related to a vast array of diseases, including cancer. Cancer research has produced increasing evidence that kinase and phosphatase activity can be compromised by mutations and also by miRNA silencing, performed by small non-coding and endogenously produced RNA molecules that lead to translational repression. miRNAs are believed to target about one-third of human mRNAs while a single miRNA may target about 200 transcripts simultaneously. Regulation of the phosphorylation balance by miRNAs has been a topic of intense research over the last years, spanning topics going as far as cancer aggressiveness and chemotherapy resistance. By addressing recent studies that have shown miRNA expression patterns as phenotypic signatures of cancers and how miRNA influence cellular processes such as apoptosis, cell cycle control, angiogenesis, inflammation and DNA repair, we discuss how kinases, phosphatases and miRNAs cooperatively act in cancer biology.

© 2014 Elsevier B.V. and Société française de biochimie et biologie Moléculaire (SFBBM). All rights reserved.

## 1. Introduction

Phosphorylation and dephosphorylation are the main type of post-translational modifications in eukaryotic cells, and are tightly regulated by protein kinases and protein phosphatases, respectively [1]. Indeed, phosphorylation and dephosphorylation are efficient strategies to modulate cellular response towards internal and external stimuli, since it is a reversible process that does not require synthesis and/or degradation of proteins. Therefore, these post-translational modifications allow cells to transiently regulate a set of proteins at the same time, in different cellular compartments and signaling pathways that in turn, define which metabolic response will be dominant under a specific stimulus. Under this context, a variety of cellular processes, such as proliferation, migration, apoptosis, autophagy, differentiation, metabolism, organelle trafficking, immunity, learning and memory depend on the action of kinases and phosphatases [2–4]. Thus, not surprisingly, abnormal kinase or phosphatase activities correlate with disease conditions such as cancer, diabetes and neurodegenerative or inflammatory disorders [5–7].


Specifically in relation to miRNAs metabolism and function, there are some evidences that kinases and phosphatases can be modulated and modulate miRNAs. Recent studies have proposed that oncogenic miRNAs can downregulate the expression of tumor suppressor kinases and phosphatases, which in turn contributes for tumor progression. The interplay between protein kinases, phosphatases and miRNAs in cancer cells is very complex and depends of many features, such as cell type and tumor context, as well as the pattern of expression of tumor suppressor and oncogenic miRNAs. Therefore, it is clear that the complex crosstalk between protein kinases, phosphatases and miRNAs can contribute to increase cancer aggressiveness, giving tumor cells the ability to survive, increase proliferation, evade cell death routines and metastasize. This review will highlight what is already known about the interplay between miRNAs, protein kinases and phosphatases and its importance in cancer. Besides, we also highlight the potential of those molecules as targets for more efficient cancer therapy.

## 1.1. Post translational modification: role of protein kinases and phosphatases

About 2% of the human genome correspond to protein kinases, encoding 518 protein kinases, which constitute one of the largest gene families in eukaryotes [3,8,9]. Protein kinases (PKs) catalyze the phosphorylation on serine, threonine or tyrosine residues.

\* Corresponding author. Tel.: +55 1935216659.

E-mail addresses: [verissimac@gmail.com](mailto:verissimac@gmail.com), [carmenv@unicamp.br](mailto:carmenv@unicamp.br) (C.V. Ferreira-Halder).



**Fig. 1.** Biogenesis of miRNA. Pre-miRNAs generated by the canonical or non-canonical pathways are transported by an exportin 5 and RAN-GTP-dependent process from the nucleus to the cytosol. The precursors are further processing by the Dicer and transactivation-response RNA-binding protein (TRBP) RNase III enzyme complex to form the mature double-stranded miRNA. Next, argonaute proteins facilitate incorporation of the mature miRNA-targeting strand into the AGO-containing RNA-induced silencing complex (RISC). The RISC-miRNA assembly is then guided to specific target sequences in mRNAs resulting either in translational repression or decreased mRNA stability.

Thus, based on the nature of the phosphorylated group, kinases are classified as protein-serine/threonine kinases (385 members), protein-tyrosine kinases (90 members), and tyrosine-kinase like proteins (43 members) [3,10].

By contrast, the human genome encodes only approximately 200 phosphatases, targeting phosphorylated proteins or lipids. Protein phosphatases (PPs) catalyze the hydrolysis of the phosphate group from its protein substrate. Thus, PPs are categorized by

structure and substrate specificity into protein serine/threonine phosphatases (PSTPs, approximately 30 members) and protein tyrosine phosphatases (PTPs, 107 members) [10,11].

## 1.2. Biogenesis and function of miRNAs

MiRNAs (miRNAs) are small non-coding RNA molecules that contain around 19–30 nucleotides (on average 20 nucleotides) and are produced by the canonical or non-canonical biogenesis pathways [12]. miRNAs generated by the canonical biogenesis pathway are synthesized as precursor RNAs from intergenic, intronic or polycistronic genomic loci by RNA polymerase II. Subsequently, Drosha and DGCR8 RNase III complex recognize and process the stem-loop structure of primary miRNA (pri-miRNA) transcript. In contrast, non-canonical miRNA biogenesis directly produces miRNAs as endogenous short hairpin RNAs or derives directly through splicing from introns that can refold into hairpins (mirtrons). Exportin 5 transports both precursors (pre-miRNA and mirtrons) (~60–100 nucleotides) from the nucleus to the cytosol in a Ran–GTP dependent manner. Subsequently, the precursors are processed by a complex consisting of Dicer and transactivation-response RNA-binding protein (TRBP) RNase III enzymes, to form the mature double-stranded miRNA. Next, argonaute proteins facilitate incorporation of the mature miRNA–targeting strand into the AGO-containing RNA-induced silencing complex (RISC). The RISC–miRNA assembly is then guided to specific target sequences in mRNAs. The initial recognition of mRNAs by the RISC–miRNA complex is driven primarily by base-pairing of nucleotides 2 to 8 in the mature miRNA with specific mRNA target sequences located in the 3′ untranslated region (3′UTR), and additional base-pairing affords greater affinity and targeting efficiency. The RISC–miRNA complex acts either in translational repression or decreasing mRNA stability [12] – (Fig. 1). However, in some cases miRNAs can enhance mRNA translation. For instance, it was reported that miR-10a binds to 5′UTR of ribosomal protein mRNAs and enhanced their translation [13].

In general miRNAs are believed to target about one-third of human mRNAs and a single miRNA is capable of targeting approximately 200 transcripts simultaneously [14–16]. Therefore, miRNAs have been implicated in almost every biological process, including development, cell cycle regulation, cell growth and differentiation, stress response, and apoptosis. In addition, miRNAs play a role in a variety of diseases, particularly in cancer [17].

## 1.3. miRNAs and cancer

In addition to maintaining the normal growth, development, and health of an organism, miRNAs can also influence the development and progression of malignancy. Since around 50% of human miRNAs are located in fragile chromosomal regions, which may exhibit DNA amplifications, deletions or translocations during tumor development, their expression is frequently deregulated in cancer [18,19]. Besides, it has been observed that cellular transformation and tumorigenesis can be promoted by suppression of key components of miRNA processing machinery. Widespread downregulation of miRNAs is often detected in human cancers such as breast, ovarian and prostate cancer. In general, miRNAs can function either as tumor suppressors or as oncogenes (oncomirs), initiating tumor growth, invasion, metastases, as well as regulating the overall *stemness* of cancer cells [20].

## 1.4. miRNA as endocrine signals

Very recent papers in the literature have reported that miRNAs can be detected in the human plasma, suggesting possibilities for

novel disease biomarkers and the intriguing notion that some miRNAs may serve regulatory purposes in target cells. In the plasma, miRNAs can be found as part of microvesicles and exosomes derived from donor plasma membranes cells, in association with the miRNA-processing enzyme argonaute 2 and in lipoproteins [21]. It has been proposed that vesicles (microvesicle/exosomes) containing miRNAs are delivered to target cells either by a process involving endocytosis or by membrane fusion [21]. Importantly, it also has been shown that circulating miRNAs might rather act as endocrine or paracrine signaling molecules to deliver a regulatory signal from donor to target cells. These mobile miRNAs, defined as *hormomirs* due to their hormone-like characteristics, could act as local or long-range signals to maintain normal homeostasis or influence the development and progression of diseases such as cancer. Different research groups have isolated cell-free circulating miRNAs from blood and demonstrated that the levels of certain miRNAs changed in response to various tumors, which pointed out these molecules as non-invasive biomarkers for different tumor entities [22]. However, the functional significance of circulating miRNAs is still not clear, and this might be, at least in part, due to non-reproducibility of data, as Jarry and collaborators have addressed in a review [23].


## 2. Influence of protein kinases and protein phosphatases in miRNAs biogenesis

### 2.1. Regulation of Drosha and Dicer stability and processing activity

Following the discovery of miRNAs and their potential significance to orchestrate the whole cellular machinery, a renewed understanding shed lights to a more comprehensive and straightforward view towards the dynamic balance of miRNAs generation. The primordial and limiting step in miRNAs biogenesis relies on the Drosha/DGCR8 holoenzyme. The translocation of Drosha to the nucleus is known to be activated by GSK3 $\beta$ , through a specific phosphorylation at the N-terminal nuclear localization domain [24]. In the nucleus Drosha selects precursor transcripts to efficiently produce a global stream of functionally relevant miRNAs *in vivo*. Therefore, the selectivity of Drosha gives insights into why, of the hundreds of thousands of potential miRNAs, only a subset is ever produced by cells [25].

As mentioned before, following translocation into the cytoplasm, the pre-miRNA is cleaved near the terminal loop by the RNase III domain of Dicer to generate a 22-nucleotide double strand mature miRNA. Several Dicer-associated proteins have been identified, including TAR RNA-binding protein (TRBP) and protein kinase R-activating protein (PACT). Dicer stability and activity are enhanced by association with TRBP and PACT [26] and depletion of these cofactors decreases the steady state levels of Dicer protein [27,28]. Furthermore, cell-signaling pathways can also modulate the expression or activity of Dicer. For instance, MAPK/ERK signaling was shown to promote the phosphorylation of TRBP [29]. Phosphorylated TRBP increases miRNA production by increasing the stability of Dicer. Besides, the activating oncogenic mutation *Braf*<sup>V600E</sup> (upstream mediator of MAPK cascade) amplifies the processing of pri-miRNAs into mature miRNAs, overcoming even the loss of Dicer expression induced by haploinsufficiency in primary mice sarcomas. Comparison with the opposite mutation in *Kras* genes reinforced that deficient MAPK activity leads to decreased steady-state levels of mature miRNAs, which has been associated with tumor progression and development of metastases [30].

Protein tyrosine phosphatase also seems to take part in miRNA biogenesis. For instance, chemoresistant chronic myeloid leukemia cells with reduced expression levels of low molecular weight protein tyrosine phosphatase (LMW-PTP) exhibit lower expression of



**Fig. 2.** Protein expression analysis of Dicer expression levels in two chronic myeloid leukemia cell lines. Lucena, derived from K562, is chemoresistant through over-expression of P-glycoprotein. Anita, derived from Lucena is a stable knockdown of LMW-PTP accomplished using shRNA in stable transfection with lentiviral particles.

Dicer. This can be observed at both mRNA and protein levels (Fig. 2). We have previously reported that LMW-PTP contributes for CML resistance [31]. However, it is still not clear if there is a connection between this phosphatase, miRNA synthesis and CML resistance.

## 2.2. miRNA decay regulation

Despite some miRNA biogenesis regulatory mechanisms having been described, the control of mature miRNAs life span through degradation has received less attention. miRNAs appear as highly stable molecules, since mature miRNAs have been found to persist for many hours or even days after their production [32–34]. Besides, mature miRNAs have been also found to be expressed in a tissue- or stage-specific manner without variation in the expression pattern of the precursor forms (pri- and pre-miRNAs), suggesting a regulatory mechanism acting on the mature miRNAs [35]. Some metabolic conditions such as cell cycle, viral infection, exposure to growth factor and endoplasmic reticulum stress have been shown to affect miRNA stability [36–38]. However, whether kinases and/or phosphatases are able to participate and mediate miRNAs degradation remain to be investigated in details, but those proteins may have crucial roles in this process. For instance, it has been shown that the exposure of breast and brain tumors to growth factor, such as epidermal growth factor (EGF), causes a fast variation on specific miRNAs and transcription factor levels. Indeed, EGFR, a tyrosine kinase receptor, activation in those cells provoked a decrease of the amount of 23 miRNAs, and consequently, rapid induction of oncogenic transcription factors [38]. Another enzyme that has been shown to play an important role in controlling miRNAs life span is the endoplasmic reticulum transmembrane kinase-endonuclease (IRE1 $\alpha$ ). IRE1 $\alpha$  is a multifunctional protein that consists of three important regions, an N-terminal sensor domain located inside of endoplasmic reticulum (ER) that detects misfolded proteins, a transmembrane region, and a cytosolic tail containing two distinct catalytic activities: a serine/threonine kinase and an endonuclease (RNase). Under ER stress, when there is an increase of misfolded proteins, an oligomerization with subsequent autophosphorylation of IRE1 $\alpha$  can be observed. This, in turn, allosterically activates its RNase activity [39,40]. Upton et al. [41] have reported that sustained IRE1 $\alpha$  RNase activation caused rapid decay of select miRNAs (miRs -17, -34a, -96, and -125b) that normally repress translation of Caspase-2 mRNA. This, consequently, elevates Caspase-2 concentration and favors the mitochondrial apoptotic pathway. Besides, in cell-free systems, recombinant IRE1 $\alpha$  endonucleolytically cleaved miRNA precursors at sites distinct from Dicer, which indicates that IRE1 $\alpha$  regulates

translation of a proapoptotic protein through terminating miRNA biogenesis [41].

## 2.3. Phosphatidylinositol 3-kinase (PI3K) modulates the level of exportin

PI3K is involved in cellular functions such as cell growth, proliferation, differentiation, motility, survival and intracellular trafficking. Therefore, when dysfunctional, PI3K displays a pivotal role in cancer progression. This enzyme produces the main positive modulator of AKT, PtdIns(3,4,5)P<sub>3</sub>. Recently, Iwasaki et al. [42] have compared the miRNA expression profiles between resting and antigen-activated lymphocytes and also in primary mouse embryonic fibroblast (MEF). In activated T-cell population an elevation of a broad numbers of miRNA was observed, and this profile was similar in MEF cells. Interestingly, in the same study the authors showed that exportin 5 was induced during cell cycle entry by a PI3K-dependent post-transcriptional mechanism. In addition, the inhibition of exportin 5 expression interfered with global miRNA elevation and resulted in a proliferation defect associated with delayed G1/S transition. Iwasaki et al. [42] also suggested that exportin 5-mediated global miRNA elevation might be involved in a broad range of cellular events associated with cell cycle control.

## 2.4. Phosphatase of regenerating liver-3 (PRL-3) upregulates the expression of miRNAs

PRL-3 a member of the classic protein tyrosine phosphatase family has received significant attention because of its involvement in colorectal cancer (CRC) metastasis. High expression of PRL-3 was observed especially in metastatic lesions derived from primary CRC, regardless of the target organ [43]; however, its expression was low or absent from normal colorectal epithelia, adenomas and primary lesions [44]. Zhang et al. [45] reported that PRL-3 upregulates the expression of miR-21, miR-17 and miR-19a in both colon cancer cells and tissues by activating signal transducer and activator of transcription 3 (STAT3) via the Csk-Src-STAT3 pathway. These data suggest a connection between PRL-3 and tumor metastasis.

## 3. Modulation of protein kinases by miRNAs

Several oncogenic and tumor suppressor kinases have been shown to be targeted by miRNAs in human cancers (Table 1). The altered expression of these proteins can result in some typical features of tumor cells, including resistance to apoptosis and an improved ability of migration and invasion. Some of these kinases and phosphatases, as well as the miRNAs described as their regulators, will be presented and their role in tumor progression will also be discussed. A summary of the main downstream signaling pathways altered by these proteins and the subsequent cellular responses is presented in Figs. 3 and 4.

### 3.1. Cell cycle kinases

In this section we addressed kinases that regulate the cell cycle progression and miRNAs that target them. The majority of those miRNAs function as tumor suppressors and they are inactivated in cancer by different mechanisms (reviewed by Bueno and Malumbres, 2011 [46]). For instance, miR-124 and miR-137 are silenced by hypermethylation in different human tumor cells which lead to CDK6 overexpression [47].

#### 3.1.1. Cyclin-dependent kinases (CDKs)

CDKs are serine/threonine kinases that play a key role in cell cycle regulation. Specific CDKs are activated in different phases of

**Table 1**  
Regulation of protein kinases by miRNAs in cancer.

Kinase	Examples of substrates	Regulatory miRNAs	Protein data	Altered miRNA expression
CDK4	Cdt1, Marcks, p107, p130, retinoblastoma, SMAD3	miR-34 [53], miR-195 [54] and miR-506 [199]	Overactive in human cancers due to genetic and epigenetic alterations [52]	Ectopic expression of miR-506 in ovarian cancer cells inhibits proliferation [199]
CDK6	Bcl2, Histone H1, retinoblastoma	miR-22 [55], miR-29 [56], miR-34 [57], miR-107 [58,59], miR-124a [60], miR-125b [61], miR-129 [62], miR-137 [63], miR-186 [64], miR-191 [65], miR-195 [66], miR-218 [67], miR-449a [68] and miR-494 [69]	Overactive in human cancers [52]	Expressive tumor growth inhibition and survival improvement were observed in mice bearing TP53-mutated multiple myeloma xenografts treated with miR-34a mimics [200]
CDK2	CDK7, E2F, p27, retinoblastoma, p53	miR-186 [64], miR-372 [70] and miR-885-5p [71]	Overactive in human cancers [52]	Ectopic expression of miR-372 suppressed cell growth and induced arrest in the S/G <sub>2</sub> phases of cell cycle in HeLa cells [70]
CDK1	Casein kinase 2, CDK7, APC	miR-16 (Takeshita et al., 2010), miR-223 [72], miR-410 [73], miR-650 [73], miR-1699, miR-1744 and miR-1798 [72]	Overactive in human cancers [52]	miR-16 significantly diminished the growth of prostate tumors in bone metastasis model [201]
Aurora	Histone H3, Aurora B, INCENP (CPC subunit), Ndc80, KNL1, Dsn1 and vimentin [202]	miR-24 [76] and let-7a [77]	Aberrant expression of Aurora kinases can disturb checkpoint functions leading to genetic instability and cancer onset. Overexpression of Aurora kinases has been detected in several cancer types and correlates to a poor prognosis [75]	Downregulation of let-7 has been detected as a common feature in several cancers, such as breast, colon, and lung cancers [78–80]; overexpression of let-7a inhibited the growth of HeLa cells by targeting Aurora kinase B [77]
Cdc7	MCM2-7 complex, ORC 1L [85]	miR-29a [88]	Cdc7 is overexpressed in human cancer cell lines and many primary tumors [86]. When overexpressed Cdc7 enhances tumor cell survival by repairing stalled replication forks [87]	Ectopic expression of miR-29a in lung cancer cells prevented Cdc7 accumulation in response to BPDE resulting in altered checkpoint signaling and increased cell death [88]
PLK1	Cdc25C, anaphase-promoting complex (APC), abnormal spindle (Asp) a microtubule-associated protein [93]	miR-10b [104], miR-100 [105,106] and miR-593 [107]	Overexpressed in cancers, including breast cancer [90], ovarian cancer [91], endometrial carcinomas [92], pancreatic cancer [93], renal carcinoma [94], colorectal cancer [95] and melanoma [96]	Ectopic expression of miR-10b decreased PLK1 expression and inhibited breast cell proliferation <i>in vitro</i> and tumor growth <i>in vivo</i> [104]. In naso-pharyngeal cancer cells transfected with miR-100 mimics exhibited reduced PLK1 expression and underwent growth inhibition, G2/M cell cycle arrest and apoptosis [106]. Synthetic miR-593 suppressed PLK1 expression in EC cells and reduced cell proliferation and increased the number of cells in the G2/M phase [107]
Wee1	cyclin B-CDK1 complex [108]; histone H2B [109]	miR-128a miR-155, miR-516a-3p [110] and miR-497 [111]	Highly active in several types of cancer, especially those ones that display p53 deficiency [203]	miR-128a, miR-155 and miR-516a-3p were shown to be overexpressed in nonfunctional pituitary adenomas [110]. Ectopic expression of these miRNAs in HeLa cells inhibited Wee1 expression, which was intriguingly accompanied by a decrease in cell proliferation [110]. siRNA knockdown of Wee1 in neuroblastoma cells increased apoptosis, while the overexpression of miR-497 reduced cell viability, increased apoptosis and sensitize neuroblastoma cells to treatment with cisplatin [111]
ATM	Chk1, Chk2 [52,56], histone H2A, p53 and Mdm2			Overexpression of miR-101 was shown to sensitize tumor cells to

(continued on next page)



Table 1 (continued)

Kinase	Examples of substrates	Regulatory miRNAs	Protein data	Altered miRNA expression
		miR-101 [115], miR-100 [114], miR-421 [116]	ATM gene alteration is a frequent event in pathogenesis of chronic lymphocytic leukemia [204]	radiation <i>in vitro</i> and <i>in vivo</i> by targeting ATM kinase [115]. Hu and colleagues [116] showed that HeLa cells overexpressing miR-421 were not able to block DNA synthesis in the S-phase of the cell cycle (S-phase checkpoint) in response to DNA damage. This effect was mediated by ATM downregulation and contributed to increase radiosensitivity in HeLa cells [116]
ATR	Chk1, Chk2 [52,56], HDAC2, p53 [205,206]	miR-185	Upregulated in Oral squamous cell carcinoma [207]	miR-185 was shown to be downregulated by ionizing radiation in renal cell carcinoma tissues and cell line [117]. Overexpression of miR-185 sensitized renal carcinoma cells to radiation both <i>in vitro</i> and <i>in vivo</i> [117]
MST2	YAP (Yes-associated protein) and TAZ (co-activator with PDZ-binding motif) [123,124]	miR-133b [120]	Overexpressed in human cervical carcinomas [120]	
LATS2	Aurora A	miR-31 [127], miR-372 [128,129] and miR-373 [128,130]	Tumor suppressor through modulating different processes such as cell proliferation and cell death ([121–123])	miR-31 was found to be overexpressed in murine and human lung cancers cells, and the knockdown of miR-31 increased LATS2 expression and repressed growth and tumorigenicity of these cells [127]. In human gastric cancer cells the inhibition of miR-372 by using antisense miR-372 oligonucleotide increased LATS2 expression, induced cell cycle arrest, suppressed proliferation and increased apoptosis of these cells [129]. miR-203 is silenced in CML; ectopic expression of miR-203 reduces ABL1 and BCR-ABL1 expression and inhibits cell proliferation [131]. Overexpression of miR-30a in K562 leukemia cells reduced ABL1 and BCR-ABL1 protein expression, proliferation, and blocked cell cycle progression between G1 and S [133]
BcrAbl	Src, Stat5 [208], uracil DNA glycosylase UNG2 [208,209]	miR-203 [131], miR-29b [132] and miR-30a [133]	hallmark in chronic myelogenous leukemia	Enhanced expression of miR-31 provokes an inhibition of the oncogenic NFκB pathway in breast [135]. Ectopic expression of miR-302-367 cluster in cervical carcinoma cells (HeLa and SiHa cells) inhibited cell proliferation and tumor formation [138]. Tao et al. [139] have shown that miR-612 inhibited hepatocellular carcinoma cells proliferation, migration, invasion, and metastasis through direct targeting AKT2. miR-99a was found to be downregulated in renal cell carcinoma (RCC) and its restoration suppressed RCC cell growth, migration and invasion, as well as induced cell cycle arrest <i>in vitro</i> . Besides, intratumoral delivery of miR-99a inhibited tumor growth in murine xenograft model of human RCC [143]
PKC	AKT1, ATF2, BAD, CX43, IRS1, MET, PKC epsilon and STAT3 [203]	miR-31 [135]	Highly activated in cancers [210]	
Akt	<b>Akt1</b> Akt1, eIF4B, H2B, HSP27, vimentin, YB-1, Wee1, PTP1B, PGC-1 alpha, PAR-4, p21, p27, mTor, Mdm2, lamin A/C, IRS1 [211] <b>Akt2</b> Substrates: Ask1, Cttnb1, ER alpha, Ezrin, histone 3, IKK alpha, PGC-1 alpha and XIAP [211]	miR-302-367 cluster regulates AKT1 expression [138], while miR-612 was related to AKT2 repression [139]	Aberrant AKT signaling has been frequently documented in human cancers and appears to play a pivotal role in their progression [137]	
mTOR	4E-BP1, Akt1, IRS-1, mTor, Myc, p70S6K, PTPN13, Stat3 and Ulk1 [212]	miR-99a [141,143] and miR-144 [144]	mTOR signaling is frequently deregulated in human cancers [142]	

Table 1 (continued)

Kinase	Examples of substrates	Regulatory miRNAs	Protein data	Altered miRNA expression
Fyn	Abl, Bcr, CDK5, CTNNB1, Fyn, H3 histone, IRS1, LKB1, SHP2 [213]	miR-125a-3p [146]	Its overexpression is associated with several types of cancer, such as glioblastoma multiforme, squamous cell carcinoma of the head and neck, and melanoma [145]	Elevated levels of miR-125a-3p causes a dramatic decrease of Fyn expression [146]
Fak	Jak2, paxilin, integrin beta5, Stat1, Pten, Src, Fyn and IRS1 [214,215]	miR-7 [150] [149]	Increased FAK expression is frequently observed in human tumors and has been associated with an increased malignant phenotype and poor prognosis [148,149]	Overexpression of miR-7 in U87 and U251 glioblastoma cells inhibited their invasiveness and migration activity [150]. Besides, the overexpression of miR-7 in breast cancer cells suppressed proliferation, anchorage independent growth, three-dimensional growth in matrigel, migration and invasion ( <i>in vitro</i> and <i>in vivo</i> models [149]
Pyk2	PTPN11, PTPN6, Src, Fyn, paxillin [214,215]		Highly activated in cancers [216]	Overexpression of miR-23b reduced PYK2 protein expression and significantly reduced glioma cell migration and invasion [151]. Ectopic expression of miR-517a and miR-517c inhibited HCC cells proliferation by blocking the G2/M transition [152]
Pak	aurora A, Bad, histone 3, Ilk, Mek1, Pak1, Raf and vimentin [153,154]. Substrates PAK2: Abl, ERK3, Jun, Mek1, Myc, Pak2 [153,154]	miR-7 targets PAK1 [157], miR-23b targets PAK2 [158] and miR-145 targets PAK4 [159]	PAKs have been related to cell transformation that results in tumor formation and cell invasiveness [153,154]	Overexpression of miR-7 in breast cancer cells inhibited motility, invasiveness, anchorage-independent growth and tumorigenic potential [157]
Rock	Ezrin, FAK, H3, IRS1, PTEN, radixin, vimentin, Rock2	miR-146a [161], miR-148b [162], miR-335 [163], miR-584 [164] and miR-1280 [165] was shown to target ROCK1, whereas miR-124 [166], miR-138 [167] and miR-139 [168] targeted ROCK2.	ROCK is frequently implicated in oncogenic transformation [160]	It was found a correlation between knockdown of ROCK kinases by the cited microRNAs and reduction of cell migration and invasion.
IGF1R	phospholipase C-gamma, mdm2, PTP1B, Src	miR-7 [171], miR-16 [172], miR-122 [173], miR-181b [174], miR-376a, miR-376c [175] and miR-497 [113,176]	Overexpressed in glioma patients [174], tongue squamous cell carcinoma cells [171], breast cancer [173], melanoma [175] and cervical carcinoma cells [176]	Shi et al. [174] have reported that miR-181b inhibited glioma proliferation, migration and invasion by targeting IGF1R and its downstream signaling pathways, PI3K/AKT and MAPK/ERK1/2. Ectopic expression of miR-7 in tongue squamous cell carcinoma (TSCC) cells reduced IGF1R expression and AKT activation, which in turn resulted in decreased cell proliferation and enhanced apoptosis [171]. miR-122, downregulated in breast cancer, also functions as a tumor suppressor <i>in vivo</i> , through targeting IGF1R and regulating PI3K/AKT/mTOR/p70S6K pathway [173]. In addition, miR-376a and miR-376c overexpression led to a significant decrease in melanoma cells migration by targeting IGF1R [175]. Luo et al. [176] have shown that ectopic expression of miR-497 in cervical carcinoma cells suppressed migration and invasiveness of these cells by

(continued on next page)

Table 1 (continued)

Kinase	Examples of substrates	Regulatory miRNAs	Protein data	Altered miRNA expression
EGFR	Ago2, Crk, EGFR, Ezrin, Fak, PTP1B	miR-7 [179–181], miR-27a [182,183] and miR-133b [184]	EGFR signaling has emerged as a major therapeutic target in many cancers, since its deregulation have been associated to tumor cell proliferation, apoptosis resistance, invasion, angiogenesis and metastasis [179]	post-transcriptionally targeting IGF1R Ectopic expression of miR-7 decreased viability and invasiveness of primary glioblastoma lines [180]
Axl	phospholipase Cgamma1 (PLCgamma), p85alpha and p85beta subunits of phosphatidylinositol 3'-kinase (PI3-kinase), c-src, Ick [217]	miR-34a and miR-199a/b	overexpression of AXL has been reported in several cancer types, such as colon, esophageal, thyroid, breast, lung, liver and astrocytoma-glioblastoma carcinomas [185]	Ectopic overexpression of these microRNAs decreased cell migration potential.
c-Met	c-Met, Fak, Shc1 [218,219]	miR-27a [182], miR-34a [191], miR-34c [192], miR-101 [193], miR-137 [194], miR-409-3p [195], miR-410 [196], miR-449a [197] and miR-7515 [198]	Deregulated in human cancers. The overexpression of c-MET and elevated levels of circulating HGF have been frequently detected in many cancers, and are associated with poor clinical outcomes [189]	

the cell cycle through phosphorylation and binding to activator proteins named cyclins [48]. The mammalian genome has twelve loci encoding CDKs, but only five of them (CDK1, CDK2, CDK3, CDK4 and CDK6) have been implicated in the regulation of cell cycle [49].

Cell cycle progression through the G1 phase is regulated by a complex mechanism involving CDK4, CDK6 and CDK2. Mitogenic signals result in increased transcription of D-type cyclins, which in turn form active complexes with CDK4 and CDK6. Then, these active kinases partially phosphorylate and inactivate the Retinoblastoma protein (pRb) allowing cyclin E synthesis. Cyclin E binds and activates CDK2, and the active complex CDK2-Cyclin E is responsible for further phosphorylation and inactivation of pRb [50]. Once phosphorylated, pRb releases E2F transcription factor allowing the transcription of genes necessary for entry into S phase [51]. CDK1 is activated by binding to A- or B-type cyclins and its activation is an essential step for mitotic entry through phosphorylation of numerous cytoplasmic and nuclear target proteins [49].

CDKs are frequently overactive in human cancers due to genetic and epigenetic alterations that affect their regulatory pathways. For this reason, CDKs have been considered as important targets for anticancer therapies [52]. Several miRNAs can influence cell cycle progression through targeting CDKs. Among them, miR-34 [53] and miR-195 [54] have been shown to target 3'UTR of CDK4. CDK6 was shown to be a direct target of miR-22 [55], miR-29 [56], miR-34 [57], miR-107 [58,59], miR-124a [60], miR-125b [61], miR-129 [62], miR-137 [63], miR-186 [64], miR-191 [65], miR-195 [66], miR-218 [67], miR-449a [68] and miR-494 [69]. Besides, CDK2 has been described as a target of miR-186 [64], miR-372 [70] and miR-885-5p [71], whereas CDK1 was found to be a target of miR-223 [72], miR-410 [73], miR-650 [73], miR-1699, miR-1744 and miR-1798 [72].

### 3.1.2. Aurora kinases

Aurora kinases are conserved serine–threonine kinases that participate as key regulators of mammalian cell division. Aurora kinase B is a chromosomal passenger protein that participates in the spindle checkpoint and chromosome function. Besides, it has been related to the prevention of chromosome instability [74]. Therefore, aberrant expression of Aurora kinases can disturb

checkpoint functions leading to genetic instability and cancer onset. The overexpression of Aurora kinases has been detected in several cancer types and correlates to a poor prognosis [75].

Two miRNAs were detected as regulators of Aurora kinase B expression, miR-24 [76] and let-7a [77]. The downregulation of let-7 has been detected as a common feature in several cancers, such as breast, colon, and lung cancers [78–80]. Liu et al. [77] showed that the level of let-7a is reduced in endometrial carcinoma cells, and the overexpression of this miRNA inhibited the growth of HeLa cells by targeting Aurora kinase B.

### 3.1.3. Cdc7 kinase


Cdc7 is a conserved serine/threonine kinase essential for the initiation of DNA replication [81,82]. Several studies have shown that Cdc7 participates in the response to DNA damage occurred during S-phase of the cell cycle [83–85], contributing to the maintenance of genomic stability. Because of that, it is expected that an altered control of Cdc7 could improve tumorigenesis. In fact, many reports have shown that Cdc7 is overexpressed in human cancer cell lines and many primary tumors [86]. In addition, it has been shown that the overexpression of Cdc7 enhances tumor cell survival by repairing stalled replication forks [87].

miR-29a was described as a regulator of Cdc7 expression by direct targeting 3'-UTR of this kinase and Cdc7 upregulation correlates with a downregulation of miR-29a [88]. miR-29a is downregulated in response to DNA damage induced by benzo[a]pyrene dihydrodiol epoxide (BPDE), a genotoxin present in cigarette smoke. Ectopic expression of miR-29a in lung cancer cells prevented Cdc7 accumulation in response to BPDE resulting in altered checkpoint signaling and increased cell death. Therefore, the loss of miR-29a might be a mechanism by which lung cancer cells acquire resistance to BPDE-induced DNA damage, allowing them to survive, proliferate and accumulate mutagenic lesions [88].

### 3.1.4. Polo-like kinases (PLKs)

PLKs comprise a highly conserved family of serine/threonine kinases involved in mitotic entry, centrosome maturation, spindle assembly and cytokinesis. PLK1 is the best-characterized member of this family and it is responsible for the phosphorylation of





**Fig. 3.** Cell cycle control by miRNAs. Some interactions were omitted for clarity. The main protein kinases and phosphatases involved in the cell cycle control and discussed above are represented in the figure. miRNAs that control their expression are presented inside blue frames.

CDC25C, which in turn activates CDK1–Cyclin B complex inducing mitotic entry [89]. PLK1 is frequently overexpressed in cancers, including breast cancer [90], ovarian cancer [91], endometrial carcinomas [92], pancreatic cancer [93], renal carcinoma [94], colorectal cancer [95] and melanoma [96]. PLK1 overexpression contributes to oncogenesis not only by inducing proliferation, but also by promoting chromosome instability [97]. When PLK1 is inhibited tumor cells undergo mitotic arrest and apoptosis, besides promoting tumor growth inhibition *in vivo* [98–103].

MiR-10b [104], miR-100 [105,106] and miR-593 [107] have been proposed to target PLK1 mRNA. miR-10b was shown to be a master regulator of breast cancer cell proliferation and is downregulated in this cancer [104]. Ectopic expression of miR-10b decreased PLK1 expression and inhibited breast cell proliferation *in vitro* and tumor growth *in vivo* [104]. In naso-pharyngeal cancer cells, miR-100 was shown to be downregulated contributing to PLK1 overexpression and tumor progression [105]. In non-small cell lung cancer (NSCLC) miR-100 was also significantly downregulated, which was correlated with an advanced tumor stage, lymph node metastasis and reduced overall survival of patients [106]. NSCLC cells (A549) transfected with miR-100 mimics exhibited reduced PLK1 expression and underwent growth inhibition, G2/M cell cycle arrest and apoptosis [106]. miR-593, another miRNA that targets 3'UTR of PLK1, was shown to be less expressed in esophageal cancer (EC) cells [107]. The introduction of synthetic miR-593 in EC cells suppressed PLK1 expression, reduced cell proliferation and increased the number of cells in the G2/M phase [107].

### 3.1.5. Wee1 kinase

Wee1 is an evolutionarily conserved nuclear tyrosine kinase protein that prevents cell entry into mitosis by negatively modulating the cyclin B–CDK1 complex through phosphorylation of the amino acids Tyr15 and Thr14 of CDK1 [108]. Besides, it has been shown that Wee1 directly phosphorylates the mammalian core


histone H2B at Tyr37 at the end of S phase, which is crucial for maintaining the correct histone–DNA stoichiometry prior to mitotic entry [109].

Wee1 has been identified as a direct target of miR-128a miR-155, miR-516a-3p [110] and miR-497 [111]. miR-128a, miR-155 and miR-516a-3p were shown to be overexpressed in nonfunctional pituitary adenomas [110]. Ectopic expression of these miRNAs in HeLa cells inhibited Wee1 expression, which was intriguingly accompanied by a decrease in cell proliferation [110].

miR-497 has been proposed to function as a tumor suppressor in several cancers [112,113], and its expression was shown to be reduced in neuroblastomas [111]. In addition, Wee1 was identified as a target of miR-497, and increased Wee1 expression was associated with poor prognosis of neuroblastoma. The siRNA knock-down of Wee1 in neuroblastoma cells increased apoptosis, while the overexpression of miR-497 reduced cell viability, increased apoptosis and sensitize neuroblastoma cells to treatment with cisplatin [111].

### 3.1.6. Ataxia-Telangiectasia Mutated kinase (ATM) and ATM and Rad 3-related kinase (ATR)

DNA Damage Checkpoint prevents cells from undergoing DNA replication or mitosis when DNA is damaged, providing an opportunity for DNA repair. In the presence of damaged DNA, a complex mechanism is triggered in mammalian cells to sense and transduce the signal to tumor suppressor pathways [48]. ATM and ATR are serine–threonine kinases members of PI3K superfamily that participates in sensing DNA damage. While ATM primarily responds to DNA double-strand breaks, caused by ionizing radiation and radiomimetic drugs, ATR also responds to damage caused by ultraviolet light and stalled replication forks [52]. These kinases phosphorylate CHK1 and CHK2, which in turn phosphorylate key proteins involved in DNA repair, cell cycle arrest and apoptosis. The major targets phosphorylated by CHKs are p53, responsible for cell



**Fig. 4.** Overview of the main signaling pathways controlled by tumor suppressor and oncogenic miRNAs. Some interactions were omitted for clarity. Inside blue frames are presented miRNAs that target kinases and phosphatases represented in the figure. After binding to specific ligands, the activated receptors c-MET, EGFR, IGF1R and AXL can activate key downstream pathways responsible for cell proliferation, growth, survival, motility and invasion. The major activated pathways include PI3K/AKT and MAPK/ERK. The kinase Fyn can be activated by RTKs and integrins and regulates PI3K/AKT, MAPK/ERK and also FAK signaling. PAK kinase can be activated by Rho GTPases, PI3K and integrins.

cycle arrest and apoptosis induction, and protein phosphatase CDC25, responsible for CDKs inhibition and cell cycle arrest [48,52].

ATM was previously identified as a target of miR-100 [114], miR-101 [115] and miR-421 [116]. The overexpression of miR-101 was shown to sensitize tumor cells to radiation *in vitro* and *in vivo* by targeting ATM [115]. The same effect of radiosensitization was shown for miR-100 overexpression in glioma cells [114]. Hu and colleagues [116] showed that HeLa cells overexpressing miR-421 were not able to block DNA synthesis in the S-phase of the cell cycle (S-phase checkpoint) in response to DNA damage. This effect was mediated by ATM downregulation and contributed to increase radiosensitivity in HeLa cells [116].

ATR has been recently identified as a target of miR-185 [117]. In renal cell carcinoma (RCC) tissues and cell line the expression of miR-185 was shown to be downregulated by ionizing radiation [117]. On the other hand, overexpression of miR-185 sensitized RCC cells to radiation both *in vitro* and *in vivo* [117].

### 3.2. Cell signaling kinases

#### 3.2.1. Mammalian STE20-like protein kinase (MST) and large tumor suppressor kinase (LATS)

STE20 and LATS are key players in the Hippo signaling pathway, recently described as a new relevant pathway to cancer development [118,119]. The core of this pathway comprises two sets of serine/threonine kinases, MST1 (mammalian STE20-like protein

kinase 1) and MST2, and the LATS1 (large tumor suppressor 1) and LATS2, together with the adaptor proteins SAV1, MOB1A (MOB kinase activator 1A) and MOB1B. Upon activation by stimuli such as cell–cell contact, the Hippo pathway kinase cascade phosphorylates the oncoproteins YAP (Yes-associated protein) and TAZ (co-activator with PDZ-binding motif). The phosphorylation of YAP/TAZ inhibits their activity and stimulates their ubiquitination and degradation. When dephosphorylated YAP/TAZ migrate to the nucleus, and modulate the activity of transcription factors such as TEADs and SMADs, thereby inducing cell proliferation and reducing apoptosis [118,119].

MST2 was shown to be downregulated by miR-133b which was found to be overexpressed in human cervical carcinomas [120]. Using the luciferase reporter assay, Qin and colleagues [120] confirmed 3'UTR of MST2 as a direct target of miR-133b. The silencing of MST2 in epidermoid cervical carcinoma (CaSki) cells using a specific siRNA enhanced cell proliferation and increased the number of colonies formed in soft agar.

LATS2, classically identified as a downstream target of MST2 on the Hippo pathway, act as a tumor suppressor through modulating different processes such as cell proliferation and cell death [121–123]. The hypermethylation of LATS2 gene and consequently, a reduction in LATS2 expression has been frequently reported in different cancers [124–126], and is related to a more aggressive tumor phenotype. Some studies have identified LATS2 mRNA as a direct target of miR-31 [127], miR-372 [128,129] and miR-373

[128,130]. miR-31 was found to be overexpressed in murine and human lung cancers cells, and the knockdown of miR-31 increased LATS2 expression and repressed growth and tumorigenicity of these cells [127]. In human gastric cancer (AGS) cells the inhibition of miR-372 using antisense miR-372 oligonucleotide increased LATS2 expression, induced cell cycle arrest, suppressed proliferation and increased apoptosis of these cells [129]. The down-regulation of LATS2 by miR-373 was first described by Voorhoeve and colleagues [128] in testicular germ cell tumors, and afterward the same interaction was proposed by Lee and colleagues [130] in esophageal cancer cell lines. In esophageal squamous cell carcinomas the level of LATS2 protein expression is inversely correlated with miR-373 expression [130].

### 3.2.2. BCR-ABL1 kinase

BCR-ABL oncoprotein, a non-receptor tyrosine kinase, is expressed from the Philadelphia chromosome upon the t(9;22)(q34;q11) chromosomal translocation and it is considered as the hallmark in chronic myelogenous leukemia (CML). BCR-ABL1 is a direct target of the miRNA miR-203. It is silenced by genetic and epigenetic mechanisms in CML, resulting in the overexpression of ABL1 and BCR-ABL1, while restoration of miR-203 expression reduces ABL1 and BCR-ABL1 expression and inhibits cell proliferation [131]. More recently, it was shown that miR-29b [132] and miR-30a [133] targeted BCR-ABL1 and promoted its downregulation in bone marrow from CML patients. In K562 leukemia cells, the overexpression of miR-29b inhibited cell growth, colony formation ability and induced apoptosis [132]. Similarly, the overexpression of miR-30a in K562 leukemia cells reduced ABL1 and BCR-ABL1 protein expression, proliferation, and blocked cell cycle progression between G1 and S [133].

### 3.2.3. PKC $\epsilon$ kinase

PKC $\epsilon$  is an isoform that belongs to the novel PKC subfamily, which lacks the calcium binding domain and is regulated by diacylglycerol [134]. In breast cancer cell lines the downregulation of PKC $\epsilon$  caused inhibition of NF $\kappa$ B signaling, augmented apoptosis, and increased sensitivity toward ionizing radiation and treatment with chemotherapeutics [134]. In breast cancer it was also found that enhanced expression of miR-31 provokes an inhibition of the oncogenic NF $\kappa$ B pathway, and miR-31 directly targets PRKCE, the gene that encodes PKC $\epsilon$  [135].

### 3.2.4. PI3K-AKT-mTOR pathway

AKT is a serine/threonine kinase downstream of phosphoinositide 3-kinase (PI3K) pathway. Activated AKT stimulates numerous processes, including cell cycle progression, survival, metabolism and migration through phosphorylation of many substrates, including mammalian target of rapamycin (mTOR) [136]. Aberrant AKT signaling has been frequently documented in human cancers and appears to play a pivotal role in their progression [137]. Previous reports have identified the miR-302-367 cluster as a regulator of AKT1 expression [138], while miR-612 was related to AKT2 repression [139]. Through direct targeting AKT1 and cyclin D1, the ectopic expression of miR-302-367 cluster in cervical carcinoma cells (HeLa and SiHa cells) inhibited cell proliferation and tumor formation [138]. Tao et al. [139] have shown that miR-612 inhibited hepatocellular carcinoma cells proliferation, migration, invasion, and metastasis through direct targeting AKT2.

mTOR controls different biological processes including protein and lipid biosynthesis, energy metabolism, autophagy, cell survival, and cytoskeletal organization [140,141]. mTOR signaling is frequently deregulated in human cancers [142]. miR-99a [141,143] and miR-144 [144] have been shown to effectively downregulated mTOR expression by direct binding the 3'UTR of its mRNA. miR-99a

was found to be downregulated in renal cell carcinoma (RCC) and its restoration suppressed RCC cell growth, migration and invasion, as well as induced cell cycle arrest *in vitro*. Besides, intratumoral delivery of miR-99a inhibited tumor growth in murine xenograft model of human RCC [143].

### 3.2.5. Cytoskeleton regulators

In this Section 4 kinases that display an important role in cytoskeleton dynamics will be discussed.

Fyn is a non-receptor tyrosine kinase belonging to the Src family kinase and has a critical role in cell adhesion, proliferation, migration, and survival. Its overexpression is associated with several types of cancer, such as glioblastoma multiforme, squamous cell carcinoma of the head and neck, and melanoma [145]. Recently, Ninio-Many et al. [146] have shown that Fyn expression is dramatically reduced by elevated levels of miR-125a-3p. Besides, the activity of proteins downstream of Fyn, such as FAK, paxillin, and AKT, was also dropped and the cellular response was cell cycle arrest at the G2/M phases and migration blockage.

Focal adhesion kinase (FAK) is the prototype for a family of non-receptor tyrosine kinases. While FAK is ubiquitously expressed, a second member of this family, referred as PYK2, has a more restricted expression [147]. FAK is an important mediator of integrin signaling and is related to cellular processes such as cell motility, cell proliferation, and cell survival. Increased FAK expression is frequently observed in human tumors and has been associated with an increased malignant phenotype and poor prognosis [148,149]. FAK and PYK2 have been described as targets for some miRNAs. MiR-7 was shown to target FAK in glioblastoma cells [150] and breast cancer cells [149]. The overexpression of miR-7 in U87 and U251 glioblastoma cells inhibited their invasiveness and migration activity [150]. miR-7 expression was shown to be decreased in malignant versus normal breast tissue. The overexpression of miR-7 in breast cancer cells suppressed proliferation, anchorage independent growth, three-dimensional growth in matrigel, migration and invasion. Besides, miR-7 overexpression inhibited primary breast tumor development, local invasion, and metastasis of breast cancer xenografts [149]. PYK2 was found to be a target of miR-23b [151], miR-517a, and miR-517c [152]. Overexpression of miR-23b reduced PYK2 protein expression and significantly reduced glioma cell migration and invasion [151]. The expression of miR-517a and miR-517c was reduced in hepatocellular carcinoma (HCC) samples. Through PYK2 regulation, ectopic expression of miR-517a and miR-517c inhibited HCC cells proliferation by blocking the G2/M transition [152].

P21 activated kinases (PAKs) are serine/threonine kinases that function as important regulators of cytoskeletal dynamics and cell motility, but have also been implicated in controlling death and survival signaling and promoting cell proliferation. Consequently, PAKs have been related to cell transformation that results in tumor formation and cell invasiveness [153,154]. PAK1 expression is widely upregulated in breast tumors and correlates with tumor invasiveness and cyclin D1 expression [155]. PAK4 was found overexpressed in cell lines derived from different human cancers types, and this finding was also related to increased tumorigenesis [156]. In breast cancer cells, 3'UTR of PAK1 mRNA was targeted by miR-7, and the introduction of miR-7 in these cells inhibited motility, invasiveness, anchorage-independent growth and tumorigenic potential [157]. PAK2 was shown to be a direct target of miR-23b in breast cancer [158], whereas miR-145 was found to be a regulator of PAK4 expression in human colon cancer cells [159]. Wang et al. [159] showed that miR-145 is downregulated in human colon cancer cells, and restoration of miR-145 in SW620 cells attenuated cell growth *in vitro*. Additionally, the findings

demonstrated that miR-145 downregulates p-ERK expression by targeting PAK4, which leads to inhibition of tumor growth.

The Rho GTPase family has a key role in regulating actin cytoskeleton organization and dynamics. RhoA and RhoC family members act primarily through two serine/threonine kinases (ROCK1 and ROCK2) that mediate the phosphorylation of numerous downstream target proteins. ROCK kinases influence several processes such as cell morphology regulation, cell adhesion, cell motility, proliferation, differentiation, and apoptosis. Rho/ROCK pathway has been frequently implicated in oncogenic transformation [160]. Several miRNAs have been described as regulators of ROCK kinases expression. Among them, miR-146a [161], miR-148b [162], miR-335 [163], miR-584 [164] and miR-1280 [165] was shown to target ROCK1, whereas miR-124 [166], miR-138 [167] and miR-139 [168] targeted ROCK2. In a general, there is a correlation between knockdown of ROCK kinases by the cited miRNAs and reduction of cell migration and invasion.

### 3.3. Kinase receptors

#### 3.3.1. Insulin-like growth factor 1 receptor (IGF1R)

IGF1R is a tyrosine kinase receptor that is mainly activated by IGF1 and IGF2. IGF1R activates multiple downstream signaling cascades (PI3K/AKT and MAPK/ERK) involved in cell proliferation, differentiation, and survival [169,170]. Several miRNAs have been pointed out as modulators of IGF1R expression, such as miR-7 [171], miR-16 [172], miR-122 [173], miR-181b [174], miR-376a, miR-376c [175] and miR-497 [113,176].

Shi et al. [174] have reported that miR-181b inhibited glioma proliferation, migration and invasion by targeting IGF1R and its downstream signaling pathways, PI3K/AKT and MAPK/ERK1/2. IGF1R was overexpressed in glioma patients, and its protein levels were inversely correlated with miR-181b expression. Chen et al. [172] have reported that miR-16 is downregulated in osteosarcoma cell lines and tissues. The overexpression of miR-16 suppresses osteosarcoma cell proliferation and tumor growth in nude mice by inhibiting Raf1-MEK1/2-ERK1/2 pathway [172]. Additionally, ectopic expression of miR-7 in tongue squamous cell carcinoma (TSCC) cells reduced IGF1R expression and AKT activation, which in turn resulted in decreased cell proliferation and enhanced apoptosis [171]. miR-122, downregulated in breast cancer, also functions as a tumor suppressor *in vivo*, through targeting IGF1R and regulating PI3K/AKT/mTOR/p70S6K pathway [173]. In addition, miR-376a and miR-376c overexpression led to a significant decrease in melanoma cells migration by targeting IGF1R [175]. Luo et al. [176] supported a similar idea, showing that ectopic expression of miR-497 in cervical carcinoma cells suppressed migration and invasiveness of these cells by post-transcriptionally targeting IGF1R.

#### 3.3.2. Epidermal growth factor receptor (EGFR)

EGFR is a member of the ErbB receptor tyrosine kinase family. Several signaling pathways are downstream targets of EGFR, such as PI3K/AKT [177] and Ras/Raf/ERK1/2 pathways [178]. For this reason, EGFR signaling has emerged as a major therapeutic target in many cancers, since its deregulation have been associated to tumor cell proliferation, apoptosis resistance, invasion, angiogenesis and metastasis [179]. Some miRNAs have been shown to directly target EGFR and, hence, have been considered potential therapeutic tools for cancer treatment. Among them are miR-7 [179–181], miR-27a [182,183] and miR-133b [184].

#### 3.3.3. AXL

Tyro3-Axl-Mer (TAM) receptor tyrosine kinase subfamily (RTKs) comprises Tyro-3 (also called Sky), MER, and AXL. Signaling pathways activated by AXL include PI3K and ERK. The overexpression of

AXL has been reported in several cancer types, such as colon, esophageal, thyroid, breast, lung, liver and astrocytoma-glioblastoma carcinomas [185]. Increased AXL expression has been associated with improved cell survival, proliferation, migration, invasion, angiogenesis and metastasis [186].

AXL was recently described as a direct target of miR-34a [186,187] and miR-199a/b [186]. The major effect promoted by ectopic overexpression of these miRNAs was the decreased cell migration potential.

#### 3.3.4. c-MET

c-MET belongs to a subfamily of RTKs, which also includes RON and SEA [188]. Hepatocyte growth factor (HGF) is the unique ligand for c-MET. After activation by HGF binding, c-MET autophosphorylates, recruits adaptor proteins and activates multiple downstream effectors. c-MET potentially activates some signaling pathways as RAS-MAPK (mitogen-activated protein kinase), PI3K, AKT, and STAT3/5 (signal transducer and activator of transcription 3/5). As a consequence, activated c-MET regulates a variety of cellular processes, including proliferation, survival, adhesion, motility and invasion [189].

The c-MET/HGF pathway is one of the most commonly deregulated pathways in human cancers. In fact, the overexpression of c-MET and elevated levels of circulating HGF have been frequently detected in many cancers, and are associated with poor clinical outcomes [189]. Different studies have shown that the downregulation or inhibition of HGF/c-MET results in reduced cell growth, survival, migration and invasion *in vitro*, and decreased tumorigenic and metastatic potential *in vivo* [189,190]. Recently, a variety of miRNAs were shown to target c-MET in cancer cells, contributing to reduce their tumorigenic potential. Among them are miR-27a [182], miR-34a [191], miR-34c [192], miR-101 [193], miR-137 [194], miR-409-3p [195], miR-410 [196], miR-449a [197] and miR-7515 [198].

## 4. Modulation of protein phosphatases by miRNAs

Protein phosphatases are not as widely encoded in the human genome as their counterpart protein kinases. However, despite having relatively few genes encoding catalytic subunits of protein phosphatases, some of them have different regulatory subunits that, combined with the catalytic subunit, allow translocation and binding to specific targets [220].

MiRNAs can also regulate protein phosphatase expression, either by targeting the regulatory or the catalytic subunit, leading to hallmarks of tumor cells (unrestricted cell proliferation, resistance to apoptosis, and an improved ability of migration and invasion). We will describe below seven protein phosphatases that have established connections to cancer. Additional information can be found in Table 2, while some of the discussed phosphatases are also present in Figs. 3 and 4, which summarize signaling pathways and cellular responses these proteins take part in.

#### 4.1. Cell division cycle 25 (CDC25)

CDC25 is a group of highly conserved dual-specificity phosphatases. In mammals there are three different CDC25 homologs: CDC25A, B and C. All three stimulate cell cycle progression, since they dephosphorylate the inhibitory site of CDK, and, in turn, activate cyclin/CDK complexes [221,222]. Therefore, abnormal expression of those phosphatases has been observed in a number of tumors [223–229].

CDC25A is the major regulator of G1/S transition and S phase progression, while CDC25B contributes to G2 progression and CDC25C is restricted to mitosis [230]. CDC25A is overexpressed in a variety of human cancers [231], which is related to aberrant



**Table 2**  
Regulation of protein phosphatases by miRNAs in cancer.

Protein	Examples of substrates	Regulatory miRNAs	Protein data	Altered miRNA expression
CDC25	CDK1 [279], CDK2 [279], p53 [280]	miR-21 [233], miR-125b [61], miR-449a [68,232] and miR-449b [68]	Highly expressed in cancer cells [231]	Overexpression leads to reduced cell proliferation and growth [61,232–235]
PTEN	Phosphatidylinositol 3,4,5-trisphosphate. Indirectly inhibits AKT [202]	miR-21, miR-23a, miR-26, miR-144 [237], miR-148a, miR-153 [238], miR-214 [239], miR-216a and miR-217 [240], miR-221 [241], miR-222 [242], miR-301 [243] as well as miR-374a [244]	Tumor suppressor. The gene is mutated in several cancers [236]	Overexpression is associated to cancer cell survival, migration, growth and proliferation [245–256]
PTPN9	N/A	miR-24 [258]	Reduced expression in cancer cells [258]	N/A
PRL1	N/A	miR-339-5p [259]	Overexpressed in cancer cells. Stimulates progression in the cell cycle [281,282]	Overexpression leads to inhibition of cell growth and reduced migration and invasion in colorectal cancer cells. It also suppressed tumor growth <i>in vivo</i> [259]
PP6	TAK1 [283]	miR-373 [260]	Negative cell cycle regulator [284]	Overexpression promotes cell proliferation of hepatocellular carcinoma cell lines [260]
PTPN1	Insulin receptor [285], BCR [286], Src [287], PDGFR [288]	miR-362-3p [263]	Both tumor suppressor and tumor promoter [261]. Leads to proliferation and metastasis in colon cancer and breast cancer cells [262,263]	Overexpression is associated with reduced risk of recurrence in adenocarcinoma patients [263]
PPM1D	ATM [267], CHK1 [265], p53 [265], CHK2 [266]	miR-16 [275]	Regulator of DNA damage signaling pathways, frequently overexpressed in many human cancers [264]	Overexpression suppressed the growth of mouse mammary tumor stem cells and sensitized MCF-7 human breast cancer cells to doxorubicin [275]

progression through G1/S and increased proliferation. miR-125b [61], miR-449a [68,232] and miR-449b [68] are all regulators of CDC25A expression. The expression of all these miRNAs was found to be reduced in cancer cells, which was related to CDC25A overexpression. miR-21, despite targeting many tumor suppressor proteins, also targets CDC25A [233].

The overexpression of miR-125b in U251 glioma stem cells decreased CDC25A expression and cell proliferation through cell cycle arrest at G1 phase [61]. Besides, the restoration of miR-449a in human bladder cancer cells inhibited cell growth and induced G1 phase arrest through downregulation of CDC25A and CDK6 [232]. Together, these results demonstrate the importance of controlling CDC25A expression to achieve tumor suppression.

Liffers et al. [234] have shown for the first time a down-regulation of CDC25B by miR-148a. They showed that miR-148a exhibited a significant 4-fold downregulation in human pancreatic ductal adenocarcinoma (PDAC) in comparison to normal pancreatic ductal cells. The same authors also reported that the overexpression of miR-148a in the pancreatic cancer cell line IMIM-PC2 inhibited tumor cell growth and colony formation. CDC25B was identified as a potential target of miR-148a by *in silico* analysis using PicTar, Targetscan and miRanda together with Gene Ontology. Accordingly, the activity of a luciferase reporter containing the 3'UTR of CDC25B was repressed in the presence of miR-148a mimics, confirming that miR-148a targets the 3'UTR of CDC25B. Yu et al. [235] have observed that miR-141 is significantly down-regulated in renal cell carcinoma and, by using miRNA target prediction software, they found that miR-141 could target the 3'UTR sequence of CDC25B. When miR-141 expression was restored in renal cell carcinoma the proliferation rate of those cells was suppressed and the endogenous CDC25B protein levels dropped. In addition, the authors suggested that the transcriptional loss of miR-141 and the resultant increase in CDC25B expression facilitates increased genomic instability at an early stage of renal cell carcinoma development.

#### 4.2. Phosphatase and tensin homolog (PTEN)

PTEN is a dual-specificity phosphatase and well-known tumor suppressor gene. As a lipid phosphatase, PTEN dephosphorylates

phosphatidylinositol 3,4,5-trisphosphate (PIP3) to phosphatidylinositol 3,4,5-diphosphate (PIP2), thereby inhibiting AKT, a key kinase that mediates proliferation and cell survival [236]. Several miRNAs, including miR-21, miR-23a, miR-26, miR-144 [237], miR-148a, miR-153 [238], miR-214 [239], miR-216a and miR-217 [240], miR-221 [241], miR-222 [242], miR-301 [243], and miR-374a [244] have been established as regulators of PTEN expression.

miR-21 was one of the first miRNAs detected in the human genome, and is also known to be up-regulated in many types of human malignancies [245]. It has been shown that several tumor suppressors including PTEN [246], tumor suppressor gene tropomyosin 1 (TPM1) [247], programmed cell death 4 (PDCD4) [248], maspin [249], and matrix metalloproteinases inhibitors RECK and TIMP3 [250] are targets of miR-21, suggesting that miR-21 is an important oncogenic miRNA closely related to tumor growth and metastasis.

Gliomas are the most common and deadly type of primary brain tumor. cAMP response element-binding protein (CREB), a proto-oncogenic transcription factor that is overexpressed in gliomas and is able to promote gliomagenesis by modulating the expression of miR-23a, which targets the tumor suppressor PTEN [251]. miR-26 can also enhance gliomagenesis by affecting PTEN levels [252].

miR-148a stimulated cell survival, cell migration, anchorage independent cell growth and tumorigenesis in SCID by targeting PTEN [253].

PTEN is downregulated by miR-214 in ovarian cancer [254]. In addition, miR-216a and miR-217, both of which target PTEN, activate AKT through PTEN down-regulation in kidney disorders [255].

miR-221 and miR-222 had an increase of expression in HCT116 and HKe3 cells (HCT116 cells in which mutated *KRAS* allele was deleted) in 3D culture. These miRNAs were regulated by oncogenic *KRAS* and were significantly overexpressed in human colorectal tumor specimens. Importantly, both miRNAs target PTEN [256].

miR-301 overexpression has been implicated as a negative prognostic indicator in lymph node negative invasive ductal breast cancer. By using luciferase reporter assay, PTEN was identified as a target of miR-301 [243].

miR-374a is markedly upregulated in primary tumor samples from patients with distant metastases and was associated with poor metastasis-free survival. This miR directly targets and



suppresses multiple negative regulators of the Wnt/ $\beta$ -catenin signaling cascade, including PTEN [244].

#### 4.3. Tyrosine-protein phosphatase non-receptor type 9 (PTPN9)

Tyrosine-protein phosphatase non-receptor type 9 (also called PTP-MEG2) is encoded by the *PTPN9* gene, in human. PTPN9 inhibits receptor PTK activation by direct dephosphorylation of EGFR and ErbB2 [257].

Du et al. [258] have observed that miR-24-expressing cells and tumors displayed higher levels of phosphorylated EGFR, whereas expression of the phosphatases PTPN9 and receptor-type tyrosine-protein phosphatase F (PTPRF) were repressed. Those authors confirmed that miR-24 directly targets both PTPN9 and PTPRF. Besides, they observed that the amount of phosphorylated EGFR was much higher in patients with metastatic breast carcinoma whereas levels of PTPN9 and PTPRF were lower.

#### 4.4. Phosphatase of regenerating liver (PRL)

PRL is a group of protein tyrosine phosphatases consisting of three members, PRL-1, -2, and -3. There is an inverse correlation between miR-339-5p and PRL-1 expressions. Zhou et al. [259] found downregulated miR-339-5p levels in colorectal cancer tissues and highly invasive CRC cell lines. Furthermore, overexpression of miR-339-5p in CRC leads to inhibition of cell growth, migration and invasion. It also suppressed tumor growth *in vivo*. Overexpression of miR-339-5p also reduces the expression of PRL-1 mRNA and protein, which was associated with low expression of phosphorylated-extracellular signal-regulated kinase 1/2 (p-ERK1/2). miR-339-5p acts as a tumor suppressor and plays a role in metastasis and inhibition of cellular growth as demonstrated in CRC cells via targeting PRL-1 and regulating p-ERK1/2.

#### 4.5. Protein phosphatase 6 (PP6)

PP6 is a serine/threonine protein phosphatase classified as a type 2A phosphatase family member based on its sequence homology to the catalytic subunit of PP2A. Wu et al. [260] found that miR-373 is upregulated in human hepatocellular carcinoma (HCC) tissues as compared with adjacent normal tissues, and promotes the proliferation of the HCC cell lines HepG2 and QGY-7703 by regulating the cell cycle progression from G1 to S-phase. These authors detected the gene encoding the protein phosphatase 6 catalytic subunit (PPP6C), a negative cell cycle regulator, as a direct target gene of miR-373. Accordingly, overexpression of PPP6C abolished the regulation of cell cycle and cell growth exercised by miR-373 in HepG2 cells.

#### 4.6. Protein-tyrosine phosphatase non-receptor type 1 (PTPN1 or PTP1B)

Protein-tyrosine phosphatase non-receptor type 1 (PTPN1 or PTP1B) has, over the last decade, attracted great attention due to its role in leptin and insulin metabolism and, more recently, in cancer [261]. Increased levels of PTPN1 have been shown to promote tumorigenicity of colon cancer cells by the activation of Src [262]. In breast cancer PTPN1 activity is required for efficient invadopodia formation and breast cancer invasion. Christensen et al. [263] have reported that high miR-362-3p expression in adenocarcinomas was found to be associated with reduced risk of recurrence in two independent patient cohorts. Functional characterization of miR-362-3p demonstrated that this miRNA inhibits proliferation. miR-362-3p transfection into colon cancer cell lines decreased among

others the amount of PTPN1, which was identified as direct miR-362-3p targets.

#### 4.7. Wild-type p53-induced phosphatase 1 (Wip1 or PPM1D)

Wip1 is a serine/threonine phosphatase that has a critical role as regulator of DNA damage signaling pathways [264]. Wip1 dephosphorylates several DNA damage-responsive proteins, such as ATM, CHK1, and CHK2 [265–267], reverting DNA damage-induced cell cycle checkpoint. Thus, Wip1 releases cells from the cell cycle arrest induced by DNA damage, allowing them to progress through the cell cycle. PPM1D is an oncogene frequently amplified in human tumors, which results in Wip1 overexpression in many different cancer types, including breast carcinomas, ovarian adenocarcinomas, neuroblastomas, pancreatic adenocarcinomas, gastric carcinomas, and medulloblastomas [268–274]. miR-16 was previously identified as a repressor of Wip1 expression [275]. After DNA damage, miR-16 is immediately induced, which in turn delays the increase in Wip1 expression. This effect prevents a premature inactivation of ATM/ATR signaling and allows DNA damage repair. miR-16 was found downregulated in mammary tumor stem cells. Its overexpression suppressed the growth of mouse mammary tumor stem cells and sensitized MCF-7 human breast cancer cells to treatment with doxorubicin [275].

#### 4.8. Src homology-2 domain-containing inositol 5-phosphatase 1 (SHIP1)

SHIP1 hydrolyzes the 5' phosphate from phosphatidylinositol (3,4,5)-triphosphate and inositol-1,3,4,5-tetrakisphosphate, affecting multiple signaling pathways. SHIP1 expression is restricted to hematopoietic cells where it negatively regulates myeloid cell proliferation and survival. O'Connell and collaborators [276] identified SHIP1 as a direct target of miR-155. Ectopic expression of miR-155 resulted in a myeloproliferative disorder. Besides, it was reported that in diffuse large B cell lymphoma high levels of miR-155 are associated with diminished SHIP1 expression [277]. Furthermore, transfection of a myeloid cell line with miR-210 resulted in loss of SHIP-1 protein expression [278].

### 5. Final considerations

We have drawn a picture of the important role miRNAs exert on regulating protein kinase and protein phosphatase expression and the effects this regulation has, or may have, in the control of cell growth, survival, proliferation, adhesion, and every other cancer-related signaling pathway. The antagonistic nature of protein kinases and protein phosphatases ensures that the knowledge of their expression and their activities is critical for the understanding of oncogenic processes. Besides, we do believe that the identification of miRNAs that target protein phosphatases and *vice versa* will open a new avenue to understanding the role of this enzyme family in cancer biology, since during the last few decades there a huge amount of information about kinases has been released but the same cannot be said for phosphatases.

Not all miRNAs regulating phosphatases and kinases, however, act through straightforward processes. The complexity of the phosphatase (but also other enzymes, such as kinases) specificity mechanism, with few catalytic subunits combined to an array of regulatory subunits, warrants a tangled regulatory network that may seem ambiguous at first look. Therefore, studies that address how microRNAs interfere in feedback loops and protein interactions in pathways formed by protein kinases and phosphatases are the next step towards understanding regulatory mechanisms that trigger the induction and development of

malignant transformations. In addition, it is important to keep in mind the complexity of miRNA action, mainly regarding multiple targets for a given miRNA, which makes the action of miRNAs dependent on cell type, metabolic context and target mRNAs that are expressed. Besides, miRNAs do not act in isolation, which is why it is essential to have an overview of the context at the miRNome, transcriptome and proteome levels in order to assess the metabolic network as a whole rather than a unique target mRNA or pathway. In addition, just to give an idea about how complex this process can be, it has become clear that some miRNAs target both oncogenic and tumor suppressor proteins. This is the case of miR-21, miR-29 and the miR-17-92 cluster [289–291].

In relation to the application of miRNAs, there are some recent reviews that comprehensively covered this subject. In general, it has been reported that miRNAs have great potential in the diagnosis, prognosis and cancer therapy [292–294]. For instance, miR-21 has been considered as a biomarker for diagnostic and prognostic assessments, as elevated blood levels of miR-21 were observed to be correlated with resistance of pancreatic and lung cancer [295,296]. More recently, Manceau and collaborators [297] reported that the expression of miR-31-3p can be used to identify patients with wild-type *KRAS* metastatic colorectal cancer who are good responders to anti-EGFR therapy. Concerning treatment, miR-34 is one of the first candidates for targeted cancer therapy. Systemic delivery of miR-34 caused a decrease in a murine model of lung cancer [298]. Despite the great potential of miRNAs as therapeutic agents, there are some challenges in using them clinically: a) development of efficient systems for delivering miR or anti-miR *in vivo* and consequently improving the biological effect and minimizing off-targets; b) understanding the miRNA network *in vivo* and be able to predict toxicity.

## Conflict of interest

The authors declare no conflict of interest.

## References

- [1] T. Hunter, J.A. Cooper, Protein-tyrosine kinases, *Annu. Rev. Biochem.* 54 (1985) 897–930, <http://dx.doi.org/10.1146/annurev.bi.54.070185.004341>.
- [2] J.D. Graves, E.G. Krebs, Protein phosphorylation and signal transduction, *Pharmacol. Ther.* 82 (1999) 111–121.
- [3] G. Manning, D.B. Whyte, R. Martinez, T. Hunter, S. Sudarsanam, The protein kinase complement of the human genome, *Science* 298 (2002) 1912–1934, <http://dx.doi.org/10.1126/science.1075762>.
- [4] G. Manning, G.D. Plowman, T. Hunter, S. Sudarsanam, Evolution of protein kinase signaling from yeast to man, *Trends Biochem. Sci.* 27 (2002) 514–520.
- [5] C.E. Gee, I.M. Mansuy, Protein phosphatases and their potential implications in neuroprotective processes, *Cell. Mol. Life Sci. CMLS* 62 (2005) 1120–1130, <http://dx.doi.org/10.1007/s00018-005-5008-4>.
- [6] N.K. Tonks, Protein tyrosine phosphatases: from genes, to function, to disease, *Nat. Rev. Mol. Cell Biol.* 7 (2006) 833–846, <http://dx.doi.org/10.1038/nrm2039>.
- [7] D. Easty, W. Gallagher, D.C. Bennett, Protein tyrosine phosphatases, new targets for cancer therapy, *Curr. Cancer Drug Targets* 6 (2006) 519–532.
- [8] S. Caenepeel, G. Charyczak, S. Sudarsanam, T. Hunter, G. Manning, The mouse kinase: discovery and comparative genomics of all mouse protein kinases, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 11707–11712, <http://dx.doi.org/10.1073/pnas.0306880101>.
- [9] H. Zhu, J.F. Klemic, S. Chang, P. Bertone, A. Casamayor, K.G. Klemic, et al., Analysis of yeast protein kinases using protein chips, *Nat. Genet.* 26 (2000) 283–289, <http://dx.doi.org/10.1038/81576>.
- [10] A. Alonso, J. Sasin, N. Bottini, I. Friedberg, I. Friedberg, A. Osterman, et al., Protein tyrosine phosphatases in the human genome, *Cell* 117 (2004) 699–711, <http://dx.doi.org/10.1016/j.cell.2004.05.018>.
- [11] Y. Shi, Serine/threonine phosphatases: mechanism through structure, *Cell* 139 (2009) 468–484, <http://dx.doi.org/10.1016/j.cell.2009.10.006>.
- [12] M. Esteller, Non-coding RNAs in human disease, *Nat. Rev. Genet.* 12 (2011) 861–874, <http://dx.doi.org/10.1038/nrg3074>.
- [13] U.A. Ørom, F.C. Nielsen, A.H. Lund, MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation, *Mol. Cell* 30 (2008) 460–471, <http://dx.doi.org/10.1016/j.molcel.2008.05.001>.
- [14] D.P. Bartel, MicroRNAs: genomics, biogenesis, mechanism, and function, *Cell* 116 (2004) 281–297.
- [15] J. Brennecke, A. Stark, R.B. Russell, S.M. Cohen, Principles of microRNA-target recognition, *PLoS Biol.* 3 (2005) e85, <http://dx.doi.org/10.1371/journal.pbio.0030085>.
- [16] M.S. Ebert, P.A. Sharp, Roles for microRNAs in conferring robustness to biological processes, *Cell* 149 (2012) 515–524, <http://dx.doi.org/10.1016/j.cell.2012.04.005>.
- [17] J.T. Mendell, E.N. Olson, MicroRNAs in stress signaling and human disease, *Cell* 148 (2012) 1172–1187, <http://dx.doi.org/10.1016/j.cell.2012.02.005>.
- [18] D.P. Bartel, MicroRNAs: target recognition and regulatory functions, *Cell* 136 (2009) 215–233, <http://dx.doi.org/10.1016/j.cell.2009.01.002>.
- [19] H. Yamamoto, Y. Adachi, H. Taniguchi, H. Kunimoto, K. Noshio, H. Suzuki, et al., Interrelationship between microsatellite instability and microRNA in gastrointestinal cancer, *World J. Gastroenterol.* WJG 18 (2012) 2745–2755, <http://dx.doi.org/10.3748/wjg.v18.i22.2745>.
- [20] A.S. Ali, A. Ahmad, S. Ali, B. Bao, P.A. Philip, F.H. Sarkar, The role of cancer stem cells and miRNAs in defining the complexities of brain metastasis, *J. Cell. Physiol.* 228 (2013) 36–42, <http://dx.doi.org/10.1002/jcp.24127>.
- [21] K.C. Vickers, A.T. Remaley, Lipid-based carriers of microRNAs and intercellular communication, *Curr. Opin. Lipidol.* 23 (2012) 91–97, <http://dx.doi.org/10.1097/MOL.0b013e328350a425>.
- [22] F. Berger, M.F. Reiser, Micro-RNAs as potential new molecular biomarkers in oncology: have they reached relevance for the clinical imaging sciences? *Theranostics* 3 (2013) 943–952, <http://dx.doi.org/10.7150/thno.7445>.
- [23] J. Jarry, D. Schadendorf, C. Greenwood, A. Spatz, L.C. van Kempen, The validity of circulating microRNAs in oncology: five years of challenges and contradictions, *Mol. Oncol.* 8 (2014) 819–829, <http://dx.doi.org/10.1016/j.molonc.2014.02.009>.
- [24] X. Tang, M. Li, L. Tucker, B. Ramratnam, Glycogen synthase kinase 3 beta (GSK3β) phosphorylates the RNAase III enzyme Drosha at S300 and S302, *PLoS One* 6 (2011) e20391, <http://dx.doi.org/10.1371/journal.pone.0020391>.
- [25] Y. Feng, X. Zhang, Q. Song, T. Li, Y. Zeng, Drosha processing controls the specificity and efficiency of global microRNA expression, *Biochim. Biophys. Acta* 1809 (2011) 700–707, <http://dx.doi.org/10.1016/j.bbarm.2011.05.015>.
- [26] A.D. Haase, L. Jaskiewicz, H. Zhang, S. Lainé, R. Sack, A. Gagnon, et al., TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA silencing, *EMBO Rep.* 6 (2005) 961–967, <http://dx.doi.org/10.1038/sj.embo.7400509>.
- [27] Y. Lee, I. Hur, S.-Y. Park, Y.-K. Kim, M.R. Suh, V.N. Kim, The role of PACT in the RNA silencing pathway, *EMBO J.* 25 (2006) 522–532, <http://dx.doi.org/10.1038/sj.emboj.7600942>.
- [28] T.P. Chendrimada, R.L. Gregory, E. Kumaraswamy, J. Norman, N. Cooch, K. Nishikura, et al., TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing, *Nature* 436 (2005) 740–744, <http://dx.doi.org/10.1038/nature03868>.
- [29] Z. Paroo, X. Ye, S. Chen, Q. Liu, Phosphorylation of the human microRNA-generating complex mediates MAPK/Erk signaling, *Cell* 139 (2009) 112–122, <http://dx.doi.org/10.1016/j.cell.2009.06.044>.
- [30] J.K. Mito, H.D. Min, Y. Ma, J.E. Carter, B.E. Brigrman, L. Dodd, et al., Oncogene-dependent control of miRNA biogenesis and metastatic progression in a model of undifferentiated pleomorphic sarcoma, *J. Pathol.* 229 (2013) 132–140, <http://dx.doi.org/10.1002/path.4099>.
- [31] P.A. Ferreira, R.R. Ruela-de-Sousa, K.C.S. Queiroz, A.C.S. Souza, R. Milani, R.A. Pilli, et al., Knocking down low molecular weight protein tyrosine phosphatase (LMW-PTP) reverts chemoresistance through inactivation of Src and Bcr-Abl proteins, *PLoS ONE* 7 (2012) e44312, <http://dx.doi.org/10.1371/journal.pone.0044312>.
- [32] E. van Rooij, L.B. Sutherland, X. Qi, J.A. Richardson, J. Hill, E.N. Olson, Control of stress-dependent cardiac growth and gene expression by a microRNA, *Science* 316 (2007) 575–579, <http://dx.doi.org/10.1126/science.1139089>.
- [33] A. Baccarini, H. Chauhan, T.J. Gardner, A.D. Jayaprakash, R. Sachidanandam, B.D. Brown, Kinetic analysis reveals the fate of a microRNA following target regulation in mammalian cells, *Curr. Biol. CB* 21 (2011) 369–376, <http://dx.doi.org/10.1016/j.cub.2011.01.067>.
- [34] M.P. Gantier, C.E. McCoy, I. Rusinova, D. Saulep, D. Wang, D. Xu, et al., Analysis of microRNA turnover in mammalian cells following Dicer1 ablation, *Nucleic Acids Res.* 39 (2011) 5692–5703, <http://dx.doi.org/10.1093/nar/gkr148>.
- [35] E.J. Lee, M. Baek, Y. Gusev, D.J. Brackett, G.J. Nuovo, T.D. Schmittgen, Systematic evaluation of microRNA processing patterns in tissues, cell lines, and tumors, *RNA N. Y.* 14 (2008) 35–42, <http://dx.doi.org/10.1261/rna.804508>.
- [36] H.-W. Hwang, E.A. Wentzel, J.T. Mendell, A hexanucleotide element directs microRNA nuclear import, *Science* 315 (2007) 97–100, <http://dx.doi.org/10.1126/science.1136235>.
- [37] L. Marcinowski, M. Tanguy, A. Krmpotic, B. Rädle, V.J. Lisnić, L. Tuddenham, et al., Degradation of cellular mir-27 by a novel, highly abundant viral transcript is important for efficient virus replication *in vivo*, *PLoS Pathog.* 8 (2012) e1002510, <http://dx.doi.org/10.1371/journal.ppat.1002510>.
- [38] R. Avraham, A. Sas-Chen, O. Manor, I. Steinfeld, R. Shalgi, G. Tarcic, et al., EGF decreases the abundance of microRNAs that restrain oncogenic transcription factors, *Sci. Signal.* 3 (2010) ra43, <http://dx.doi.org/10.1126/scisignal.2000876>.
- [39] W. Tirasophon, A.A. Welihinda, R.J. Kaufman, A stress response pathway from the endoplasmic reticulum to the nucleus requires a novel bifunctional protein kinase/endoribonuclease (Ire1p) in mammalian cells, *Genes Dev.* 12 (1998) 1812–1824.
- [40] X.Z. Wang, H.P. Harding, Y. Zhang, E.M. Jolicoeur, M. Kuroda, D. Ron, Cloning of mammalian Ire1 reveals diversity in the ER stress responses, *EMBO J.* 17 (1998) 5708–5717, <http://dx.doi.org/10.1093/emboj/17.19.5708>.

- [41] J.-P. Upton, L. Wang, D. Han, E.S. Wang, N.E. Huskey, L. Lim, et al., IRE1 $\alpha$  cleaves select microRNAs during ER stress to derepress translation of pro-apoptotic Caspase-2, *Science* 338 (2012) 818–822, <http://dx.doi.org/10.1126/science.1226191>.
- [42] Y.W. Iwasaki, K. Kiga, H. Kayo, Y. Fukuda-Yuzawa, J. Weise, T. Inada, et al., Global microRNA elevation by inducible exportin 5 regulates cell cycle entry, *RNA* N. Y. 19 (2013) 490–497, <http://dx.doi.org/10.1261/rna.036608.112>.
- [43] A. Bardelli, S. Saha, J.A. Sager, K.E. Romans, B. Xin, S.D. Markowitz, et al., PRL-3 expression in metastatic cancers, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 9 (2003) 5607–5615.
- [44] S. Saha, A. Bardelli, P. Buckhaults, V.E. Velculescu, C. Rago, B. St Croix, et al., A phosphatase associated with metastasis of colorectal cancer, *Science* 294 (2001) 1343–1346, <http://dx.doi.org/10.1126/science.1065817>.
- [45] J. Zhang, Z. Xiao, D. Lai, J. Sun, C. He, Z. Chu, et al., miR-21, miR-17 and miR-19a induced by phosphatase of regenerating liver-3 promote the proliferation and metastasis of colon cancer, *Br. J. Cancer* 107 (2012) 352–359, <http://dx.doi.org/10.1038/bjc.2012.251>.
- [46] M.J. Bueno, M. Malumbres, MicroRNAs and the cell cycle, *Biochim. Biophys. Acta* 1812 (2011) 592–601, <http://dx.doi.org/10.1016/j.bbdis.2011.02.002>.
- [47] A. Lujambio, S. Ropero, E. Ballestar, M.F. Fraga, C. Cerrato, F. Setién, et al., Genetic unmasking of an epigenetically silenced microRNA in human cancer cells, *Cancer Res.* 67 (2007) 1424–1429, <http://dx.doi.org/10.1158/0008-5472.CAN-06-4218>.
- [48] G. de Cárcer, I. Pérez de Castro, M. Malumbres, Targeting cell cycle kinases for cancer therapy, *Curr. Med. Chem.* 14 (2007) 969–985.
- [49] M. Malumbres, M. Barbacid, Mammalian cyclin-dependent kinases, *Trends Biochem. Sci.* 30 (2005) 630–641, <http://dx.doi.org/10.1016/j.tibs.2005.09.005>.
- [50] M. Malumbres, M. Barbacid, To cycle or not to cycle: a critical decision in cancer, *Nat. Rev. Cancer* 1 (2001) 222–231, <http://dx.doi.org/10.1038/35106065>.
- [51] K.A. Schafer, The cell cycle: a review, *Vet. Pathol.* 35 (1998) 461–478.
- [52] S. Lapenna, A. Giordano, Cell cycle kinases as therapeutic targets for cancer, *Nat. Rev. Drug Discov.* 8 (2009) 547–566, <http://dx.doi.org/10.1038/nrd2907>.
- [53] L. He, X. He, S.W. Lowe, G.J. Hannon, microRNAs join the p53 network – another piece in the tumour-suppression puzzle, *Nat. Rev. Cancer* 7 (2007) 819–822, <http://dx.doi.org/10.1038/nrc2232>.
- [54] Y. Lin, J. Wu, H. Chen, Y. Mao, Y. Liu, Q. Mao, et al., Cyclin-dependent kinase 4 is a novel target in microRNA-195-mediated cell cycle arrest in bladder cancer cells, *FEBS Lett.* 586 (2012) 442–447, <http://dx.doi.org/10.1016/j.febslet.2012.01.027>.
- [55] M. Hafner, M. Landthaler, L. Burger, M. Khorshid, J. Hausser, P. Berninger, et al., Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP, *Cell* 141 (2010) 129–141, <http://dx.doi.org/10.1016/j.cell.2010.03.009>.
- [56] J.-J. Zhao, J. Lin, T. Lwin, H. Yang, J. Guo, W. Kong, et al., microRNA expression profile and identification of miR-29 as a prognostic marker and pathogenetic factor by targeting CDK6 in mantle cell lymphoma, *Blood* 115 (2010) 2630–2639, <http://dx.doi.org/10.1182/blood-2009-09-243147>.
- [57] G.T. Bommer, I. Gerin, Y. Feng, A.J. Kaczorowski, R. Kuick, R.E. Love, et al., p53-mediated activation of miRNA34 candidate tumor-suppressor genes, *Curr. Biol. CB* 17 (2007) 1298–1307, <http://dx.doi.org/10.1016/j.cub.2007.06.068>.
- [58] L. Feng, Y. Xie, H. Zhang, Y. Wu, miR-107 targets cyclin-dependent kinase 6 expression, induces cell cycle G1 arrest and inhibits invasion in gastric cancer cells, *Med. Oncol. Northwood Lond. Engl.* 29 (2012) 856–863, <http://dx.doi.org/10.1007/s12032-011-9823-1>.
- [59] L. Böhlig, M. Friedrich, K. Engeland, p53 activates the PANK1/miRNA-107 gene leading to downregulation of CDK6 and p130 cell cycle proteins, *Nucleic Acids Res.* 39 (2011) 440–453, <http://dx.doi.org/10.1093/nar/gkq796>.
- [60] X. Agirre, A. Vilas-Zornoza, A. Jiménez-Velasco, J.I. Martín-Subero, L. Cordeu, L. Gárate, et al., Epigenetic silencing of the tumor suppressor microRNA Hsa-miR-124a regulates CDK6 expression and confers a poor prognosis in acute lymphoblastic leukemia, *Cancer Res.* 69 (2009) 4443–4453, <http://dx.doi.org/10.1158/0008-5472.CAN-08-4025>.
- [61] L. Shi, J. Zhang, T. Pan, J. Zhou, W. Gong, N. Liu, et al., MiR-125b is critical for the suppression of human U251 glioma stem cell proliferation, *Brain Res.* 1312 (2010) 120–126, <http://dx.doi.org/10.1016/j.brainres.2009.11.056>.
- [62] J. Wu, J. Qian, C. Li, L. Kwok, F. Cheng, P. Liu, et al., miR-129 regulates cell proliferation by downregulating Cdk6 expression, *Cell Cycle Georget. Tex.* 9 (2010) 1809–1818.
- [63] X. Zhu, Y. Li, H. Shen, H. Li, L. Long, L. Hui, et al., miR-137 inhibits the proliferation of lung cancer cells by targeting Cdc42 and Cdk6, *FEBS Lett.* 587 (2013) 73–81, <http://dx.doi.org/10.1016/j.febslet.2012.11.004>.
- [64] J. Cai, J. Wu, H. Zhang, L. Fang, Y. Huang, Y. Yang, et al., miR-186 down-regulation correlates with poor survival in lung adenocarcinoma, where it interferes with cell-cycle regulation, *Cancer Res.* 73 (2013) 756–766, <http://dx.doi.org/10.1158/0008-5472.CAN-12-2651>.
- [65] M. Colamaio, E. Borbone, L. Russo, M. Bianco, A. Federici, D. Califano, et al., miR-191 down-regulation plays a role in thyroid follicular tumors through CDK6 targeting, *J. Clin. Endocrinol. Metab.* 96 (2011) E1915–E1924, <http://dx.doi.org/10.1210/jc.2011-0408>.
- [66] H. Deng, Y. Guo, H. Song, B. Xiao, W. Sun, Z. Liu, et al., MicroRNA-195 and microRNA-378 mediate tumor growth suppression by epigenetical regulation in gastric cancer, *Gene* 518 (2013) 351–359, <http://dx.doi.org/10.1016/j.gene.2012.12.103>.
- [67] S. Venkataraman, D.K. Birks, I. Balakrishnan, I. Alimova, P.S. Harris, P.R. Patel, et al., MicroRNA 218 acts as a tumor suppressor by targeting multiple cancer phenotype-associated genes in medulloblastoma, *J. Biol. Chem.* 288 (2013) 1918–1928, <http://dx.doi.org/10.1074/jbc.M112.396762>.
- [68] X. Yang, M. Feng, X. Jiang, Z. Wu, Z. Li, M. Aau, et al., miR-449a and miR-449b are direct transcriptional targets of E2F1 and negatively regulate pRb-E2F1 activity through a feedback loop by targeting CDK6 and CDC25A, *Genes Dev.* 23 (2009) 2388–2393, <http://dx.doi.org/10.1101/gad.1819009>.
- [69] A.V. Olaru, G. Ghiaur, S. Yamanaka, D. Luvsanjav, F. An, I. Popescu, et al., MicroRNA down-regulated in human cholangiocarcinoma control cell cycle through multiple targets involved in the G1/S checkpoint, *Hepatol. Baltim. Md.* 54 (2011) 2089–2098, <http://dx.doi.org/10.1002/hep.24591>.
- [70] R.-Q. Tian, X.-H. Wang, L.-J. Hou, W.-H. Jia, Q. Yang, Y.-X. Li, et al., MicroRNA-372 is down-regulated and targets cyclin-dependent kinase 2 (CDK2) and cyclin A1 in human cervical cancer, which may contribute to tumorigenesis, *J. Biol. Chem.* 286 (2011) 25556–25563, <http://dx.doi.org/10.1074/jbc.M111.221564>.
- [71] E.A. Afanasyeva, P. Mestdagh, C. Kumps, J. Vandesompele, V. Ehemann, J. Theissen, et al., MicroRNA miR-885-5p targets CDK2 and MCM5, activates p53 and inhibits proliferation and survival, *Cell Death Differ.* 18 (2011) 974–984, <http://dx.doi.org/10.1038/cdd.2010.164>.
- [72] J.-Y. Lee, W. Jeong, J.-H. Kim, J. Kim, F.W. Bazer, J.Y. Han, et al., Distinct expression pattern and post-transcriptional regulation of cell cycle genes in the glandular epithelia of avian ovarian carcinomas, *PLoS One* 7 (2012) e51592, <http://dx.doi.org/10.1371/journal.pone.0051592>.
- [73] W.W. Chien, C. Domenech, R. Catallo, T. Kaddar, J.-P. Magaud, G. Salles, et al., Cyclin-dependent kinase 1 expression is inhibited by p16(INK4a) at the post-transcriptional level through the microRNA pathway, *Oncogene* 30 (2011) 1880–1891, <http://dx.doi.org/10.1038/onc.2010.570>.
- [74] M. Malumbres, M. Barbacid, Cell cycle kinases in cancer, *Curr. Opin. Genet. Dev.* 17 (2007) 60–65, <http://dx.doi.org/10.1016/j.gde.2006.12.008>.
- [75] K.R. Kelly, J. Ecsedy, D. Mahalingam, S.T. Nawrocki, S. Padmanabhan, F.J. Giles, et al., Targeting aurora kinases in cancer treatment, *Curr. Drug Targets* 12 (2011) 2067–2078.
- [76] A. Lal, F. Navarro, C.A. Maher, L.E. Maliszewski, N. Yan, E. O'Day, et al., miR-24 inhibits cell proliferation by targeting E2F2, MYC, and other cell-cycle genes via binding to "seedless" 3'UTR microRNA recognition elements, *Mol. Cell* 35 (2009) 610–625, <http://dx.doi.org/10.1016/j.molcel.2009.08.020>.
- [77] P. Liu, M. Qi, C. Ma, G. Lao, Y. Liu, Y. Liu, et al., Let7a inhibits the growth of endometrial carcinoma cells by targeting aurora-B, *FEBS Lett.* 587 (2013) 2523–2529, <http://dx.doi.org/10.1016/j.febslet.2013.05.065>.
- [78] J. Takamizawa, H. Konishi, K. Yanagisawa, S. Tomida, H. Osada, H. Endoh, et al., Reduced expression of the let-7 microRNAs in human lung cancers in association with shortened postoperative survival, *Cancer Res.* 64 (2004) 3753–3756, <http://dx.doi.org/10.1158/0008-5472.CAN-04-0637>.
- [79] Y. Akao, Y. Nakagawa, T. Naoe, let-7 microRNA functions as a potential growth suppressor in human colon cancer cells, *Biol. Pharm. Bull.* 29 (2006) 903–906.
- [80] F. Yu, H. Yao, P. Zhu, X. Zhang, Q. Pan, C. Gong, et al., let-7 regulates self renewal and tumorigenicity of breast cancer cells, *Cell* 131 (2007) 1109–1123, <http://dx.doi.org/10.1016/j.cell.2007.10.054>.
- [81] H. Masai, K.-I. Arai, Cdc7 kinase complex: a key regulator in the initiation of DNA replication, *J. Cell. Physiol.* 190 (2002) 287–296, <http://dx.doi.org/10.1002/jcp.10070>.
- [82] S. Tanaka, H. Araki, Regulation of the initiation step of DNA replication by cyclin-dependent kinases, *Chromosoma* 119 (2010) 565–574, <http://dx.doi.org/10.1007/s00412-010-0291-8>.
- [83] P. Tenca, D. Brotherton, A. Montagnoli, S. Rainoldi, C. Albanese, C. Santocanale, Cdc7 is an active kinase in human cancer cells undergoing replication stress, *J. Biol. Chem.* 282 (2007) 208–215, <http://dx.doi.org/10.1074/jbc.M604457200>.
- [84] J.M. Kim, N. Kakusho, M. Yamada, Y. Kanoh, N. Takemoto, H. Masai, Cdc7 kinase mediates Claspin phosphorylation in DNA replication checkpoint, *Oncogene* 27 (2008) 3475–3482, <http://dx.doi.org/10.1038/sj.onc.1210994>.
- [85] T.A. Day, K. Palle, L.R. Barkley, N. Kakusho, Y. Zou, S. Tateishi, et al., Phosphorylated Rad18 directs DNA polymerase  $\eta$  to sites of stalled replication, *J. Cell Biol.* 191 (2010) 953–966, <http://dx.doi.org/10.1083/jcb.201006043>.
- [86] D. Bonte, C. Lindvall, H. Liu, K. Dykema, K. Furge, M. Weinreich, Cdc7-Dbf4 kinase overexpression in multiple cancers and tumor cell lines is correlated with p53 inactivation, *Neoplasia* N. Y. 10 (2008) 920–931.
- [87] A. Montagnoli, J. Moll, F. Colotta, Targeting cell division cycle 7 kinase: a new approach for cancer therapy, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 16 (2010) 4503–4508, <http://dx.doi.org/10.1158/1078-0432.CCR-10-0185>.
- [88] L.R. Barkley, C. Santocanale, MicroRNA-29a regulates the benzo[a]pyrene dihydrodiol epoxide-induced DNA damage response through Cdc7 kinase in lung cancer cells, *Oncogenesis* 2 (2013) e57, <http://dx.doi.org/10.1038/oncsis.2013.20>.
- [89] D.M. Glover, I.M. Hagan, A.A. Tavares, Polo-like kinases: a team that plays throughout mitosis, *Genes Dev.* 12 (1998) 3777–3787.
- [90] G. Wolf, R. Hildenbrand, C. Schwar, R. Grobholz, M. Kaufmann, H.J. Stutte, et al., Polo-like kinase: a novel marker of proliferation: correlation with estrogen-receptor expression in human breast cancer, *Pathol. Res. Pract.* 196 (2000) 753–759, [http://dx.doi.org/10.1016/S0344-0338\(00\)80107-7](http://dx.doi.org/10.1016/S0344-0338(00)80107-7).



- [91] N. Takai, T. Miyazaki, K. Fujisawa, K. Nasu, R. Hamanaka, I. Miyakawa, Expression of polo-like kinase in ovarian cancer is associated with histological grade and clinical stage, *Cancer Lett.* 164 (2001) 41–49.
- [92] N. Takai, T. Miyazaki, K. Fujisawa, K. Nasu, R. Hamanaka, I. Miyakawa, Polo-like kinase (PLK) expression in endometrial carcinoma, *Cancer Lett.* 169 (2001) 41–49.
- [93] P.J. Gray Jr., D.J. Bearss, H. Han, R. Nagle, M.-S. Tsao, N. Dean, et al., Identification of human polo-like kinase 1 as a potential therapeutic target in pancreatic cancer, *Mol. Cancer Ther.* 3 (2004) 641–646.
- [94] G. Zhang, Z. Zhang, Z. Liu, Polo-like kinase 1 is overexpressed in renal cancer and participates in the proliferation and invasion of renal cancer cells, *Tumour Biol. J. Int. Soc. Oncodevelopmental Biol. Med.* 34 (2013) 1887–1894, <http://dx.doi.org/10.1007/s13277-013-0732-0>.
- [95] T. Takahashi, B. Sano, T. Nagata, H. Kato, Y. Sugiyama, K. Kunieda, et al., Polo-like kinase 1 (PLK1) is overexpressed in primary colorectal cancers, *Cancer Sci.* 94 (2003) 148–152.
- [96] L. Kneisel, K. Strebhardt, A. Bernd, M. Wolter, A. Binder, R. Kaufmann, Expression of polo-like kinase (PLK1) in thin melanomas: a novel marker of metastatic disease, *J. Cutan. Pathol.* 29 (2002) 354–358.
- [97] Y. Degenhardt, T. Lampkin, Targeting Polo-like kinase in cancer therapy, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 16 (2010) 384–389, <http://dx.doi.org/10.1158/1078-0432.CCR-09-1380>.
- [98] B. Spänkuch-Schmitt, G. Wolf, C. Solbach, S. Loibl, R. Knecht, M. Stegmüller, et al., Downregulation of human polo-like kinase activity by antisense oligonucleotides induces growth inhibition in cancer cells, *Oncogene* 21 (2002) 3162–3171, <http://dx.doi.org/10.1038/sj.onc.1205412>.
- [99] K. Gumireddy, M.V.R. Reddy, S.C. Cosenza, R. Boominathan, R. Boomi Nathan, S.J. Baker, et al., ONO1910, a non-ATP-competitive small molecule inhibitor of Plk1, is a potent anticancer agent, *Cancer Cell* 7 (2005) 275–286, <http://dx.doi.org/10.1016/j.ccr.2005.02.009>.
- [100] M. Nogawa, T. Yuasa, S. Kimura, M. Tanaka, J. Kuroda, K. Sato, et al., Intravesical administration of small interfering RNA targeting PLK-1 successfully prevents the growth of bladder cancer, *J. Clin. Invest.* 115 (2005) 978–985, <http://dx.doi.org/10.1172/JCI23043>.
- [101] S. Reagan-Shaw, N. Ahmad, Silencing of polo-like kinase (Plk) 1 via siRNA causes induction of apoptosis and impairment of mitosis machinery in human prostate cancer cells: implications for the treatment of prostate cancer, *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 19 (2005) 611–613, <http://dx.doi.org/10.1096/fj.04-2910fje>.
- [102] M. Steegmaier, M. Hoffmann, A. Baum, P. Lénárt, M. Petronczki, M. Krssák, et al., BI 2536, a potent and selective inhibitor of polo-like kinase 1, inhibits tumor growth in vivo, *Curr. Biol. CB* 17 (2007) 316–322, <http://dx.doi.org/10.1016/j.cub.2006.12.037>.
- [103] A.T. Tandle, T. Kramp, W.J. Kil, A. Halthore, K. Gehlhaus, U. Shankavaram, et al., Inhibition of polo-like kinase 1 in glioblastoma multiforme induces mitotic catastrophe and enhances radiosensitisation, *Eur. J. Cancer Oxf. Engl.* 1990 (49) (2013) 3020–3028, <http://dx.doi.org/10.1016/j.ejca.2013.05.013>.
- [104] F. Biagioni, N. Bossel Ben-Moshe, G. Fontemaggi, V. Canu, F. Mori, B. Antoniani, et al., miR-10b\*, a master inhibitor of the cell cycle, is down-regulated in human breast tumours, *EMBO Mol. Med.* 4 (2012) 1214–1229, <http://dx.doi.org/10.1002/emmm.201201483>.
- [105] W. Shi, N.M. Alajez, C. Bastianutto, A.B.Y. Hui, J.D. Mocanu, E. Ito, et al., Significance of Plk1 regulation by miR-100 in human nasopharyngeal cancer, *Int. J. Cancer* 126 (2010) 2036–2048, <http://dx.doi.org/10.1002/ijc.24880>.
- [106] J. Liu, K.-H. Lu, Z.-L. Liu, M. Sun, W. De, Z.-X. Wang, MicroRNA-100 is a potential molecular marker of non-small cell lung cancer and functions as a tumor suppressor by targeting polo-like kinase 1, *BMC Cancer* 12 (2012) 519, <http://dx.doi.org/10.1186/1471-2407-12-519>.
- [107] T. Ito, F. Sato, T. Kan, Y. Cheng, S. David, R. Agarwal, et al., Polo-like kinase 1 regulates cell proliferation and is targeted by miR-593\* in esophageal cancer, *Int. J. Cancer* 129 (2011) 2134–2146, <http://dx.doi.org/10.1002/ijc.25874>.
- [108] C.H. McGowan, P. Russell, Human Wee1 kinase inhibits cell division by phosphorylating p34cdc2 exclusively on Tyr15, *EMBO J.* 12 (1993) 75–85.
- [109] K. Mahajan, B. Fang, J.M. Koomen, N.P. Mahajan, H2B Tyr37 phosphorylation suppresses expression of replication-dependent core histone genes, *Nat. Struct. Mol. Biol.* 19 (2012) 930–937, <http://dx.doi.org/10.1038/nsmb.2356>.
- [110] H. Butz, I. Likó, S. Cziráj, P. Igaz, M.M. Khan, V. Zivkovic, et al., Down-regulation of Wee1 kinase by a specific subset of microRNA in human sporadic pituitary adenomas, *J. Clin. Endocrinol. Metab.* 95 (2010) E181–E191, <http://dx.doi.org/10.1210/jc.2010-0581>.
- [111] L. Creevey, J. Ryan, H. Harvey, I.M. Bray, M. Meehan, A.R. Khan, et al., MicroRNA-497 increases apoptosis in MYCN amplified neuroblastoma cells by targeting the key cell cycle regulator WEE1, *Mol. Cancer* 12 (2013) 23, <http://dx.doi.org/10.1186/1476-4598-12-23>.
- [112] D. Li, Y. Zhao, C. Liu, X. Chen, Y. Qi, Y. Jiang, et al., Analysis of MiR-195 and MiR-497 expression, regulation and role in breast cancer, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 17 (2011) 1722–1730, <http://dx.doi.org/10.1158/1078-0432.CCR-10-1800>.
- [113] S.T. Guo, C.C. Jiang, G.P. Wang, Y.P. Li, C.Y. Wang, X.Y. Guo, et al., MicroRNA-497 targets insulin-like growth factor 1 receptor and has a tumour suppressive role in human colorectal cancer, *Oncogene* 32 (2013) 1910–1920, <http://dx.doi.org/10.1038/ncr.2012.214>.
- [114] W.L. Ng, D. Yan, X. Zhang, Y.-Y. Mo, Y. Wang, Over-expression of miR-100 is responsible for the low-expression of ATM in the human glioma cell line: M059], *DNA Repair* 9 (2010) 1170–1175, <http://dx.doi.org/10.1016/j.dnarep.2010.08.007>.
- [115] D. Yan, W.L. Ng, X. Zhang, P. Wang, Z. Zhang, Y.-Y. Mo, et al., Targeting DNA-PKcs and ATM with miR-101 sensitizes tumors to radiation, *PLoS One* 5 (2010) e11397, <http://dx.doi.org/10.1371/journal.pone.0011397>.
- [116] H. Hu, L. Du, G. Nagabayashi, R.C. Seeger, R.A. Gatti, ATM is down-regulated by N-Myc-regulated microRNA-421, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 1506–1511, <http://dx.doi.org/10.1073/pnas.0907763107>.
- [117] J. Wang, J. He, F. Su, N. Ding, W. Hu, B. Yao, et al., Repression of ATR pathway by miR-185 enhances radiation-induced apoptosis and proliferation inhibition, *Cell Death Dis.* 4 (2013) e699, <http://dx.doi.org/10.1038/cddis.2013.227>.
- [118] B. Zhao, L. Li, Q. Lei, K.-L. Guan, The Hippo-YAP pathway in organ size control and tumorigenesis: an updated version, *Genes Dev.* 24 (2010) 862–874, <http://dx.doi.org/10.1101/gad.1909210>.
- [119] K.F. Harvey, X. Zhang, D.M. Thomas, The Hippo pathway and human cancer, *Nat. Rev. Cancer* 13 (2013) 246–257, <http://dx.doi.org/10.1038/nrc3458>.
- [120] W. Qin, P. Dong, C. Ma, K. Mitchelson, T. Deng, L. Zhang, et al., MicroRNA-133b is a key promoter of cervical carcinoma development through the activation of the ERK and AKT1 pathways, *Oncogene* 31 (2012) 4067–4075, <http://dx.doi.org/10.1038/ncr.2011.561>.
- [121] Y. Li, J. Pei, H. Xia, H. Ke, H. Wang, W. Tao, Lats2, a putative tumor suppressor, inhibits G1/S transition, *Oncogene* 22 (2003) 4398–4405, <http://dx.doi.org/10.1038/sj.onc.1206603>.
- [122] H. Ke, J. Pei, Z. Ni, H. Xia, H. Qi, T. Woods, et al., Putative tumor suppressor Lats2 induces apoptosis through downregulation of Bcl-2 and Bcl-x(L), *Exp. Cell Res.* 298 (2004) 329–338, <http://dx.doi.org/10.1016/j.yexcr.2004.04.031>.
- [123] K. Tschöp, A.R. Conery, L. Litovchick, J.A. Decaprio, J. Settleman, E. Harlow, et al., A kinase shRNA screen links LATS2 and the pRB tumor suppressor, *Genes Dev.* 25 (2011) 814–830, <http://dx.doi.org/10.1101/gad.2000211>.
- [124] A. Jiménez-Velasco, J. Román-Gómez, X. Agirre, M. Barrios, G. Navarro, I. Vázquez, et al., Downregulation of the large tumor suppressor 2 (LATS2/KPM) gene is associated with poor prognosis in acute lymphoblastic leukemia, *Leukemia* 19 (2005) 2347–2350, <http://dx.doi.org/10.1038/sj.leu.2403974>.
- [125] Y. Takahashi, Y. Miyoshi, C. Takahata, N. Irahara, T. Taguchi, Y. Tamaki, et al., Down-regulation of LATS1 and LATS2 mRNA expression by promoter hypermethylation and its association with biologically aggressive phenotype in human breast cancers, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 11 (2005) 1380–1385, <http://dx.doi.org/10.1158/1078-0432.CCR-04-1773>.
- [126] Z. Jiang, X. Li, J. Hu, W. Zhou, Y. Jiang, G. Li, et al., Promoter hypermethylation-mediated down-regulation of LATS1 and LATS2 in human astrocytoma, *Neurosci. Res.* 56 (2006) 450–458, <http://dx.doi.org/10.1016/j.neures.2006.09.006>.
- [127] X. Liu, L.F. Sempere, H. Ouyang, V.A. Memoli, A.S. Andrew, Y. Luo, et al., MicroRNA-31 functions as an oncogenic microRNA in mouse and human cancer cells by repressing specific tumor suppressors, *J. Clin. Invest.* 120 (2010) 1298–1309, <http://dx.doi.org/10.1172/JCI39566>.
- [128] P.M. Voorhoeve, C. le Sage, M. Schrier, A.J.M. Gillis, H. Stoop, R. Nagel, et al., A genetic screen implicates miRNA-372 and miRNA-373 as oncogenes in testicular germ cell tumors, *Cell* 124 (2006) 1169–1181, <http://dx.doi.org/10.1016/j.cell.2006.02.037>.
- [129] W.J. Cho, J.M. Shin, J.S. Kim, M.R. Lee, K.S. Hong, J.-H. Lee, et al., miR-372 regulates cell cycle and apoptosis of ags human gastric cancer cell line through direct regulation of LATS2, *Mol. Cells* 28 (2009) 521–527, <http://dx.doi.org/10.1007/s10059-009-0158-0>.
- [130] K.-H. Lee, Y.-G. Goan, M. Hsiao, C.-H. Lee, S.-H. Jian, J.-T. Lin, et al., MicroRNA-373 (miR-373) post-transcriptionally regulates large tumor suppressor, homolog 2 (LATS2) and stimulates proliferation in human esophageal cancer, *Exp. Cell Res.* 315 (2009) 2529–2538, <http://dx.doi.org/10.1016/j.yexcr.2009.06.001>.
- [131] M.J. Bueno, I. Pérez de Castro, M. Gómez de Cedrón, J. Santos, G.A. Calin, J.C. Cigudosa, et al., Genetic and epigenetic silencing of MicroRNA-203 enhances ABL1 and BCR-ABL1 oncogene expression, *Cancer Cell* 13 (2008) 496–506, <http://dx.doi.org/10.1016/j.ccr.2008.04.018>.
- [132] Y. Li, H. Wang, K. Tao, Q. Xiao, Z. Huang, L. Zhong, et al., miR-29b suppresses CML cell proliferation and induces apoptosis via regulation of BCR/ABL1 protein, *Exp. Cell Res.* 319 (2013) 1094–1101, <http://dx.doi.org/10.1016/j.yexcr.2013.02.002>.
- [133] Y. Liu, Y. Song, W. Ma, W. Zheng, H. Yin, Decreased microRNA-30a levels are associated with enhanced ABL1 and BCR-ABL1 expression in chronic myeloid leukemia, *Leuk. Res.* 37 (2013) 349–356, <http://dx.doi.org/10.1016/j.leukres.2012.12.003>.
- [134] N. Duquesnes, F. Lezoualc'h, B. Crozatier, PKC-delta and PKC-epsilon: foes of the same family or strangers? *J. Mol. Cell. Cardiol.* 51 (2011) 665–673, <http://dx.doi.org/10.1016/j.yjmcc.2011.07.013>.
- [135] C. Körner, I. Keklikoglou, C. Bender, A. Wörner, E. Münstermann, S. Wiemann, MicroRNA-31 sensitizes human breast cells to apoptosis by direct targeting of protein kinase C epsilon (PKCepsilon), *J. Biol. Chem.* 288 (2013) 8750–8761, <http://dx.doi.org/10.1074/jbc.M112.414128>.
- [136] A. Carnero, The PKB/AKT pathway in cancer, *Curr. Pharm. Des.* 16 (2010) 34–44.
- [137] K.M. Nicholson, N.G. Anderson, The protein kinase B/Akt signalling pathway in human malignancy, *Cell Signal.* 14 (2002) 381–395.

- [138] N. Cai, Y.-D. Wang, P.-S. Zheng, The microRNA-302-367 cluster suppresses the proliferation of cervical carcinoma cells through the novel target AKT1, *RNA N. Y.* 19 (2013) 85–95, <http://dx.doi.org/10.1261/rna.035295.112>.
- [139] Z.-H. Tao, J.-L. Wan, L.-Y. Zeng, L. Xie, H.-C. Sun, L.-X. Qin, et al., miR-612 suppresses the invasive-metastatic cascade in hepatocellular carcinoma, *J. Exp. Med.* 210 (2013) 789–803, <http://dx.doi.org/10.1084/jem.20120153>.
- [140] M. Laplante, D.M. Sabatini, mTOR signaling in growth control and disease, *Cell* 149 (2012) 274–293, <http://dx.doi.org/10.1016/j.cell.2012.03.017>.
- [141] J. Sun, Z. Chen, X. Tan, F. Zhou, F. Tan, Y. Gao, et al., MicroRNA-99a/100 promotes apoptosis by targeting mTOR in human esophageal squamous cell carcinoma, *Med. Oncol. Northwood Lond. Engl.* 30 (2013) 411, <http://dx.doi.org/10.1007/s12032-012-0411-9>.
- [142] D.A. Guertin, D.M. Sabatini, An expanding role for mTOR in cancer, *Trends Mol. Med.* 11 (2005) 353–361, <http://dx.doi.org/10.1016/j.molmed.2005.06.007>.
- [143] L. Cui, H. Zhou, H. Zhao, Y. Zhou, R. Xu, X. Xu, et al., MicroRNA-99a induces G1-phase cell cycle arrest and suppresses tumorigenicity in renal cell carcinoma, *BMC Cancer* 12 (2012) 546, <http://dx.doi.org/10.1186/1471-2407-12-546>.
- [144] T. Iwaya, T. Yokobori, N. Nishida, R. Kogo, T. Sudo, F. Tanaka, et al., Down-regulation of miR-144 is associated with colorectal cancer progression via activation of mTOR signaling pathway, *Carcinogenesis* 33 (2012) 2391–2397, <http://dx.doi.org/10.1093/carcin/bgs288>.
- [145] Y.D. Saito, A.R. Jensen, R. Salgia, E.M. Posadas, Fyn: a novel molecular target in cancer, *Cancer* 116 (2010) 1629–1637, <http://dx.doi.org/10.1002/cncr.24879>.
- [146] L. Ninio-Many, H. Grossman, N. Shomron, D. Chuderland, R. Shalgi, microRNA-125a-3p reduces cell proliferation and migration by targeting Fyn, *J. Cell Sci.* 126 (2013) 2867–2876, <http://dx.doi.org/10.1242/jcs.123414>.
- [147] J.T. Parsons, K.H. Martin, J.K. Slack, J.M. Taylor, S.A. Weed, Focal adhesion kinase: a regulator of focal adhesion dynamics and cell movement, *Oncogene* 19 (2000) 5606–5613, <http://dx.doi.org/10.1038/sj.onc.1203877>.
- [148] W.W. Ma, Development of focal adhesion kinase inhibitors in cancer therapy, *Anticancer Agents Med. Chem.* 11 (2011) 638–642.
- [149] X. Kong, G. Li, Y. Yuan, Y. He, X. Wu, W. Zhang, et al., MicroRNA-7 inhibits epithelial-to-mesenchymal transition and metastasis of breast cancer cells via targeting FAK expression, *PLoS One* 7 (2012) e41523, <http://dx.doi.org/10.1371/journal.pone.0041523>.
- [150] D. Wu, Y. Wang, L. Fan, H. Luo, B. Han, L. Sun, et al., MicroRNA-7 regulates glioblastoma cell invasion via targeting focal adhesion kinase expression, *Chin. Med. J. (Engl.)* 124 (2011) 2616–2621.
- [151] J.C. Loftus, J.T.D. Ross, K.M. Paquette, V.M. Paulino, S. Nasser, Z. Yang, et al., miRNA expression profiling in migrating glioblastoma cells: regulation of cell migration and invasion by miR-23b via targeting of Pyk2, *PLoS One* 7 (2012) e39818, <http://dx.doi.org/10.1371/journal.pone.0039818>.
- [152] R.-F. Liu, X. Xu, J. Huang, Q.-L. Fei, F. Chen, Y.-D. Li, et al., Down-regulation of miR-517a and miR-517c promotes proliferation of hepatocellular carcinoma cells via targeting Pyk2, *Cancer Lett.* 329 (2013) 164–173, <http://dx.doi.org/10.1016/j.canlet.2012.10.027>.
- [153] G.M. Bokoch, Biology of the p21-activated kinases, *Annu. Rev. Biochem.* 72 (2003) 743–781, <http://dx.doi.org/10.1146/annurev.biochem.72.121801.161742>.
- [154] R. Kumar, A.E. Gururaj, C.J. Barnes, p21-activated kinases in cancer, *Nat. Rev. Cancer* 6 (2006) 459–471, <http://dx.doi.org/10.1038/nrc1892>.
- [155] S. Balasenthil, A.A. Sahin, C.J. Barnes, R.-A. Wang, R.G. Pestell, R.K. Vadlamudi, et al., p21-activated kinase-1 signaling mediates cyclin D1 expression in mammary epithelial and cancer cells, *J. Biol. Chem.* 279 (2004) 1422–1428, <http://dx.doi.org/10.1074/jbc.M309937200>.
- [156] M.G. Callow, F. Clairvoyant, S. Zhu, B. Schryver, D.B. Whyte, J.R. Bischoff, et al., Requirement for PAK4 in the anchorage-independent growth of human cancer cell lines, *J. Biol. Chem.* 277 (2002) 550–558, <http://dx.doi.org/10.1074/jbc.M105732200>.
- [157] S.D.N. Reddy, K. Ohshiro, S.K. Rayala, R. Kumar, MicroRNA-7, a homeobox D10 target, inhibits p21-activated kinase 1 and regulates its functions, *Cancer Res.* 68 (2008) 8195–8200, <http://dx.doi.org/10.1158/0008-5472.CAN-08-2103>.
- [158] L. Pellegrino, J. Stebbing, V.M. Braga, A.E. Frampton, J. Jacob, L. Buluwela, et al., miR-23b regulates cytoskeletal remodeling, motility and metastasis by directly targeting multiple transcripts, *Nucleic Acids Res.* 41 (2013) 5400–5412, <http://dx.doi.org/10.1093/nar/gkt245>.
- [159] Z. Wang, X. Zhang, Z. Yang, H. Du, Z. Wu, J. Gong, et al., MiR-145 regulates PAK4 via the MAPK pathway and exhibits an antitumor effect in human colon cells, *Biochem. Biophys. Res. Commun.* 427 (2012) 444–449, <http://dx.doi.org/10.1016/j.bbrc.2012.06.123>.
- [160] N. Rath, M.F. Olson, Rho-associated kinases in tumorigenesis: re-considering ROCK inhibition for cancer therapy, *EMBO Rep.* 13 (2012) 900–908, <http://dx.doi.org/10.1038/embor.2012.127>.
- [161] S.-L. Lin, A. Chiang, D. Chang, S.-Y. Ying, Loss of mir-146a function in hormone-refractory prostate cancer, *RNA N. Y.* 14 (2008) 417–424, <http://dx.doi.org/10.1261/rna.874808>.
- [162] D. Cimino, C. De Pittà, F. Orso, M. Zampini, S. Casara, E. Penna, et al., miR148b is a major coordinator of breast cancer progression in a relapse-associated microRNA signature by targeting ITGA5, ROCK1, PIK3CA, NRAS, and CSF1, *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 27 (2013) 1223–1235, <http://dx.doi.org/10.1096/fj.12-214692>.
- [163] J. Lynch, J. Fay, M. Meehan, K. Bryan, K.M. Watters, D.M. Murphy, et al., miRNA-335 suppresses neuroblastoma cell invasiveness by direct targeting of multiple genes from the non-canonical TGF- $\beta$  signalling pathway, *Carcinogenesis* 33 (2012) 976–985, <http://dx.doi.org/10.1093/carcin/bgs114>.
- [164] K. Ueno, H. Hirata, V. Shahryari, Y. Chen, M.S. Zaman, K. Singh, et al., Tumour suppressor microRNA-584 directly targets oncogene Rock-1 and decreases invasion ability in human clear cell renal cell carcinoma, *Br. J. Cancer* 104 (2011) 308–315, <http://dx.doi.org/10.1038/sj.bjc.6606028>.
- [165] S. Majid, A.A. Dar, S. Saini, V. Shahryari, S. Arora, M.S. Zaman, et al., MicroRNA-1280 inhibits invasion and metastasis by targeting ROCK1 in bladder cancer, *PLoS One* 7 (2012) e46743, <http://dx.doi.org/10.1371/journal.pone.0046743>.
- [166] F. Zheng, Y.-J. Liao, M.-Y. Cai, Y.-H. Liu, T.-H. Liu, S.-P. Chen, et al., The putative tumour suppressor microRNA-124 modulates hepatocellular carcinoma cell aggressiveness by repressing ROCK2 and EZH2, *Gut* 61 (2012) 278–289, <http://dx.doi.org/10.1136/gut.2011.239145>.
- [167] L. Jiang, X. Liu, A. Kolokythas, J. Yu, A. Wang, C.E. Heidbreder, et al., Down-regulation of the Rho GTPase signaling pathway is involved in the microRNA-138-mediated inhibition of cell migration and invasion in tongue squamous cell carcinoma, *Int. J. Cancer* 127 (2010) 505–512, <http://dx.doi.org/10.1002/ijc.25320>.
- [168] C.C.-L. Wong, C.-M. Wong, E.K.-K. Tung, S.L.-K. Au, J.M.-F. Lee, R.T.-P. Poon, et al., The microRNA miR-139 suppresses metastasis and progression of hepatocellular carcinoma by down-regulating Rho-kinase 2, *Gastroenterology* 140 (2011) 322–331, <http://dx.doi.org/10.1053/j.gastro.2010.10.006>.
- [169] R. Baserga, F. Peruzzi, K. Reiss, The IGF-1 receptor in cancer biology, *Int. J. Cancer* 107 (2003) 873–877, <http://dx.doi.org/10.1002/ijc.11487>.
- [170] Z. Cao, L.-Z. Liu, D.A. Dixon, J.Z. Zheng, B. Chandran, B.-H. Jiang, Insulin-like growth factor-I induces cyclooxygenase-2 expression via PI3K, MAPK and PKC signaling pathways in human ovarian cancer cells, *Cell Signal.* 19 (2007) 1542–1553, <http://dx.doi.org/10.1016/j.cellsig.2007.01.028>.
- [171] L. Jiang, X. Liu, Z. Chen, Y. Jin, C.E. Heidbreder, A. Kolokythas, et al., MicroRNA-7 targets IGF1R (insulin-like growth factor 1 receptor) in tongue squamous cell carcinoma cells, *Biochem. J.* 432 (2010) 199–205, <http://dx.doi.org/10.1042/BJ20100859>.
- [172] L. Chen, Q. Wang, G. Wang, H. Wang, Y. Huang, X. Liu, et al., miR-16 inhibits cell proliferation by targeting IGF1R and the Raf1-MEK1/2-ERK1/2 pathway in osteosarcoma, *FEBS Lett.* 587 (2013) 1366–1372, <http://dx.doi.org/10.1016/j.febslet.2013.03.007>.
- [173] B. Wang, H. Wang, Z. Yang, MiR-122 inhibits cell proliferation and tumorigenesis of breast cancer by targeting IGF1R, *PLoS One* 7 (2012) e47053, <http://dx.doi.org/10.1371/journal.pone.0047053>.
- [174] Z.-M. Shi, X.-F. Wang, X. Qian, T. Tao, L. Wang, Q.-D. Chen, et al., MiRNA-181b suppresses IGF-1R and functions as a tumor suppressor gene in gliomas, *RNA N. Y.* 19 (2013) 552–560, <http://dx.doi.org/10.1261/rna.035972.112>.
- [175] L. Zehavi, R. Avraham, A. Barzilai, D. Bar-Ilan, R. Navon, Y. Sidi, et al., Silencing of a large microRNA cluster on human chromosome 14q32 in melanoma: biological effects of mir-376a and mir-376c on insulin growth factor 1 receptor, *Mol. Cancer* 11 (2012) 44, <http://dx.doi.org/10.1186/1476-4598-11-44>.
- [176] M. Luo, D. Shen, X. Zhou, X. Chen, W. Wang, MicroRNA-497 is a potential prognostic marker in human cervical cancer and functions as a tumor suppressor by targeting the insulin-like growth factor 1 receptor, *Surgery* 153 (2013) 836–847, <http://dx.doi.org/10.1016/j.surg.2012.12.004>.
- [177] I. Vivanco, C.L. Sawyers, The phosphatidylinositol 3-Kinase AKT pathway in human cancer, *Nat. Rev. Cancer* 2 (2002) 489–501, <http://dx.doi.org/10.1038/nrc839>.
- [178] T.S. Lewis, P.S. Shapiro, N.G. Ahn, Signal transduction through MAP kinase cascades, *Adv. Cancer Res.* 74 (1998) 49–139.
- [179] F.C. Kalinowski, K.M. Giles, P.A. Candy, A. Ali, C. Ganda, M.R. Epis, et al., Regulation of epidermal growth factor receptor signaling and erlotinib sensitivity in head and neck cancer cells by miR-7, *PLoS One* 7 (2012) e47067, <http://dx.doi.org/10.1371/journal.pone.0047067>.
- [180] B. Kefas, J. Godlewski, L. Comeau, Y. Li, R. Abounader, M. Hawkinson, et al., microRNA-7 inhibits the epidermal growth factor receptor and the Akt pathway and is down-regulated in glioblastoma, *Cancer Res.* 68 (2008) 3566–3572, <http://dx.doi.org/10.1158/0008-5472.CAN-07-6639>.
- [181] R.J. Webster, K.M. Giles, K.J. Price, P.M. Zhang, J.S. Mattick, P.J. Leedman, Regulation of epidermal growth factor receptor signaling in human cancer cells by microRNA-7, *J. Biol. Chem.* 284 (2009) 5731–5741, <http://dx.doi.org/10.1074/jbc.M804280200>.
- [182] M. Acunzo, G. Romano, D. Palmieri, A. Laganà, M. Garofalo, V. Balatti, et al., Cross-talk between MET and EGFR in non-small cell lung cancer involves miR-27a and Sprouty2, *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 8573–8578, <http://dx.doi.org/10.1073/pnas.1302107110>.
- [183] W. Wang, B. Cheng, L. Miao, Y. Mei, M. Wu, Mutant p53-R273H gains new function in sustained activation of EGFR signaling via suppressing miR-27a expression, *Cell Death Dis.* 4 (2013) e574, <http://dx.doi.org/10.1038/cddis.2013.97>.
- [184] L. Liu, X. Shao, W. Gao, Z. Zhang, P. Liu, R. Wang, et al., MicroRNA-133b inhibits the growth of non-small-cell lung cancer by targeting the epidermal growth factor receptor, *FEBS J.* 279 (2012) 3800–3812, <http://dx.doi.org/10.1111/j.1742-4658.2012.08741.x>.
- [185] A. Verma, S.L. Warner, H. Vankayalapati, D.J. Bearss, S. Sharma, Targeting Axl and Src kinases in cancer, *Mol. Cancer Ther.* 10 (2011) 1763–1773, <http://dx.doi.org/10.1158/1535-7163.MCT-11-0116>.



- [186] G. Mudduluru, P. Ceppi, R. Kumarswamy, G.V. Scagliotti, M. Papotti, H. Allgayer, Regulation of Axl receptor tyrosine kinase expression by miR-34a and miR-199a/b in solid cancer, *Oncogene* 30 (2011) 2888–2899, <http://dx.doi.org/10.1038/ncr.2011.13>.
- [187] M. Mackiewicz, K. Huppi, J.J. Pitt, T.H. Dorsey, S. Ambs, N.J. Caplen, Identification of the receptor tyrosine kinase AXL in breast cancer as a target for the human miR-34a microRNA, *Breast Cancer Res. Treat.* 130 (2011) 663–679, <http://dx.doi.org/10.1007/s10549-011-1690-0>.
- [188] C. Birchmeier, W. Birchmeier, E. Gherardi, G.F. Vande Woude, Met, metastasis, motility and more, *Nat. Rev. Mol. Cell Biol.* 4 (2003) 915–925, <http://dx.doi.org/10.1038/nrm1261>.
- [189] X. Liu, W. Yao, R.C. Newton, P.A. Scherle, Targeting the c-MET signaling pathway for cancer therapy, *Expert Opin. Invest. Drugs* 17 (2008) 997–1011, <http://dx.doi.org/10.1517/13543784.17.7.997>.
- [190] J.G. Christensen, J. Burrows, R. Sargia, c-Met as a target for human cancer and characterization of inhibitors for therapeutic intervention, *Cancer Lett.* 225 (2005) 1–26, <http://dx.doi.org/10.1016/j.canlet.2004.09.044>.
- [191] D. Yan, X. Zhou, X. Chen, D.-N. Hu, X.D. Dong, J. Wang, et al., MicroRNA-34a inhibits uveal melanoma cell proliferation and migration through down-regulation of c-Met, *Invest. Ophthalmol. Vis. Sci.* 50 (2009) 1559–1565, <http://dx.doi.org/10.1167/jovs.08-2681>.
- [192] Z. Hagman, B.S. Hafliadottir, M. Ansari, M. Persson, A. Bjartell, A. Edsjö, et al., The tumour suppressor miR-34c targets MET in prostate cancer cells, *Br. J. Cancer* 109 (2013) 1271–1278, <http://dx.doi.org/10.1038/bjc.2013.449>.
- [193] Z. Hu, Y. Lin, H. Chen, Y. Mao, J. Wu, Y. Zhu, et al., MicroRNA-101 suppresses motility of bladder cancer cells by targeting c-Met, *Biochem. Biophys. Res. Commun.* 435 (2013) 82–87, <http://dx.doi.org/10.1016/j.bbrc.2013.04.042>.
- [194] C. Luo, P.W. Tetteh, P.R. Merz, E. Dickes, A. Abukiwan, A. Hotz-Wagenblatt, et al., miR-137 inhibits the invasion of melanoma cells through downregulation of multiple oncogenic target genes, *J. Invest. Dermatol.* 133 (2013) 768–775, <http://dx.doi.org/10.1038/jid.2012.357>.
- [195] X. Xu, H. Chen, Y. Lin, Z. Hu, Y. Mao, J. Wu, et al., MicroRNA-409-3p inhibits migration and invasion of bladder cancer cells via targeting c-Met, *Mol. Cells* 36 (2013) 62–68, <http://dx.doi.org/10.1007/s10059-013-0044-7>.
- [196] L. Chen, J. Zhang, Y. Feng, R. Li, X. Sun, W. Du, et al., MiR-410 regulates MET to influence the proliferation and invasion of glioma, *Int. J. Biochem. Cell Biol.* 44 (2012) 1711–1717, <http://dx.doi.org/10.1016/j.biocel.2012.06.027>.
- [197] W. Luo, B. Huang, Z. Li, H. Li, L. Sun, Q. Zhang, et al., MicroRNA-449a is downregulated in non-small cell lung cancer and inhibits migration and invasion by targeting c-Met, *PLoS One* 8 (2013) e64759, <http://dx.doi.org/10.1371/journal.pone.0064759>.
- [198] J.M. Lee, J.K. Yoo, H. Yoo, H.Y. Jung, D.R. Lee, H.C. Jeong, et al., The novel miR-7515 decreases the proliferation and migration of human lung cancer cells by targeting c-Met, *Mol. Cancer Res. MCR* 11 (2013) 43–53, <http://dx.doi.org/10.1158/1541-7786.MCR-12-0355>.
- [199] G. Liu, Y. Sun, P. Ji, X. Li, D. Cogdell, D. Yang, et al., MiR-506 suppresses proliferation and induces senescence by directly targeting the CDK4/6-FOXO1 axis in ovarian cancer, *J. Pathol.* 233 (2014) 308–318, <http://dx.doi.org/10.1002/path.4348>.
- [200] M.T. Di Martino, E. Leone, N. Amodio, U. Foresta, M. Lionetti, M.R. Pitari, et al., Synthetic miR-34a mimics as a novel therapeutic agent for multiple myeloma: in vitro and in vivo evidence, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 18 (2012) 6260–6270, <http://dx.doi.org/10.1158/1078-0432.CCR-12-1708>.
- [201] F. Takeshita, L. Patrawala, M. Osaki, R. Takahashi, Y. Yamamoto, N. Kosaka, et al., Systemic delivery of synthetic microRNA-16 inhibits the growth of metastatic prostate tumors via downregulation of multiple cell-cycle genes, *Mol. Ther. J. Am. Soc. Gene Ther.* 18 (2010) 181–187, <http://dx.doi.org/10.1038/mt.2009.207>.
- [202] M.A. Lampson, I.M. Cheeseman, Sensing centromere tension: aurora B and the regulation of kinetochore function, *Trends Cell Biol.* 21 (2011) 133–140, <http://dx.doi.org/10.1016/j.tcb.2010.10.007>.
- [203] P.C. De Witt Hamer, S.E. Mir, D. Noske, C.J.F. Van Noorden, T. Würdinger, WEE1 kinase targeting combined with DNA-damaging cancer therapy catalyzes mitotic catastrophe, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 17 (2011) 4200–4207, <http://dx.doi.org/10.1158/1078-0432.CCR-10-2537>.
- [204] T. Stankovic, A. Skowronska, The role of ATM mutations and 11q deletions in disease progression in chronic lymphocytic leukemia, *Leuk. Lymphoma* 55 (2014) 1227–1239, <http://dx.doi.org/10.3109/10428194.2013.829919>.
- [205] D.R. Schmidt, S.L. Schreiber, Molecular association between ATR and two components of the nucleosome remodeling and deacetylating complex, HDAC2 and CHD4, *Biochemistry (Moscow)* 38 (1999) 14711–14717.
- [206] M. Fabbro, K. Savage, K. Hobson, A.J. Deans, S.N. Powell, G.A. McArthur, et al., BRCA1-BARD1 complexes are required for p53Ser-15 phosphorylation and a G1/S arrest following ionizing radiation-induced DNA damage, *J. Biol. Chem.* 279 (2004) 31251–31258, <http://dx.doi.org/10.1074/jbc.M405372200>.
- [207] M. Sankunni, R.A. Parikh, D.W. Lewis, W.E. Gooding, W.S. Saunders, S.M. Gollin, Targeted inhibition of ATR or CHEK1 reverses radioresistance in oral squamous cell carcinoma cells with distal chromosome arm 11q loss, *Genes, Chromosomes Cancer* 53 (2014) 129–143, <http://dx.doi.org/10.1002/gcc.22125>.
- [208] N. Chatain, P. Ziegler, D. Fahrenkamp, E. Jost, R. Moriggl, H. Schmitz-Van de Leur, et al., Src family kinases mediate cytoplasmic retention of activated STAT5 in BCR-ABL-positive cells, *Oncogene* 32 (2013) 3587–3597, <http://dx.doi.org/10.1038/ncr.2012.369>.
- [209] A. Slupianek, R. Falinski, P. Znojek, T. Stoklosa, S. Flis, V. Doneddu, et al., BCR-ABL kinase inhibits uracil DNA glycosylase UNG2 to enhance oxidative DNA damage and stimulate genomic instability, *Leukemia* 27 (2013) 629–634, <http://dx.doi.org/10.1038/leu.2012.294>.
- [210] M.E. Sobhia, B.K. Grewal, M.L.S. Paul, J. Patel, A. Kaur, T. Haokip, et al., Protein kinase C inhibitors: a patent review (2010–present), *Expert Opin. Ther. Pat.* 23 (2013) 1451–1468, <http://dx.doi.org/10.1517/13543776.2013.812073>.
- [211] I. Hers, E.E. Vincent, J.M. Tavaré, Akt signalling in health and disease, *Cell Signal.* 23 (2011) 1515–1527, <http://dx.doi.org/10.1016/j.cellsig.2011.05.004>.
- [212] C. Betz, M.N. Hall, Where is mTOR and what is it doing there? *J. Cell Biol.* 203 (2013) 563–574, <http://dx.doi.org/10.1083/jcb.201306041>.
- [213] S. Schenone, C. Brullo, F. Musumeci, M. Biava, F. Falchi, M. Botta, Fyn kinase in brain diseases and cancer: the search for inhibitors, *Curr. Med. Chem.* 18 (2011) 2921–2942.
- [214] S.-T.S. Lim, Nuclear FAK: a new mode of gene regulation from cellular adhesions, *Mol. Cells* 36 (2013) 1–6, <http://dx.doi.org/10.1007/s10059-013-0139-1>.
- [215] J.E. Hall, W. Fu, M.D. Schaller, Focal adhesion kinase: exploring Fak structure to gain insight into function, *Int. Rev. Cell Mol. Biol.* 288 (2011) 185–225, <http://dx.doi.org/10.1016/B978-0-12-386041-5.00005-4>.
- [216] C.A. Lipinski, J.C. Loftus, Targeting Pyk2 for therapeutic intervention, *Expert Opin. Ther. Targets* 14 (2010) 95–108, <http://dx.doi.org/10.1517/14728220903473194>.
- [217] J. Braunger, L. Schleithoff, A.S. Schulz, H. Kessler, R. Lammers, A. Ullrich, et al., Intracellular signaling of the Ufo/Axl receptor tyrosine kinase is mediated mainly by a multi-substrate docking-site, *Oncogene* 14 (1997) 2619–2631, <http://dx.doi.org/10.1038/sj.onc.1201123>.
- [218] L. Trusolino, A. Bertotti, P.M. Comoglio, A signaling adapter function for alpha6beta4 integrin in the control of HGF-dependent invasive growth, *Cell* 107 (2001) 643–654.
- [219] D.A. Ucar, A.T. Magis, D.-H. He, N.J. Lawrence, S.M. Sebti, E. Kurenova, et al., Inhibiting the interaction of cMET and IGF-1R with FAK effectively reduces growth of pancreatic cancer cells in vitro and in vivo, *Anticancer Agents Med. Chem.* 13 (2013) 595–602.
- [220] D.M. Virshup, S. Shenolikar, From promiscuity to precision: protein phosphatases get a makeover, *Mol. Cell* 33 (2009) 537–545, <http://dx.doi.org/10.1016/j.molcel.2009.02.015>.
- [221] C. Lammner, S. Wagerer, R. Saffrich, D. Mertens, W. Ansorge, I. Hoffmann, The cdc25B phosphatase is essential for the G2/M phase transition in human cells, *J. Cell Sci.* 111 (Pt. 16) (1998) 2445–2453.
- [222] A. Lavecchia, C. Di Giovanni, E. Novellino, CDC25A and B dual-specificity phosphatase inhibitors: potential agents for cancer therapy, *Curr. Med. Chem.* 16 (2009) 1831–1849.
- [223] K. Kristjánssdóttir, J. Rudolph, Cdc25 phosphatases and cancer, *Chem. Biol.* 11 (2004) 1043–1051, <http://dx.doi.org/10.1016/j.chembiol.2004.07.007>.
- [224] J. Guo, J. Kleeff, J. Li, J. Ding, J. Hammer, Y. Zhao, et al., Expression and functional significance of CDC25B in human pancreatic ductal adenocarcinoma, *Oncogene* 23 (2004) 71–81, <http://dx.doi.org/10.1038/sj.onc.1206926>.
- [225] Y. Ito, H. Yoshida, T. Urano, Y. Takamura, A. Miya, K. Kuma, et al., Expression of cdc25A and cdc25B phosphatase in breast carcinoma, *Breast Cancer Tokyo Jpn.* 11 (2004) 295–300.
- [226] E.S.W. Ngan, Y. Hashimoto, Z.-Q. Ma, M.-J. Tsai, S.Y. Tsai, Overexpression of Cdc25B, an androgen receptor coactivator, in prostate cancer, *Oncogene* 22 (2003) 734–739, <http://dx.doi.org/10.1038/sj.onc.1206121>.
- [227] I. Takemasa, H. Yamamoto, M. Sekimoto, M. Ohue, S. Noura, Y. Miyake, et al., Overexpression of CDC25B phosphatase as a novel marker of poor prognosis of human colorectal carcinoma, *Cancer Res.* 60 (2000) 3043–3050.
- [228] S. Hernández, X. Bessa, S. Beà, L. Hernández, A. Nadal, C. Mallofré, et al., Differential expression of cdc25 cell-cycle-activating phosphatases in human colorectal carcinoma, *Lab. Invest. J. Tech. Methods Pathol.* 81 (2001) 465–473.
- [229] Y. Kudo, W. Yasui, T. Ue, S. Yamamoto, H. Yokozaki, H. Nikai, et al., Overexpression of cyclin-dependent kinase-activating CDC25B phosphatase in human gastric carcinomas, *Jpn. J. Cancer Res. Gann* 88 (1997) 947–952.
- [230] A.P. Ducruet, A. Vogt, P. Wipf, J.S. Lazo, Dual specificity protein phosphatases: therapeutic targets for cancer and Alzheimer's disease, *Annu. Rev. Pharmacol. Toxicol.* 45 (2005) 725–750, <http://dx.doi.org/10.1146/annurev.pharmtox.45.120403.100040>.
- [231] R. Boutros, V. Lobjois, B. Ducommun, CDC25 phosphatases in cancer cells: key players? Good targets? *Nat. Rev. Cancer* 7 (2007) 495–507, <http://dx.doi.org/10.1038/nrc2169>.
- [232] H. Chen, Y. Lin, Y. Mao, J. Wu, Y. Liu, X. Zheng, et al., MicroRNA-449a acts as a tumor suppressor in human bladder cancer through the regulation of pocket proteins, *Cancer Lett.* 320 (2012) 40–47, <http://dx.doi.org/10.1016/j.canlet.2012.01.027>.
- [233] P. Wang, F. Zou, X. Zhang, H. Li, A. Dulak, R.J. Tomko Jr., et al., microRNA-21 negatively regulates Cdc25A and cell cycle progression in colon cancer cells, *Cancer Res.* 69 (2009) 8157–8165, <http://dx.doi.org/10.1158/0008-5472.CAN-09-1996>.
- [234] S.-T. Liffers, J.B. Munding, M. Vogt, J.D. Kuhlmann, B. Verdoodt, S. Nambiar, et al., MicroRNA-148a is down-regulated in human pancreatic ductal adenocarcinomas and regulates cell survival by targeting CDC25B, *Lab. Invest. J. Tech. Methods Pathol.* 91 (2011) 1472–1479, <http://dx.doi.org/10.1038/abinvest.2011.99>.

- [235] X.-Y. Yu, Z. Zhang, J. Liu, B. Zhan, C.-Z. Kong, MicroRNA-141 is downregulated in human renal cell carcinoma and regulates cell survival by targeting CDC25B, *OncoTargets Ther.* 6 (2013) 349–354, <http://dx.doi.org/10.2147/OTT.S41343>.
- [236] V. Stambolic, A. Suzuki, J.L. de la Pompa, G.M. Brothers, C. Mirtsos, T. Sasaki, et al., Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN, *Cell* 95 (1998) 29–39.
- [237] L.-Y. Zhang, Y. Ho-Fun Lee, A.M.G. Wong, D.L.-W. Kwong, Y.-H. Zhu, S.-S. Dong, et al., MicroRNA-144 promotes cell proliferation, migration and invasion in nasopharyngeal carcinoma through repression of PTEN, *Carcinogenesis* 34 (2013) 454–463, <http://dx.doi.org/10.1093/carcin/bgs346>.
- [238] Z. Wu, B. He, J. He, X. Mao, Upregulation of miR-153 promotes cell proliferation via downregulation of the PTEN tumor suppressor gene in human prostate cancer, *Prostate* 73 (2013) 596–604, <http://dx.doi.org/10.1002/pros.22600>.
- [239] Y.-S. Wang, Y.-H. Wang, H.-P. Xia, S.-W. Zhou, G. Schmid-Bindert, C.-C. Zhou, MicroRNA-214 regulates the acquired resistance to gefitinib via the PTEN/AKT pathway in EGFR-mutant cell lines, *Asian Pac. J. Cancer Prev. APJCP* 13 (2012) 255–260.
- [240] H. Xia, L.L.P.J. Ooi, K.M. Hui, MicroRNA-216a/217-induced epithelial-mesenchymal transition targets PTEN and SMAD7 to promote drug resistance and recurrence of liver cancer, *Hepatol. Baltim. Md.* 58 (2013) 629–641, <http://dx.doi.org/10.1002/hep.26369>.
- [241] G. Zhao, C. Cai, T. Yang, X. Qiu, B. Liao, W. Li, et al., MicroRNA-221 induces cell survival and cisplatin resistance through PI3K/Akt pathway in human osteosarcoma, *PLoS One* 8 (2013) e53906, <http://dx.doi.org/10.1371/journal.pone.0053906>.
- [242] S. Zhou, D. Shen, Y. Wang, L. Gong, X. Tang, B. Yu, et al., microRNA-222 targeting PTEN promotes neurite outgrowth from adult dorsal root ganglion neurons following sciatic nerve transection, *PLoS One* 7 (2012) e44768, <http://dx.doi.org/10.1371/journal.pone.0044768>.
- [243] W. Shi, K. Gerster, N.M. Alajez, J. Tsang, L. Waldron, M. Pintilie, et al., MicroRNA-301 mediates proliferation and invasion in human breast cancer, *Cancer Res.* 71 (2011) 2926–2937, <http://dx.doi.org/10.1158/0008-5472.CAN-10-3369>.
- [244] J. Cai, H. Guan, L. Fang, Y. Yang, X. Zhu, J. Yuan, et al., MicroRNA-374a activates Wnt/ $\beta$ -catenin signaling to promote breast cancer metastasis, *J. Clin. Invest.* 123 (2013) 566–579, <http://dx.doi.org/10.1172/JCI65871>.
- [245] S. Volinia, G.A. Calin, C.-G. Liu, S. Ambros, A. Cimmino, F. Petrocca, et al., A microRNA expression signature of human solid tumors defines cancer gene targets, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 2257–2261, <http://dx.doi.org/10.1073/pnas.0510565103>.
- [246] F. Meng, R. Henson, H. Wehbe-Janeke, K. Ghoshal, S.T. Jacob, T. Patel, MicroRNA-21 regulates expression of the PTEN tumor suppressor gene in human hepatocellular cancer, *Gastroenterology* 133 (2007) 647–658, <http://dx.doi.org/10.1053/j.gastro.2007.05.022>.
- [247] S. Zhu, M.-L. Si, H. Wu, Y.-Y. Mo, MicroRNA-21 targets the tumor suppressor gene tropomyosin 1 (TPM1), *J. Biol. Chem.* 282 (2007) 14328–14336, <http://dx.doi.org/10.1074/jbc.M611393200>.
- [248] I.A. Asangani, S.A.K. Rasheed, D.A. Nikolova, J.H. Leupold, N.H. Colburn, S. Post, et al., MicroRNA-21 (miR-21) post-transcriptionally downregulates tumor suppressor Pdc4d and stimulates invasion, intravasation and metastasis in colorectal cancer, *Oncogene* 27 (2008) 2128–2136, <http://dx.doi.org/10.1038/sj.onc.1210856>.
- [249] S. Zhu, H. Wu, F. Wu, D. Nie, S. Sheng, Y.-Y. Mo, MicroRNA-21 targets tumor suppressor genes in invasion and metastasis, *Cell Res.* 18 (2008) 350–359, <http://dx.doi.org/10.1038/cr.2008.24>.
- [250] G. Gabrieli, T. Wurdinger, S. Kesari, C.C. Esau, J. Burchard, P.S. Linsley, et al., MicroRNA 21 promotes glioma invasion by targeting matrix metalloproteinase regulators, *Mol. Cell. Biol.* 28 (2008) 5369–5380, <http://dx.doi.org/10.1128/MCB.00479-08>.
- [251] X. Tan, S. Wang, L. Zhu, C. Wu, B. Yin, J. Zhao, et al., cAMP response element-binding protein promotes gliomagenesis by modulating the expression of oncogenic microRNA-23a, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 15805–15810, <http://dx.doi.org/10.1073/pnas.1207787109>.
- [252] J.T. Huse, C. Brennan, D. Hambardzumyan, B. Wee, J. Pena, S.H. Rouhanifard, et al., The PTEN-regulating microRNA miR-26a is amplified in high-grade glioma and facilitates gliomagenesis in vivo, *Genes Dev.* 23 (2009) 1327–1337, <http://dx.doi.org/10.1101/gad.1777409>.
- [253] A.M. Cheng, M.W. Byrom, J. Shelton, L.P. Ford, Antisense inhibition of human miRNAs and indications for an involvement of miRNA in cell growth and apoptosis, *Nucleic Acids Res.* 33 (2005) 1290–1297, <http://dx.doi.org/10.1093/nar/gki200>.
- [254] H. Yang, W. Kong, L. He, J.-J. Zhao, J.D. O'Donnell, J. Wang, et al., MicroRNA expression profiling in human ovarian cancer: miR-214 induces cell survival and cisplatin resistance by targeting PTEN, *Cancer Res.* 68 (2008) 425–433, <http://dx.doi.org/10.1158/0008-5472.CAN-07-2488>.
- [255] M. Kato, S. Putta, M. Wang, H. Yuan, L. Lanting, I. Nair, et al., TGF-beta activates Akt kinase through a microRNA-dependent amplifying circuit targeting PTEN, *Nat. Cell Biol.* 11 (2009) 881–889, <http://dx.doi.org/10.1038/ncb1897>.
- [256] T. Tsunoda, Y. Takashima, Y. Yoshida, K. Doi, Y. Tanaka, T. Fujimoto, et al., Oncogenic KRAS regulates miR-200c and miR-221/222 in a 3D-specific manner in colorectal cancer cells, *Anticancer Res.* 31 (2011) 2453–2459.
- [257] T. Yuan, Y. Wang, Z.J. Zhao, H. Gu, Protein-tyrosine phosphatase PTPN9 negatively regulates ErbB2 and epidermal growth factor receptor signaling in breast cancer cells, *J. Biol. Chem.* 285 (2010) 14861–14870, <http://dx.doi.org/10.1074/jbc.M109.099879>.
- [258] W.W. Du, L. Fang, M. Li, X. Yang, Y. Liang, C. Peng, et al., MicroRNA miR-24 enhances tumor invasion and metastasis by targeting PTPN9 and PTPRF to promote EGF signaling, *J. Cell Sci.* 126 (2013) 1440–1453, <http://dx.doi.org/10.1242/jcs.118299>.
- [259] C. Zhou, G. Liu, L. Wang, Y. Lu, L. Yuan, L. Zheng, et al., MiR-339-5p regulates the growth, colony formation and metastasis of colorectal cancer cells by targeting PRL-1, *PLoS One* 8 (2013) e63142, <http://dx.doi.org/10.1371/journal.pone.0063142>.
- [260] N. Wu, X. Liu, X. Xu, X. Fan, M. Liu, X. Li, et al., MicroRNA-373, a new regulator of protein phosphatase 6, functions as an oncogene in hepatocellular carcinoma, *FEBS J.* 278 (2011) 2044–2054, <http://dx.doi.org/10.1111/j.1742-4658.2011.08120.x>.
- [261] L. Lessard, M. Stuiblé, M.L. Tremblay, The two faces of PTP1B in cancer, *Biochim. Biophys. Acta* 1804 (2010) 613–619, <http://dx.doi.org/10.1016/j.bbapap.2009.09.018>.
- [262] S. Zhu, J.D. Bjorge, D.J. Fujita, PTP1B contributes to the oncogenic properties of colon cancer cells through Src activation, *Cancer Res.* 67 (2007) 10129–10137, <http://dx.doi.org/10.1158/0008-5472.CAN-06-4338>.
- [263] L.L. Christensen, H. Tobiasen, A. Holm, T. Schepele, M.S. Ostensfeld, K. Thorsen, et al., MiRNA-362-3p induces cell cycle arrest through targeting of E2F1, USF2 and PTPN1 and is associated with recurrence of colorectal cancer, *Int. J. Cancer* 133 (2013) 67–78, <http://dx.doi.org/10.1002/ijc.28010>.
- [264] X. Lu, T.-A. Nguyen, S.-H. Moon, Y. Darlington, M. Sommer, L.A. Donehower, The type 2C phosphatase Wip1: an oncogenic regulator of tumor suppressor and DNA damage response pathways, *Cancer Metastasis Rev.* 27 (2008) 123–135, <http://dx.doi.org/10.1007/s10555-008-9127-x>.
- [265] X. Lu, B. Nannenga, L.A. Donehower, PPM1D dephosphorylates Chk1 and p53 and abrogates cell cycle checkpoints, *Genes Dev.* 19 (2005) 1162–1174, <http://dx.doi.org/10.1101/gad.1291305>.
- [266] H. Fujimoto, N. Onishi, N. Kato, M. Takekawa, X.Z. Xu, A. Kosugi, et al., Regulation of the antioncogenic Chk2 kinase by the oncogenic Wip1 phosphatase, *Cell Death Differ.* 13 (2006) 1170–1180, <http://dx.doi.org/10.1038/sj.cdd.4401801>.
- [267] S. Shreeram, O.N. Demidov, W.K. Hee, H. Yamaguchi, N. Onishi, C. Kek, et al., Wip1 phosphatase modulates ATM-dependent signaling pathways, *Mol. Cell* 23 (2006) 757–764, <http://dx.doi.org/10.1016/j.molcel.2006.07.010>.
- [268] J. Li, Y. Yang, Y. Peng, R.J. Austin, W.G. van Eyndhoven, K.C.Q. Nguyen, et al., Oncogenic properties of PPM1D located within a breast cancer amplification epicenter at 17q23, *Nat. Genet.* 31 (2002) 133–134, <http://dx.doi.org/10.1038/ng888>.
- [269] A. Hirasawa, F. Saito-Ohara, J. Inoue, D. Aoki, N. Susumu, T. Yokoyama, et al., Association of 17q21-q24 gain in ovarian clear cell adenocarcinomas with poor prognosis and identification of PPM1D and APPBP2 as likely amplification targets, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 9 (2003) 1995–2004.
- [270] F. Saito-Ohara, I. Imoto, J. Inoue, H. Hosoi, A. Nakagawara, T. Sugimoto, et al., PPM1D is a potential target for 17q gain in neuroblastoma, *Cancer Res.* 63 (2003) 1876–1883.
- [271] A. Ehrbrecht, U. Müller, M. Wolter, A. Hoischen, A. Koch, B. Radlwimmer, et al., Comprehensive genomic analysis of desmoplastic medulloblastomas: identification of novel amplified genes and separate evaluation of the different histological components, *J. Pathol.* 208 (2006) 554–563, <http://dx.doi.org/10.1002/path.1925>.
- [272] T. Fukui, S. Semba, H. Yutori, H. Yokozaki, Increased wild-type p53-induced phosphatase 1 (Wip1 or PPM1D) expression correlated with down-regulation of checkpoint kinase 2 in human gastric carcinoma, *Pathol. Int.* 57 (2007) 566–571, <http://dx.doi.org/10.1111/j.1440-1827.2007.02140.x>.
- [273] P. Loukopoulou, T. Shibata, H. Katoh, A. Kokubu, M. Sakamoto, K. Yamazaki, et al., Genome-wide array-based comparative genomic hybridization analysis of pancreatic adenocarcinoma: identification of genetic indicators that predict patient outcome, *Cancer Sci.* 98 (2007) 392–400, <http://dx.doi.org/10.1111/j.1349-7006.2007.00395.x>.
- [274] R.C. Castellino, M. De Bortoli, X. Lu, S.-H. Moon, T.-A. Nguyen, M.A. Shepard, et al., Medulloblastomas overexpress the p53-inactivating oncogene WIP1/PPM1D, *J. Neurooncol.* 86 (2008) 245–256, <http://dx.doi.org/10.1007/s11060-007-9470-8>.
- [275] X. Zhang, G. Wan, S. Mlotshwa, V. Vance, F.G. Berger, H. Chen, et al., Oncogenic Wip1 phosphatase is inhibited by miR-16 in the DNA damage signaling pathway, *Cancer Res.* 70 (2010) 7176–7186, <http://dx.doi.org/10.1158/0008-5472.CAN-10-0697>.
- [276] R.M. O'Connell, A.A. Chaudhuri, D.S. Rao, D. Baltimore, Inositol phosphatase SHIP1 is a primary target of miR-155, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 7113–7118, <http://dx.doi.org/10.1073/pnas.0902636106>.
- [277] I.M. Pedersen, D. Otero, E. Kao, A.V. Miletic, C. Hother, E. Ralfkiaer, et al., Onco-miR-155 targets SHIP1 to promote TNFalpha-dependent growth of B cell lymphomas, *EMBO Mol. Med.* 1 (2009) 288–295, <http://dx.doi.org/10.1002/emmm.200900028>.
- [278] D.W. Lee, M. Futami, M. Carroll, Y. Feng, Z. Wang, M. Fernandez, et al., Loss of SHIP-1 protein expression in high-risk myelodysplastic syndromes is associated with miR-210 and miR-155, *Oncogene* 31 (2012) 4085–4094, <http://dx.doi.org/10.1038/onc.2011.579>.
- [279] Z. Qu, J.N. Weiss, W.R. MacLellan, Regulation of the mammalian cell cycle: a model of the G1-to-S transition, *Am. J. Physiol. Cell Physiol.* 284 (2003) C349–C364, <http://dx.doi.org/10.1152/ajpcell.00066.2002>.

- [280] J. Chen, G. Dai, Y.-Q. Wang, S. Wang, F.-Y. Pan, B. Xue, et al., Polo-like kinase 1 regulates mitotic arrest after UV irradiation through dephosphorylation of p53 and inducing p53 degradation, *FEBS Lett.* 580 (2006) 3624–3630, <http://dx.doi.org/10.1016/j.febslet.2006.05.047>.
- [281] J. Wang, C.E. Kirby, R. Herbst, The tyrosine phosphatase PRL-1 localizes to the endoplasmic reticulum and the mitotic spindle and is required for normal mitosis, *J. Biol. Chem.* 277 (2002) 46659–46668, <http://dx.doi.org/10.1074/jbc.M206407200>.
- [282] M.K. Pathak, D. Dhawan, D.J. Lindner, E.C. Borden, C. Farver, T. Yi, Pentamidine is an inhibitor of PRL phosphatases with anticancer activity, *Mol. Cancer Ther.* 1 (2002) 1255–1264.
- [283] T. Kajino, H. Ren, S.-I. Iemura, T. Natsume, B. Stefansson, D.L. Brautigan, et al., Protein phosphatase 6 down-regulates TAK1 kinase activation in the IL-1 signaling pathway, *J. Biol. Chem.* 281 (2006) 39891–39896, <http://dx.doi.org/10.1074/jbc.M608155200>.
- [284] B. Stefansson, D.L. Brautigan, Protein phosphatase PP6 N terminal domain restricts G1 to S phase progression in human cancer cells, *Cell Cycle Georget. Tex.* 6 (2007) 1386–1392.
- [285] S. Nouaille, C. Blanquart, V. Zilberfarb, N. Boute, D. Perdureau, J. Roix, et al., Interaction with Grb14 results in site-specific regulation of tyrosine phosphorylation of the insulin receptor, *EMBO Rep.* 7 (2006) 512–518, <http://dx.doi.org/10.1038/sj.embor.7400668>.
- [286] K.R. LaMontagne Jr., A.J. Flint, B.R. Franza Jr., A.M. Pandergast, N.K. Tonks, Protein tyrosine phosphatase 1B antagonizes signalling by oncoprotein tyrosine kinase p210 bcr-abl in vivo, *Mol. Cell. Biol.* 18 (1998) 2965–2975.
- [287] J.D. Borge, A. Pang, D.J. Fujita, Identification of protein-tyrosine phosphatase 1B as the major tyrosine phosphatase activity capable of dephosphorylating and activating c-Src in several human breast cancer cell lines, *J. Biol. Chem.* 275 (2000) 41439–41446, <http://dx.doi.org/10.1074/jbc.M004852200>.
- [288] R.A. Klinghoffer, A. Kazlauskas, Identification of a putative Syp substrate, the PDGF beta receptor, *J. Biol. Chem.* 270 (1995) 22208–22217.
- [289] E.-J. Jung, G.A. Calin, The meaning of 21 in the MicroRNA world: perfection rather than destruction? *Cancer Cell* 18 (2010) 203–205, <http://dx.doi.org/10.1016/j.ccr.2010.08.015>.
- [290] O.A. Kent, J.T. Mendell, A small piece in the cancer puzzle: microRNAs as tumor suppressors and oncogenes, *Oncogene* 25 (2006) 6188–6196, <http://dx.doi.org/10.1038/sj.onc.1209913>.
- [291] C.A. Gebeshuber, K. Zatloukal, J. Martinez, miR-29a suppresses tristetraprolin, which is a regulator of epithelial polarity and metastasis, *EMBO Rep.* 10 (2009) 400–405, <http://dx.doi.org/10.1038/embor.2009.9>.
- [292] S.P. Nana-Sinkam, C.M. Croce, Clinical applications for microRNAs in cancer, *Clin. Pharmacol. Ther.* 93 (2013) 98–104, <http://dx.doi.org/10.1038/clpt.2012.192>.
- [293] N. Ahmad, S. Haider, S. Jagannathan, E. Anaissie, J.J. Driscoll, MicroRNA theragnostics for the clinical management of multiple myeloma, *Leukemia* 28 (2014) 732–738, <http://dx.doi.org/10.1038/leu.2013.262>.
- [294] M. Llauradó, B. Majem, T. Altadill, L. Lanau, J. Castellví, J.L. Sánchez-Iglesias, et al., MicroRNAs as prognostic markers in ovarian cancer, *Mol. Cell. Endocrinol.* 390 (2014) 73–84, <http://dx.doi.org/10.1016/j.mce.2014.03.006>.
- [295] R. Liu, X. Chen, Y. Du, W. Yao, L. Shen, C. Wang, et al., Serum microRNA expression profile as a biomarker in the diagnosis and prognosis of pancreatic cancer, *Clin. Chem.* 58 (2012) 610–618, <http://dx.doi.org/10.1373/clinchem.2011.172767>.
- [296] W. Gao, X. Lu, L. Liu, J. Xu, D. Feng, Y. Shu, MiRNA-21: a biomarker predictive for platinum-based adjuvant chemotherapy response in patients with non-small cell lung cancer, *Cancer Biol. Ther.* 13 (2012) 330–340, <http://dx.doi.org/10.4161/cbt.19073>.
- [297] G. Manceau, S. Imbeaud, R. Thiébaud, F. Liébaert, K. Fontaine, F. Rousseau, et al., Hsa-miR-31-3p expression is linked to progression-free survival in patients with KRAS wild-type metastatic colorectal cancer treated with anti-EGFR therapy, *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* 20 (2014) 3338–3347, <http://dx.doi.org/10.1158/1078-0432.CCR-13-2750>.
- [298] A.G. Bader, miR-34-a microRNA replacement therapy is headed to the clinic, *Front. Genet.* 3 (2012) 120, <http://dx.doi.org/10.3389/fgene.2012.00120>.