


Fish type I and type II interferons: composition, receptor usage, production and function

Zhen Gan^{1,2,*}, Shan Nan Chen^{1,*}, Bei Huang^{1,†}, Jun Zou³ and Pin Nie^{1,4,5} 

- 1 State Key Laboratory of Freshwater Ecology and Biotechnology, and Key Laboratory of Aquaculture Disease Control, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei Province, China
- 2 University of Chinese Academy of Sciences, Beijing, China
- 3 National Pathogen Collection Center of Aquatic Animals, College of Fisheries and Life Science, Shanghai Ocean University, Shanghai, China
- 4 Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, Shandong Province, China
- 5 School of Marine Science and Engineering, Qingdao Agricultural University, Qingdao, Shandong Province, China

Correspondence

Pin Nie, School of Marine Science and Engineering, Qingdao Agricultural University, Qingdao, Shandong Province, 266109, China; or Institute of Hydrobiology, Chinese Academy of Sciences, 7 South Donghu Road, Wuchang District, Wuhan, Hubei Province, 430072, China. Emails: pinnie@qau.edu.cn, pinnie@ihb.ac.cn

*Indicating the two-first authors.

†Present address: College of Fisheries, Jimei University, Xiamen, Fujian Province, 361021, China.

Received 13 December 2018; accepted 27 March 2019.

Abstract

Interferons (IFNs) have been reported in vertebrates from fish to mammals. To date, two types of IFNs, i.e. type I and type II IFNs have been reported in a range of fish species in aquaculture. In fish, type I IFNs are clarified into group I and group II, with two cysteines and four cysteines, and further into subgroups a, d, e, h, and b, c, f, respectively. Group I IFNs appear to exist in all fish species, whereas group II IFNs are discovered only in certain species such as cyprinids, salmonids and perciform fish. It seems preferentially functional that group I IFNs signal through cytokine receptor family B (CRFB) 5 and CRFB1, while group II IFNs through CRFB5 and CRFB2. But they all initiate a same signalling pathway for the expression of interferon induced genes. The information about the production and the function of type I IFNs in fish are further summarized in the review. Interestingly, two members of type II IFNs, IFN- γ and a unique member, IFN- γ related gene (IFN- γ rel) exist in fish, and IFN- γ and IFN- γ rel can separately become homodimers for signalling through CRFB13, CRFB17, CRFB6 and CRFB17, respectively. In addition, the variation in the composition of type I and type II IFNs as well as in their receptors in some cyprinid and salmonid fish has also been reviewed. It is suggested that future perspectives for the research on type I and type II IFNs in fish are outlined from a comparative and evolutionarily point of view.

Key words: antiviral response, fish, IFN receptor, Interferon, type I IFN, type II IFN.

Introduction

Interferons (IFNs) are a subset of the class II cytokines playing crucial roles in host immune defense, especially in immune response against viruses (Pestka *et al.* 2004). It is perhaps not an exaggeration to say that the discovery and clinical use of IFNs have been a major advance in immunology and biomedicine over the past sixty years. Based on sequence similarity, genomic organisations, receptor usage and biological function, IFNs are classified into three types, i.e. type I, type II and type III IFNs (Pestka *et al.* 2004). Type I and type III IFNs are specialized as innate antiviral cytokines (Lazear *et al.* 2015; Teijaro 2016), and their main difference lies in their distribution and

target cells: almost all nucleated cells respond to type I IFNs, whereas type III IFNs are restricted in tissues with high risk of viral exposure and infection, such as those with epithelial surfaces (Lazear *et al.* 2015; Teijaro 2016). By contrast, type II IFN is now known to represent a regulatory cytokine orchestrating both innate and adaptive immunity against viral and intracellular bacterial infections (Schoenborn & Wilson 2007).

From an evolutionary point of view, IFNs have been identified in vertebrates from fish to mammals (Zou & Secombes 2011; Gan *et al.* 2017, 2018; Langevin *et al.* 2013a; Santhakumar *et al.* 2017; Secombes & Zou 2017). Fish can be classified into three classes, i.e. Agnatha (jawless fish), Chondrichthyes (cartilaginous fish) and Osteichthyes

(bony fish), and the latter two are considered as the earliest vertebrates possessing the IFN system, representing an important model to expand the knowledge about the origin and evolution of the IFN system in vertebrates (Zou & Secombes 2011; Zhang & Gui 2012; Langevin *et al.* 2013a; Boudinot *et al.* 2016; Secombes & Zou 2017; Robertsen 2018). On the other hand, as a vital protein source for human consumption, farmed fish are now threatened with viral infectious diseases, and the understanding of IFN system in fish with aquaculture importance may provide clues for the development of strategies in relation with the prevention of diseases (Workenhe *et al.* 2010; Evensen & Leong 2013; Collet 2014; Collins *et al.* 2018; Oidtmann *et al.* 2018). Hence, the fish IFN system has been one of the research focuses for comparative immunologists, and significant progress has been achieved in understanding the fish IFN system in recent years. Although type III IFN genes are not to date identified in fish (Zou & Secombes 2011; Zhang & Gui 2012; Langevin *et al.* 2013a; Boudinot *et al.* 2016; Secombes & Zou 2017; Robertsen 2018), IFN ligands, receptors, signalling pathways, regulatory mechanism and functional properties of type I and II IFN systems have been extensively studied. This review attempts to summarize the recent discoveries on the IFN systems in fish.

Type I IFNs in fish

Discovery and phylogeny of type I IFNs

Although IFN was first discovered by Alick Isaacs and Jean Lindenmann as early as in 1957, the first fish IFN gene was identified in 2003 by three independent groups in zebrafish (*Danio rerio*) (Altmann *et al.* 2003), Atlantic salmon (*Salmo salar*) (Robertsen *et al.* 2003) and green spotted pufferfish (*Tetraodon nigroviridis*) (Lutfalla *et al.* 2003). Afterwards, the IFN genes have been identified in a multitude of fish species, including grass carp (*Ctenopharyngodon idella*), common carp (*Cyprinus carpio*), gibel carp (*Carassius auratus gibelio*), red crucian carp (*C. auratus* red var.), black carp (*Mylopharyngodon piceus*), rohu (*Labeo rohita*), rainbow trout (*Oncorhynchus mykiss*), mandarin fish (*Siniperca chuatsi*), large yellow croaker (*Larimichthys crocea*), orange-spotted grouper (*Epinephelus coioides*), seven band grouper (*E. septemfasciatus*), European sea bass (*Dicentrarchus labrax*), gilthead seabream (*Sparus aurata*), rock bream (*Oplegnathus fasciatus*), meagre (*Argyrosomus regius*), turbot (*Scophthalmus maximus*), Japanese flounder (*Paralichthys olivaceus*), channel catfish (*Ictalurus punctatus*), medaka (*Oryzias latipes*) and Japanese eel (*Anguilla japonica*) (Table 1) (Gonzalez *et al.* 2007; Aggad *et al.* 2009; Casani *et al.* 2009; Chang *et al.* 2009; Chen *et al.* 2014; Huang *et al.* 2015b; Ding *et al.* 2016, 2017; Hu *et al.* 2017; Huang *et al.* 2018a; Long *et al.* 2004, 2006; Zou *et al.* 2007, 2014a; Kitao *et al.* 2009; Lopez-Munoz *et al.* 2009; Purcell *et al.* 2009;

Sun *et al.* 2009; Li *et al.* 2012; Wan *et al.* 2012; Svingerud *et al.* 2012; Parhi *et al.* 2014; Pereiro *et al.* 2014; Valero *et al.* 2015; Kuo *et al.* 2016; Liao *et al.* 2016; Maekawa *et al.* 2016; Wei *et al.* 2016; Yan *et al.* 2016; Yang *et al.* 2017; Laghari *et al.* 2018; Milne *et al.* 2018; Wu *et al.* 2018; Xia *et al.* 2018). In addition, the IFN genes are also found in cartilaginous fish, such as elephant shark (*Callorhynchus milii*) (Chang *et al.* 2009; Zou & Secombes 2011).

Nevertheless, the phylogenetic relationship of these fish IFNs with mammalian type I and III IFNs had been heavily debated among comparative immunologists in the last ten to twenty years (Lutfalla *et al.* 2003; Robertsen 2006; Levraud *et al.* 2007; Zou *et al.* 2007; Aggad *et al.* 2009; Chang *et al.* 2009). Several aspects of evidence based on sequence analysis suggested that fish IFNs are likely to be the homologues of mammalian type I IFNs. In particular, they possess two important sequence features which are conserved between fish and mammalian type I IFNs, including the cysteine pattern and the C-terminal CAWE motif (Robertsen 2006; Zou *et al.* 2007; Chang *et al.* 2009). In contrast, it was argued that fish IFNs were likely to be the homologues of mammalian type III IFNs, as these fish IFNs are encoded by intron-containing genes with five exons and four introns as seen in mammalian type III IFNs, being entirely different from mammalian type I IFN genes which are intronless (Lutfalla *et al.* 2003). Consistently, the receptors of these fish IFNs share features of mammalian type III IFN receptors rather than type I IFN receptors in terms of the domain organization (Lutfalla *et al.* 2003; Aggad *et al.* 2009; Levraud *et al.* 2007). The short cytoplasmic receptor chain of fish IFNs, named as cytokine receptor family B (CRFB) 5, only possesses two fibronectin type III (FNIII) domains, as observed in the corresponding receptor chain of mammalian type III IFNs (IL-10R2), being distinct from the one of mammalian type I IFNs (IFNAR1) which has a uniquely large extracellular region containing four FNIII domains (Lutfalla *et al.* 2003; Aggad *et al.* 2009; Levraud *et al.* 2007). It was once hard to claim with certainty that these fish IFNs were the orthologues of mammalian type I or type III IFNs before the possible emergence of new evidence, so some researchers proposed that fish IFNs should be named as IFN ϕ , representing a unique type of virus-induced IFNs in fish (Stein *et al.* 2007).

This puzzling controversy has been clarified in a crystallographic study to determinate the three-dimensional structure of zebrafish IFNs in 2011, in which both zebrafish IFN1 and IFN2 exhibit typical type I IFN structure with a straight F helix rather than a bend in type III IFN architecture (Hamming *et al.* 2011). Now, it is generally accepted that these fish IFNs belong to type I IFNs (Zou & Secombes 2011; Zhang & Gui 2012; Langevin *et al.* 2013a; Boudinot *et al.* 2016; Secombes & Zou 2017; Robertsen 2018), but two new intriguing questions need to be further resolved:

Table 1 Currently reported type I IFNs in fish

Order	Species	Gene	GenBank accession number	References
Cypriniformes	Zebrafish (<i>D. rerio</i>)	Dr-IFN1	NM_207640	(Aggad <i>et al.</i> 2009; Altmann <i>et al.</i> 2003; Lopez-Munoz <i>et al.</i> 2009)
		Dr-IFN2	NM_0011111082	
		Dr-IFN3	NM_0011111083	
		Dr-IFN4	NM_001161740	
	Grass carp (<i>C. idella</i>)	Ci-IFN1	DQ357216	(Li <i>et al.</i> 2012; Liao <i>et al.</i> 2016)
		Ci-IFN2-4	KU182641–KU182643	
	Common carp (<i>C. carpio</i>)	Cc-IFN1-2	AB376666–AB376667	(Gonzalez <i>et al.</i> 2007; Kitao <i>et al.</i> 2009; Wei <i>et al.</i> 2016)
	Crucian carp (<i>C. auratus</i>)	Ca-IFNa	AY452069	(Xia <i>et al.</i> 2018)
		Ca-IFNc	MF574165	
	Red crucian carp (<i>C. auratus</i> red var.)	Car-IFNa	KU561831	(Yan <i>et al.</i> 2017; Yan <i>et al.</i> 2016)
	Black carp (<i>M. piceus</i>)	Car-IFNa2	KX839490	(Huang <i>et al.</i> 2015b; Wu <i>et al.</i> 2018)
		Mp-IFNa	KR265208	
Salmoniformes	Rohu (<i>L. rohita</i>)	Mp-IFNb	MF497811	(Parhi <i>et al.</i> 2014)
		Lr-IFN	HQ667143	
		Om-IFNa1 (IFN1)	CAM28541	
	Rainbow trout (<i>O. mykiss</i>)	Om-IFNa2 (IFN2)	NP_001153977	(Chang <i>et al.</i> 2009; Purcell <i>et al.</i> 2009; Zou <i>et al.</i> 2007, 2014a)
		Om-IFNa3–a4	HF931021–HF931022	
		Om-IFNb1 (IFN3)	NP_001153974	
		Om-IFNb2 (IFN4)	NP_001158515	
		Om-IFNb3–b5	HF931023–HF931025	
		Om-IFNc1–c4	HF931026–HF931029	
		Om-IFNd1 (IFN5)	NP_001152811	
		Om-IFNe1–e7	HF931030–HF931036	
	Atlantic salmon (<i>S. salar</i>)	Om-IFNf1–f2	HF931037–HF931038	(Robertsen <i>et al.</i> 2003; Sun <i>et al.</i> 2009; Svingerud <i>et al.</i> 2012)
		Ss-IFNa	AY216594	
		Ss-IFNb	JX524152	
		Ss-IFNc	JX524153	
		Ss-IFNd	JX524151	
Perciformes	Mandarin fish (<i>S. chuatsi</i>)	Sc-IFNc	KY768914	(Laghari <i>et al.</i> 2018)
		Sc-IFNd	KY768915	
		Sc-IFNh	KY768916	
	Large yellow croaker (<i>L. crocea</i>)	Lc-IFNd	KU144880	(Ding <i>et al.</i> 2016, 2017)
	Orange-spotted grouper (<i>E. coioides</i>)	Lc-IFNh	KU144879	(Chen <i>et al.</i> 2014)
		Ec-IFN	JX443623	
	Seven band grouper (<i>E. septemfasciatus</i>)	Es-IFN	AB585968	(Kuo <i>et al.</i> 2016)
Pleuronectiformes	European sea bass (<i>D. labrax</i>)	Sb-IFN1–2	AM765846–AM765847	(Casani <i>et al.</i> 2009; Valero <i>et al.</i> 2015)
		Sa-IFN1–4	FM882241–FM882244	
	Gilthead seabream (<i>S. aurata</i>)	Rb-IFN	JX020703	(Wan <i>et al.</i> 2012)
		Ar-IFNc	MG489872	
	Rock bream (<i>O. fasciatus</i>)	Ar-IFNd	MG489873	(Milne <i>et al.</i> 2018)
		Ar-IFNh	MG489874	
	Meagre (<i>A. regius</i>)			
Pleuronectiformes	Turbot (<i>S. maximus</i>)	Sm-IFN1–2	KJ150677–KJ150678	(Pereiro <i>et al.</i> 2014)
		Jf-IFN1	BAH84776	
	Japanese flounder (<i>P. olivaceus</i>)	Jf-IFN2	AHB59752	(Hu <i>et al.</i> 2017)
		Jf-IFN3–4	LC222627–LC222628	

Table 1 (continued)

Order	Species	Gene	GenBank accession number	References
Tetraodontiformes	Pufferfish (<i>T. nigroviridis</i>)	Tn-IFN	AJ544889	(Lutfalla <i>et al.</i> 2003)
Siluriformes	Channel catfish (<i>I. punctatus</i>)	Ip-IFN1	AY267538	(Long <i>et al.</i> 2004, 2006)
		Ip-IFN2	AY847295	
		IpIFN4	AY847296	
Beloniformes	Medaka (<i>O. latipes</i>)	OI-IFNa	LC066594	(Maekawa <i>et al.</i> 2016)
		OI-IFNd	LC066595	
Anguilliformes	Japanese eel (<i>A. japonica</i>)	Aj-IFN1–4	MH822129–MH822132	(Huang <i>et al.</i> 2018a)

how intron-containing type I IFN genes in fish might have evolved into intronless type I IFN genes in mammals, and CRFB5 with two FNIII domains in fish evolved into IFNAR1 with four FNIII domains in mammals.

Recently, studies on amphibian type I IFNs and their receptors provide reasonable evidence to explain the difference in genomic organisations between fish and mammalian type I IFNs as well as in the copy number of FNIII domains between the receptors (Gan *et al.* 2017, 2018). It is revealed that intronless and intron-containing type I IFN genes coexist in the amphibian species, the clawed frog *Xenopus* and the Tibetan frog *Nanorana parkeri*, providing key evidence to support the hypothesis that retroposition events may have resulted in the generation of intronless type I IFNs in vertebrates (Gan *et al.* 2017, 2018). More importantly, several lines of evidence based on homology, synteny, phylogeny and divergence time estimation indicate that the intronless type I IFN genes in *Xenopus*, *N. parkeri* and amniotes might have arisen from three independent retroposition events occurred in these three lineages (Gan *et al.* 2017, 2018). It can then be proposed that the intron-containing type I IFN genes in fish might have lost their introns and evolved into the intronless type I IFN genes in amniotes through the retroposition event (Gan *et al.* 2017, 2018). On the other hand, it is interesting to note that the extracellular region of IFNAR1 in *N. parkeri* contains only two FNIII domains, which is similar to CRFB5 in fish, suggesting that the two FNIII domain-containing CRFB5 gene in fish might have gained other two FNIII domains and evolved into the four FNIII domain-containing IFNAR1 gene in mammals via a tandem domain duplication event during a transit period from cold-blooded to warm-blooded tetrapods (Gan *et al.* 2018). As a result of these events during the evolution of tetrapods, fish type I IFNs harbour a combination of features observed in mammalian type I and type III IFNs.

Classification and nomenclature of type I IFNs

As described above, owing to the profound impacts of retroposition events in the evolution of type I IFNs, as well

as to the lineage-specific expansion and divergence of type I IFNs in vertebrates, it is difficult to establish orthologous relationships between fish type I IFNs with their mammalian counterparts. It is clear that fish type I IFNs cannot be simply classified into IFN α or IFN β or any other subtypes of type I IFNs according to the classification and nomenclature in mammals, and great efforts have been made already by fish immunologists to establish a new classification system for the type I IFNs in fish.

Like other vertebrates, fish also possess multiple copies of type I IFN genes which are linked in the genome, and the gene copy number varies in different species. For example, zebrafish appears to possess four type I IFN genes (Altmann *et al.* 2003; Aggad *et al.* 2009; Lopez-Munoz *et al.* 2009), whereas at least eleven type I IFN genes have been identified in Atlantic salmon (Robertsen *et al.* 2003; Sun *et al.* 2009; Svingerud *et al.* 2012). Based on the number of cysteine residues predicted to be involved in the formation of disulfide bond, fish type I IFNs were originally classified into two main groups, two cysteine-containing group I and four cysteine-containing group II (Zou *et al.* 2007; Chang *et al.* 2009). Group I IFNs appear to exist in all fish species, whereas group II IFNs are discovered only in certain species such as cyprinids, salmonids and perciform fish (Zou & Secombes 2011; Langevin *et al.* 2013a; Boudinot *et al.* 2016; Secombes & Zou 2017; Laghari *et al.* 2018; Robertsen 2018). Undoubtedly, it is rational to classify fish type I IFNs according to this criterion, because group I and group II IFNs have been proven to exhibit certain degrees of difference in their receptor usage, expression pattern, and antiviral activity, which will be discussed in the following sections. However, the copies of type I IFN genes in this classification system are distinguished by Arabic numerals according to their discovery order and chromosome location. In the case of zebrafish, four type I IFN genes are named as IFN1, IFN2, IFN3 and IFN4, with IFN1 and IFN4 being in group I, and IFN2 and IFN3 in group II (Altmann *et al.* 2003; Aggad *et al.* 2009; Lopez-Munoz *et al.* 2009). Obviously, this nomenclature does not distinguish group I nor group II IFNs, and may lead to the inconsistent classification among type I IFN genes from different species. For

instance, IFN1 or IFN2 from other species may fall in different groups as observed in zebrafish, and both IFN1 and IFN2 from the Japanese eel all belong to group I IFNs (Huang *et al.* 2018a).

To avoid the above-mentioned problem, a more complex classification system has been established, in which group I IFNs are further divided into subgroups a, d, e and h, and group II IFNs into subgroups b, c and f (Sun *et al.* 2009; Svingerud *et al.* 2012; Zou *et al.* 2014a; Ding *et al.* 2016; Laghari *et al.* 2018; Milne *et al.* 2018). Among the investigated fish species, salmonids appear to harbour the most complex type I IFN repertoire, containing subgroups a, b, c, d, e and f (Sun *et al.* 2009; Svingerud *et al.* 2012; Zou *et al.* 2014a). In addition, a recently identified IFNh, is only reported in perciformes (Ding *et al.* 2016; Laghari *et al.* 2018; Milne *et al.* 2018). According to the current sequence information, the type I IFN repertoire in most fish species appears to contain only three subgroups, including subgroups a, c and d (Zou & Secombes 2011; Langevin *et al.* 2013a; Boudinot *et al.* 2016; Secombes & Zou 2017; Robertsen 2018). Although it is convenient to recognize the subgroup of IFN genes in this classification system, this nomenclature may be easily confused with mammalian type I IFNs, as IFN α , IFN β etc. are present in mammals. The names of fish type I IFNs should be clarified to the subgroup and any possible confusion should be avoided with mammalian denomination, and a better nomenclature still remains to be established.

Transcript variants and intracellular forms of type I IFNs

In mammals, the IFN proteins consist of a predicted signal peptide and a mature region, which are secreted into extracellular space to perform their functions via a classic protein secretion pathway (Pestka *et al.* 2004). Unexpectedly, type I IFN variants which lack a signal peptide have been reported in several fish species (Long *et al.* 2006; Levraud *et al.* 2007; Purcell *et al.* 2009; Chang *et al.* 2013). These variants are transcribed from a single gene that generates two kinds of variants, referred to as the shorter and longer transcripts, with the shorter transcript encoding the normal form of IFN with a signal peptide, and the longer transcript encoding the intracellular form of IFN without a signal peptide (Levraud *et al.* 2007; Purcell *et al.* 2009; Chang *et al.* 2013). In zebrafish, the shorter transcript of IFN1 was induced by viral infection, whereas the longer transcript was constitutively expressed at a relatively high level and was poorly inducible after viral infection (Levraud *et al.* 2007). By contrast, both shorter and longer transcripts of IFN1 in rainbow trout could be detected in most tissues and were induced following viral infection, and it is proven that these two forms were derived from alternative

transcription under the control of two different promoters (Purcell *et al.* 2009).

Remarkably, the intracellular form of type I IFNs encoded by the longer transcript has been proven to be functional in rainbow trout, and its recombinant intracellular IFN can induce the expression of Mx and provide protection against viral infection *in vitro* and *in vivo* (Chang *et al.* 2013; Cao *et al.* 2016). Similar to the IFN genes, the IFN receptor genes in rainbow trout also have transcript variants translating into the intracellular form of IFN receptors without a signal peptide. The intracellular receptors seem to be mainly found in the perinuclear region and co-localized with the intracellular IFN. The overexpression of intracellular IFNs and receptors in HEK293 cells can lead to the phosphorylation of signal transducers and activators of transcription (STAT) 1 and STAT2 (Chang *et al.* 2013), confirming that intracellular IFN and receptors can indeed form a cellular ligand–receptor complex to induce the expression of interferon induced genes (ISGs) and a cellular antiviral state in a STAT-dependent manner. However, it remains to be determined whether intracellular IFNs are widely present in different fish species. Notably, all the fish intracellular IFNs stated above belong to the IFNa subgroup of group I IFNs. It is not clear whether fish type I IFN genes from other subgroups can also generate transcript variants encoding the intracellular IFN proteins.

Type I IFN receptors

In mammals, the initiation of type I IFN-mediated signalling pathway depends on the interaction between type I IFNs and their heterodimeric receptor complex comprised of IFNAR1 and IFNAR2 (Pestka *et al.* 2004). IFNAR1 is regarded as the low-affinity receptor chain with a uniquely large extracellular region containing four FNIII domains and a short cytoplasmic region, whereas the high-affinity receptor chain IFNAR2 has a two FNIII domain-containing extracellular region and a long cytoplasmic region (Piehler *et al.* 2012). Both IFNAR1 and IFNAR2 belong to the class II cytokine receptor family, which is called the cytokine receptor family B (CRFB) in fish by comparative immunologists (Lutfalla *et al.* 2003; Stein *et al.* 2007). Genome-wide sequence analyses revealed that at least 17 CRFB members are present in zebrafish and pufferfish (Lutfalla *et al.* 2003; Stein *et al.* 2007; Aggad *et al.* 2010). Phylogenetically, it is indicated that CRFB1 and CRFB2 can be the homologues of mammalian IFNAR2, and CRFB5 is of the mammalian IFNAR1 (Stein *et al.* 2007). The two group I IFNs in zebrafish, which are named as IFN1/4 in zebrafish and should be IFNa and IFNd, respectively, were found to signal through a receptor complex consisted of CRFB1 and CRFB5, while the two group II IFNs, i.e. IFN2/3 which should all be

classified as IFNc, signal through a distinct receptor composed of CRFB2 and CRFB5 (Levraud *et al.* 2007; Aggad *et al.* 2009).

The evidence obtained in other species of fish is almost consistent with the observation in zebrafish. As revealed by cross-linking reaction, recombinant grass carp IFNa (belonging to group I IFNs) can bind the recombinant extracellular region of CRFB1 and CRFB5 to form a ternary complex *in vitro* (Chen *et al.* 2015a,b). In mandarin fish, the co-transfection assays showed that IFNd and IFNh (group I IFNs) use preferentially the receptor complex comprising of CRFB1 and CRFB5, and IFNc (group II IFN) use preferentially the one consisting of CRFB2 and CRFB5, for their efficacious induction of ISGs (Laghari *et al.* 2018). Curiously, the co-transfection of mandarin fish IFNh, CRFB2 and CRFB5 expressing plasmids also had inductive effect on the expression of ISGs (Laghari *et al.* 2018). In consistency with the complex type I IFN repertoire, Atlantic salmon appears to have a more extended repertoire of type I IFN receptors than other fish species, with two CRFB clusters containing two CRFB1 members (CRFB1a and CRFB1b), one CRFB2 gene, and four CRFB5 members (CRFB5a, CRFB5b, CRFB5c and CRFB5x) (Sun *et al.* 2014). The Mx reporter assays suggested that salmon IFNa (group I IFN) signals through a receptor complex consisted of CRFB1a as the long cytoplasmic chain and either CRFB5a, or CRFB5b or CRFB5c as the short cytoplasmic chain. By contrast, IFNc (group II IFN) signals through the one composed of CRFB2 as the long chain and either CRFB5a or CRFB5c as the short chain, and that IFNb (group II IFN) through the one comprised of CRFB2 as the long chain and CRFB5x as the short chain (Sun *et al.* 2014).

It is therefore demonstrated that fish group I and group II IFNs may signal through distinct receptor complexes (Fig. 1), which is obviously different from the manner in mammals, since all mammalian type I IFN subtypes initiate signalling via the same receptor complex consisting of IFNAR1 and IFNAR2.

Type I IFN-mediated signalling pathway

Mammalian type I IFNs exert their effects mainly through the JAK-STAT pathway (Stark & Darnell 2012). The interaction between type I IFNs and their receptors triggers recruitment and binding of tyrosine kinase 2 (TYK2) and Janus kinase 1 (JAK1) to IFNAR1 and IFNAR2, respectively. Subsequently, these activated kinases promote the phosphorylation of STAT1 and STAT2, which are dimerized and assembled with IFN regulatory factor (IRF) 9 to form a complex called IFN-stimulated gene factor 3 (ISGF3). This trimolecular complex is translocated to the nucleus and binds to IFN-stimulated response elements (ISREs), thus activating the




Figure 1 Signalling and ligand-receptor models for type I IFNs in fish. The two groups of type I IFNs, two cysteine-containing group I and four cysteines-containing group II IFNs, bind to receptor complexes with a common short cytoplasmic receptor chain CRFB5, but two distinct long cytoplasmic receptor chains, CRFB1 and CRFB2. The interaction between type I IFNs and their receptors triggers the recruitment and activation of TYK2 and JAK1. Subsequently, these activated kinases promote the phosphorylation of STAT1 and STAT2, which are dimerized and assembled with IRF9 to form a complex called ISGF3. This trimolecular complex translocates to the nucleus and binds to ISREs, thus activating the transcription of ISGs.

transcription of several hundred ISGs (Platanias 2005). As revealed in several immunogenetics studies, the essential components involved in this pathway, including TYK2, JAK1, STAT1, STAT2 and IRF9, also exist in fish (Guo *et al.* 2009; Huang *et al.* 2010; Liongue *et al.* 2012; Jin *et al.* 2018; Xu *et al.* 2016). Mammalian IFNAR1 and IFNAR2 have no intrinsic enzymatic activity, and instead depend on the kinase activity of TYK2 and JAK1 for signal initialization (Piehler *et al.* 2012). Two recent studies suggested that this mechanism is likely conserved in fish. It is shown that the overexpression of Atlantic salmon TYK2 led to the autophosphorylation by a kinase domain-dependent way as seen in mammals, leading to the up-regulation of Mx expression (Sobhkhez *et al.* 2013). In grass carp, both JAK1 and TYK2 were up-regulated and phosphorylated following the stimulation by polyinosinic-polycytidylic acid [poly(I:C)], and the

overexpression of type I IFN receptor chains CRFB1 and CRFB5 also facilitated the phosphorylation of JAK1 and TYK2 (Hou *et al.* 2017).

Intriguingly, two STAT1 genes, named as STAT1a and STAT1b, were identified in zebrafish (Song *et al.* 2011). Zebrafish STAT1a contains all five domains of human STAT1a, whereas zebrafish STAT1b lacks the C-terminal transcriptional activation domain, resembling human STAT1b, a splice variant of human STAT1a (Song *et al.* 2011). Several reports from different fish species showed that fish STAT1a may play a vital role in type I IFN-mediated signalling (Oates *et al.* 1999; Skjesol *et al.* 2010; Zhang *et al.* 2017). Zebrafish STAT1a is able to rescue IFN-mediated growth suppression in a STAT1-deficient human cell line (Oates *et al.* 1999). Consistent with this finding, Atlantic salmon STAT1a was phosphorylated and transported rapidly into the nucleus following the treatment with recombinant IFN α 1 (Skjesol *et al.* 2010), and the overexpression of orange-spotted grouper STAT1a significantly up-regulated the expression level of ISGs, thereby exerting antiviral activity against iridovirus and nodavirus (Zhang *et al.* 2017). However, gibel carp STAT1, which is similar to zebrafish STAT1b with the lack of the C-terminal transcriptional activation domain, has also been shown to be involved in the IFN α -triggered ISG induction as well as in the inhibition of viral infection (Yu *et al.* 2010). Similar to the highly extended repertoire of type I IFNs and their receptors, STAT1 and STAT2 genes exist as multiple copies in Atlantic salmon, including STAT1a, STAT1b and STAT1c, and STAT2a, STAT2b and STAT2c (Sobhkhez *et al.* 2014). Both Atlantic salmon STAT2a and STAT2b were phosphorylated and co-localized with STAT1a in the nucleus following the stimulation of recombinant IFN α 1 (Sobhkhez *et al.* 2014). As revealed by Co-IP assay, both salmon STAT2a and STAT2b can bind to STAT1a and IRF9, suggesting that the ISGF3 complex composed of STAT1, STAT2 and IRF9 may conservatively exist in fish (Sobhkhez *et al.* 2014). This notion is further confirmed in a recent study in which mandarin fish STAT1, STAT2 and IRF9 can interact with each other *in vitro* (Laghari *et al.* 2018). Consistently, it has been demonstrated in studies from different fish species that fish IRF9 is essential for the type I IFN-mediated signalling (Shi *et al.* 2012; Huang *et al.* 2017a; Wu *et al.* 2017).

It is thus suggested that fish type I IFNs may mediate signalling downstream of the receptors in a similar way as in mammals (Fig. 1). Nevertheless, it remains to be determined whether multicopy STAT1 and STAT2 genes exist in a broad range of taxa in fish. Furthermore, it will be interesting to investigate the functional similarity and divergence of fish multicopy STAT1 and STAT2 genes in type I IFN signalling.

Regulation of type I IFN response

Extensive studies have shown that at least three families of pattern recognition receptors (PRRs), including the retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs), Toll-like receptors (TLRs), and cytosolic DNA sensors, are involved in the activation of type I IFN response in mammals (Kawai & Akira 2011; Loo & Gale 2011; Wu & Chen 2014). When these PRRs are engaged with their corresponding ligands, they recruit downstream signalling molecules, resulting in the activation of key transcription factors, such as IRFs, for initiating eventually the transcription of type I IFN genes (Kawai & Akira 2011; Loo & Gale 2011; Wu & Chen 2014). The PRR-induced mechanism of type I IFN production seems to be conserved in fish (Aoki *et al.* 2013; Poynter *et al.* 2015) (Fig. 2), which will be discussed in this section.

RLR-mediated type I IFN response

Among all known PRRs, RLRs are a family of cytosolic receptors recognizing viral RNAs with three members, including RIG-I, melanoma differentiation-associated gene 5 (MDA5) and laboratory of genetics and physiology 2 (LGP2) (Loo & Gale 2011). In mammals, RIG-I senses primarily 5' triphosphated double-stranded RNA (dsRNA), whereas MDA5 recognizes long dsRNA (Bruns & Horvath 2014). After the detection of appropriate viral RNAs, RIG-I and MDA5 can interact with the adaptor, mitochondrial antiviral signalling protein (MAVS), and recruit cytosolic protein kinase TANK binding kinase 1 (TBK1) through their caspase activation and recruitment domains (CARDs), leading to the activation of IRF3 and IRF7 and the transcription of type I IFN genes (Loo & Gale 2011). In contrast, LGP2 differs from RIG-I and MDA5 in lacking CARDs, but retains the ability to bind to dsRNA and the function to regulate signalling by RIG-I and MDA5 (Rothenfusser *et al.* 2005; Satoh *et al.* 2010). All three members in the RLR family, RIG-I, MDA5 and LGP2, as well as the key downstream molecules MAVS and TBK1, are found conservatively in fish (Zou *et al.* 2009; Chen *et al.* 2017b). Although MDA5 and LGP2 appear to exist in all fish species, RIG-I seems to be lost in fish of Acanthopterygii (Zou *et al.* 2009; Chen *et al.* 2017b). With regard to the ligand recognition by RLRs in fish, only a report from rainbow trout proved that MDA5 and LGP2 can bind synthetic dsRNA poly(I:C) by pull down assay *in vitro* (Chang *et al.* 2011). To date, there has been no experimental evidence for the ligand-binding activity of fish RIG-I.

Functionally, like its mammalian counterpart, fish RIG-I possesses the ability to induce the expression of type I IFNs, ISGs, and antiviral state, as supported by studies from




Figure 2 Regulation of type I IFN response in fish. Upon the recognition of dsRNA from viruses, RIG-I/MDA5 recruits MAVS, an adaptor protein located on mitochondria, which then become associated with TRAF3 and TBK1, leading to the phosphorylation and activation of IRF3/IRF7. LGP2 can be a positive or negative regulator of RIG-I/MDA5-mediated type I IFN response. TLR3, TLR19 and TLR22 can also recognize dsRNA from viruses and recruit TRIF and TBK1 to elicit type I IFN response. In addition, STING recruits TBK1 and IRF3 to trigger the expression of type I IFNs. IRF1, IRF3 and IRF7 can become either homo- or hetero-dimerized to recruit MyD88, for translocating into nucleus and then for binding ISRE motif, and for the production of type I IFNs and ISGs. Both IRF2 and IRF10 can inhibit type I IFN production by binding ISRE motif, and IRF10 can also inhibit type I IFN response through interacting with STING. FOXO3 can interact directly with IRF3 and IRF7, thereby suppressing IRF3/7-mediated type I IFN response. PIAS4a, TBK1L, TBK1_tv1, TBK1_tv2 and STAT6 negatively regulate the RLRs-MAVS-TBK1-mediated signalling pathway. In the signalling schematics, the negative regulation cascades are marked with red lines, and broken lines indicate the direct interaction or the molecules whose interaction need to be confirmed.

several fish species (Biacchesi *et al.* 2009, 2012; Sun *et al.* 2011a,b; Li *et al.* 2014a; Wang *et al.* 2014; Chen *et al.* 2015d; Zou *et al.* 2015; Wan *et al.* 2017). The transfection in EPC cells with N-terminal region of EPC RIG-I can activate strongly the EPC type I IFN promoter, and induce the expression of type I IFN and ISGs, thus resisting viral haemorrhagic septicaemia virus (VHSV) infection (Biacchesi *et al.* 2009). Similarly, the overexpression of gibel carp RIG-I enhanced significantly the gibel carp and zebrafish type I IFN promoters as well as the expression of viperin in CAB cells (Sun *et al.* 2011b; Wang *et al.* 2014), and the grass carp RIG-I when overexpressed in CIK cells increased significantly the expression of IFN2 and IFN4 (Wan *et al.*

2017). Research from zebrafish showed that the N-terminal region or wild-type of zebrafish RIG-I when overexpressed in EPC cells strongly activated zebrafish type I IFN promoter (Biacchesi *et al.* 2012; Li *et al.* 2014a; Zou *et al.* 2015). In addition, EPC cells transfected with wild-type RIG-I in zebrafish are resistant to spring viraemia of carp virus (SVCV) infection, with the increasing expression of Mx and IRF7, and these effects can be further improved by the co-transfection of zebrafish RIG-Ia, a longer spliced variant of wild-type RIG-I (Zou *et al.* 2015). Likewise, zebrafish IFN1, IFN4 and MxC genes were significantly induced when wild-type RIG-I was overexpressed in embryos (Chen *et al.* 2015d).

Like RIG-I, MDA5 in fish has also been proven to be capable of triggering type I IFN-mediated antiviral response, which was reported initially in rainbow trout and Japanese flounder, in which the overexpression of MDA5 in fish cells caused the increase in the expression of type I IFNs and ISGs as well as antiviral activity against viral infection (Chang *et al.* 2011; Ohtani *et al.* 2011). Similar results were reported in recent studies from other fish species. MDA5 in orange-spotted grouper when overexpressed in GS cells enhanced significantly type I IFN promoter activities as well as the expression of IRF3 and IRF7, thereby inhibiting cytopathic effect (CPE) progression and viral gene transcription during red-spotted grouper nervous necrosis virus (RGNNV) and Singapore grouper iridovirus (SGIV) infection (Huang *et al.* 2016a). Ectopic expression of black carp MDA5 in EPC cells resulted in the induction of promoter activity of zebrafish and fathead minnow type I IFNs, and improved antiviral ability against SVCV and grass carp reovirus (GCRV) infection (Liu *et al.* 2017). The overexpression of grass carp MDA5 in CIK cells significantly enhanced the expression of type I IFNs (Wan *et al.* 2017). Consistently, the knock down of MDA5 from sea perch (*Lateolabrax japonicus*) significantly increased the viral burden in RGNNV infected LJF cells (Jia *et al.* 2016a). More convincingly, several recent studies have shown that zebrafish MDA5 can enhance type I IFN-mediated antiviral response *in vitro* and *in vivo*. The activation of type I IFN promoters and the resistance against viral infection were found in fish cells transfected with zebrafish wild-type MDA5 (Li *et al.* 2014a; Zou *et al.* 2014b; Gabor *et al.* 2015), and such effect can be enhanced by the co-transfection of zebrafish MDA5b, a shorter spliced variant of wild-type MDA5 (Zou *et al.* 2014b). It is further confirmed that zebrafish MDA5 when overexpressed in embryos increased the expression of type I IFNs and ISGs (Chen *et al.* 2015d), and dominant negative MDA5 gene transgenic zebrafish displayed higher viral burden and mortality after snakehead vesiculovirus (SHVV) infection, which could be rescued by recombinant full-length MDA5 (Gabor *et al.* 2015).

Unlike RIG-I and MDA5, fish LGP2 may play dual roles in the type I IFN-mediated antiviral response, being similar to its mammalian counterpart (Rothenfusser *et al.* 2005; Satoh *et al.* 2010). Some reports showed that fish LGP2 can be a positive regulator of type I IFN response (Ohtani *et al.* 2010; Chang *et al.* 2011; Xiao *et al.* 2016). For example, the overexpression of Japanese flounder LGP2 in HINAE cells induced the expression of type I IFNs and ISGs, improving the antiviral ability against VHSV and hiramé rhabdovirus (HIRRV) infection (Ohtani *et al.* 2010). Likewise, RTG2 cells transfected with rainbow trout LGP2 are resistant to infection by VHSV, with the enhancement of Mx expression (Chang *et al.* 2011), and ectopic expression of black carp LGP2 in EPC cells provides an immune protection in

cells against SVCV and GCRV (Xiao *et al.* 2016). Conversely, there are some other reports which showed that fish LGP2 exerts inhibitory effect on antiviral response triggered by type I IFNs (Sun *et al.* 2011b; Yu *et al.* 2016a). For instance, the overexpression of gibel carp LGP2 in CAB cells decreased the activity of type I IFN promoter induced by poly(I:C), RIG-I or MDA5 (Sun *et al.* 2011b), and significant down-regulation of ISGs and increased severity of SGIV and RGNNV infections were observed in GS cells transfected with grouper LGP2 (Yu *et al.* 2016a).

The factor influencing the positive or negative role of fish LGP2 function in regulating type I IFN response is not yet fully understood, but may be related to the type and dose of pathogens as well as the stage of infection. In grass carp, the overexpression of LGP2 in CIK cells diminished the promoter activity and the expression of type I IFNs, as well as the synthesis and phosphorylation of IRF3 and IRF7 in the early phase of GCRV infection (Rao *et al.* 2017), but inhibited virus replication and protected cells from death in the late phase of GCRV infection (Chen *et al.* 2015e). Conversely, zebrafish LGP2 functions as a positive regulator of type I IFN signalling at the beginning of SVCV infection, but it switches to a negative regulator along with the replication of SVCV in cells up to a certain titre (Zhang *et al.* 2018). In addition, the functional switch of fish LGP2 has been revealed to be relevant to the dose of stimulator, which is supported by the fact that zebrafish LGP2 enhanced the production of type I IFNs in the absence or presence of poly(I:C) at limited concentrations, but inhibited type I IFN response when the concentrations of poly(I:C) is relatively high (Zhang *et al.* 2018). Thus, fish LGP2 may serve as a regulator to control the magnitude of type I IFN response, not only guaranteeing sufficient production of IFN molecules to elicit effective protection, and also preventing excessive production of IFN molecules to cause potential toxicities.

Unsurprisingly, as a key downstream molecule, MAVS seems to have conserved function on RLR-mediated type I IFN response in fish. It has been demonstrated that fish MAVS is associated with both RIG-I and MDA5 (Zou *et al.* 2014b, 2015), and can enhance both RIG-I- and MDA5-mediated type I IFNs response (Zhang *et al.* 2014; Zou *et al.* 2014b, 2015); Chen *et al.* 2015d. Mechanistically, fish MAVS is located at mitochondria in dimmers, and is likely to activate IRF3 and IRF7 to trigger the expression of type I IFNs through the recruitment of TBK1 (Lauksund *et al.* 2009; Biacchesi *et al.* 2009, 2012; Simora *et al.* 2010; Xiang *et al.* 2011; Kasthuri *et al.* 2014; Zhang *et al.* 2014; Zou *et al.* 2014b, 2015); Chen *et al.* 2015d; Feng *et al.* 2015b; Lu *et al.* 2015; Zhou *et al.* 2015; Jia *et al.* 2016b; Xiao *et al.* 2017; Huang *et al.* 2018b; Krishnan *et al.* 2018. Consistently, fish TBK1 can interact with IRF3 and IRF7 (Sun *et al.* 2011b; Hu *et al.* 2018b; Yu *et al.* 2018b), and the

overexpression of TBK1 alone can significantly up-regulate the expression of type I IFNs and ISGs *in vitro* and *in vivo*, thereby providing an efficient protection against virus infection (Sun *et al.* 2011b; Feng *et al.* 2014a; Zhang *et al.* 2016; Yan *et al.* 2017a; Hu *et al.* 2018a,b; Yu *et al.* 2018b). In addition, tumour necrosis factor receptor-associated factor (TRAF) 2 and TRAF3 may be also involved in the MAVS-mediated antiviral signalling (Xiang *et al.* 2011; Chen *et al.* 2017a).

However, two important and intriguing questions regarding fish RLR-mediated type I IFN response remain to be answered. Firstly, since RIG-I appears to be lost in certain groups of fish species (Zou *et al.* 2009; Chen *et al.* 2017b), it would be interesting to clarify the mechanism to compensate the deficiency of RIG-I in these species. Secondly, much more research is still required to fully elucidate the dual roles of fish LGP2 in type I IFN-mediated antiviral response, and to understand the factors influencing the functional switch of LGP2 in fish.

TLR-mediated type I IFN response

TLRs are type I integral membrane proteins comprising of an ectodomain, which contains leucine-rich repeats (LRRs) for the recognition of ligands, a transmembrane region, and cytosolic Toll-IL-1 receptor (TIR) domains which mediate downstream signalling pathways (Kawai & Akira 2010). Currently, 13 functional TLRs, named as TLR1–13, have been identified in human and mouse (Kawai & Akira 2010), in which TLR3, TLR7, TLR8 and TLR9 are the sensors of nucleic acids with the ability to initiate type I IFN response (Kawai & Akira 2011). TLR3 senses double-stranded RNA (dsRNA) and signals via TRIF, leading to the activation of IRF3 and the production of type I IFNs (Kawai & Akira 2011). By comparison, TLR7/8 and TLR9 recognize single-strand RNA (ssRNA) and cytosine-phosphate-guanosine (CpG)-containing DNA, respectively, and recruit MyD88 to activate IRF7-dependent type I IFN response (Kawai & Akira 2011).

Fish possess a larger TLR repertoire, with almost all orthologues of mammalian TLRs (TLR1–5, and 7–9) and non-mammalian TLRs (at least TLR5S, 18–23, and 25, 26) (Pietretti & Wiegertjes 2014). Among them, TLR3, TLR19 and TLR22 are verified to be involved in the activation of type I IFN response (Ji *et al.* 2018; Matsuo *et al.* 2008). The studies from fugu (*Takifugu rubripes*) and grass carp have shown that fish TLR3, TLR19 and TLR22 can all recognize dsRNA and recruit TRIF to elicit type I IFN response (Matsuo *et al.* 2008; Ji *et al.* 2018). TBK1 may be also involved in the TRIF-mediated type I IFN response, because it was proven that TRIF can bind TBK1 *in vitro* (Sullivan *et al.* 2007). The main difference among these three TLRs lies in their cellular localization and ligand preference. Firstly, fish

TLR3 resides in endoplasmic reticulum, whereas TLR19 is present in endosome, and TLR22 is localized exclusively in plasma membrane (Matsuo *et al.* 2008; Ji *et al.* 2018). Secondly, fish TLR3 senses preferentially short dsRNA, while TLR22 recognizes relatively long dsRNA (Matsuo *et al.* 2008). Given that mammalian TLR3 is expressed in cytoplasmic compartments as well as on the plasma membrane of fibroblasts (Funami *et al.* 2004), TLR22 in fish might be a functional substitute for cell surface-expressed mammalian TLR3.

However, the ligand recognition and type I IFN-inducing activity of TLR3 and TLR22 were only defined in fugu, and TLR19 only in grass carp. The functional properties of these TLRs remain to be further characterized in other species of fish. Additionally, although fish TLR9 can bind CpG-containing DNA as in mammals (Iliev *et al.* 2013; Yeh *et al.* 2013), it remains to be demonstrated whether it can activate the production of type I IFNs. Future studies are also required to determine whether fish TLR7 and TLR8 can conservatively recognize ssRNA and trigger the type I IFN response.

Cytosolic DNA sensor-mediated type I IFN response

Apart from TLR9-mediated CpG-containing DNA sensing, a series of cytosolic DNA sensors are discovered and characterized functionally in mammals, including absent in melanoma 2 (AIM2), DNA-dependent activator of IFN-regulatory factors (DAI), interferon gamma-inducible protein 16 (IFI16), cyclic GMP-AMP synthase (cGAS), DEAD-box helicase 41 (DDX41) and DExH-box helicase 9 (DHX9) (Wu & Chen 2014). In downstream of these DNA sensors, the stimulator of IFN genes (STING, also known as MITA), which is a transmembrane protein located in the endoplasmic reticulum, acts as an adapter and triggers the production of type I IFNs through the activation of IRF3 (Ran *et al.* 2014).

In contrast with the significant advance in mammals, the understanding of cytosolic DNA sensors in fish is rather limited. Among the above-mentioned DNA sensors, only DDX41 and DHX9 have been shown to be involved in type I IFN-mediated antiviral response in fish (Ge *et al.* 2015; Quynh *et al.* 2015; Lazarte *et al.* 2017; Ma *et al.* 2018). In zebrafish, the induced expression of type I IFNs and ISGs following the stimulation of dsDNA virus infection or synthetic dsDNA analogue was significantly reduced by the separate knockdown of DDX41 and DHX9 (Ge *et al.* 2015; Ma *et al.* 2018), and was almost abolished in the DDX41/DHX9 double knockdown morphant (Ge *et al.* 2015). Consistently, the activity of the type I IFN promoter as well as the expression of type I IFNs were enhanced in Japanese flounder DDX41-overexpressing HINAE cells with the treatment of C-di-GMP (Quynh *et al.* 2015), and Japanese

flounder DDX41 is an effective adjuvant for the viral glycoprotein-based DNA vaccine against VHSV infection (Lazarte *et al.* 2017). Surprisingly, zebrafish cGAS seems to be dispensable for type I IFN production in response to dsDNA virus infection (Ge *et al.* 2015). In consideration of the importance of cGAS-STING pathway in the mammalian DNA sensing system (Chen *et al.* 2016), it appears possible that the foreign DNA sensing system may differ at least to some extent in fish.

Expectedly, fish STING is similar to the mammalian counterpart, functioning as a critical adapter molecule in type I IFN signalling. Mechanistically, fish STING recruits TBK1 and IRF3 to trigger the expression of type I IFNs and ISGs (Sun *et al.* 2011b; Biacchesi *et al.* 2012; Feng *et al.* 2014b; Ge *et al.* 2015; Huang *et al.* 2015a; Lu *et al.* 2017; Xu *et al.* 2017). Two serine residues (S₃₇₉ and S₃₈₇ in orange-spotted grouper, and S₃₇₁ and S₃₇₉ in black carp) in fish STING are essential for STING-induced type I IFN activation (Huang *et al.* 2015a; Lu *et al.* 2017). A new mechanism of fish STING-mediated type I IFN response was uncovered by a recent study from grass carp, in which STING was co-localized with zinc finger DHHC-type containing 1 (ZDHHC1) in endoplasmic reticulum, where it interacts with ZDHHC1 to produce the STING/ZDHHC1 heterodimer (Xu *et al.* 2017). This heterodimer complex elicits nuclear translocation of IRF3, thus initiating the transcription of type I IFNs (Xu *et al.* 2017). Interestingly, it has been reported that fish STING is also involved in RLR-mediated type I IFN response (Sun *et al.* 2011b; Biacchesi *et al.* 2012), suggesting that there exists a crosstalk between the viral RNA and DNA sensing system in fish.

Nevertheless, there is no direct experimental evidence to define the ligand recognition by cytosolic DNA sensors in fish. Moreover, the importance of fish STING in viral DNA sensor-mediated signalling remains to be established, and it is still unclear how fish cytosolic DNA sensors and RLRs cooperatively regulate type I IFN-mediated antiviral response.

Regulation of type I IFN response by IRFs

IRFs are a family of transcription factors playing diverse roles in the regulation of innate and adaptive immune responses (Ikushima *et al.* 2013). To date, 11 IRF members have been identified in vertebrates (Huang *et al.* 2010), among which, IRF3 and IRF7 are two best-characterized members in controlling the transcription of type I IFN genes (Ikushima *et al.* 2013). Based on the canonical model of IRF3/7-mediated type I IFN induction in mammals, IRF3 is constitutively expressed and is not up-regulated by type I IFNs, while IRF7 is considered as a typical type I IFN-induced gene. At the beginning of viral infection, constitutively expressed IRF3 is activated by virus-induced

phosphorylation, resulting in the rapid induction of IFN β and some IFN α genes, which are called early-phase IFN genes. These induced early-phase IFN genes then trigger a strong expression of IRF7, thus priming the expression of most IFN α genes, which exhibit delayed induction and are called late-phase IFN genes (Marie *et al.* 1998; Sato *et al.* 2000; Honda *et al.* 2005).

Analogously, IRF3 and IRF7 are also primary transcriptional factors priming the type I IFN response in fish. It has been shown that fish IRF3 or IRF7 can bind and activate type I IFN promoter, thereby increasing the expression of type I IFNs and ISGs as well as antiviral activity against viral infection (Holland *et al.* 2008; Kileng *et al.* 2009; Bergan *et al.* 2010; Sun *et al.* 2010, 2011b; Cui *et al.* 2011; Hu *et al.* 2011; Iliev *et al.* 2011; Feng *et al.* 2016; Zhao *et al.* 2017; Wang *et al.* 2018). A striking divergence between fish and mammals is that both IRF3 and IRF7 are type I IFN-induced genes in fish (Holland *et al.* 2008; Sun *et al.* 2010; Ding *et al.* 2016; Laghari *et al.* 2018). Importantly, different fish type I IFN members appear to be regulated by distinct IRFs. In the case of zebrafish, IFN1 (group I IFN) promoter is primarily primed by IRF3, being similar to the fashion of mammalian IFN β , and IFN3 (group II IFN) promoter is predominantly primed by IRF7, resembling those of mammalian IFN α (Sun *et al.* 2011b; Feng *et al.* 2016). Additionally, fish IRF1 has been shown to be a key transcriptional factor involved in the activation of type I IFNs (Caipang *et al.* 2005; Bergan *et al.* 2010; Lai *et al.* 2013; Kung *et al.* 2014; Feng *et al.* 2015a, 2016). Akin to IRF3 and IRF7, fish IRF1 can bind directly to type I IFN promoter and activate type I IFN-mediated antiviral response (Caipang *et al.* 2005; Bergan *et al.* 2010; Lai *et al.* 2013; Kung *et al.* 2014; Feng *et al.* 2015a, 2016). Interestingly, these three IRFs can become either homo- or hetero-dimerized to recruit the adaptor myeloid differentiation primary response 88 (MyD88), cooperatively tuning the type I IFN response (Iliev *et al.* 2011; Feng *et al.* 2015a, 2016). Recent studies have shown that fish IRF5 and IRF6 are likely to be the positive regulators of type I IFN response (Wickramaarachchi *et al.* 2014; Li *et al.* 2016; Ai *et al.* 2018), but more experimental data from different fish species may be needed to further support this notion.

Negative regulation of type I IFN response

Type I IFNs have the capacity to activate the expression of ISGs (Schoggins *et al.* 2011), but uncontrolled type I IFN response can be harmful to host and may cause autoinflammation and tissue damage (Kretschmer & Lee-Kirsch 2017). To minimize potential detrimental outcome, the strength and duration of type I IFN response must be tightly controlled by a range of negative regulatory

molecules (Hertzog & Williams 2013). Fish appear to possess a finely balanced regulatory network to tune the immune response (Rebl & Goldammer 2018), and several negative regulation mechanisms of type I IFN response have been elucidated recently in fish.

Among several hundred ISGs induced by mammalian type I IFNs, suppressor of cytokine signalling (SOCS) 1 and ubiquitin carboxy-terminal hydrolase (USP) 18 are typical negative regulators of type I IFN signalling (Hertzog & Williams 2013; Yim *et al.* 2016; Yu *et al.* 2018a). The inhibitory role of SOCS1 and USP18 on type I IFN response seems to be conserved in fish as observed in mammals. Unsurprisingly in fish, both SOCS1 and USP18 can be significantly induced by type I IFNs (Nie *et al.* 2014; Skjesol *et al.* 2014; Chen *et al.* 2015a; Sobhkhez *et al.* 2017a). A preliminary study from zebrafish has shown that the overexpression of SOCS1 in embryos suppressed markedly the IFN1-induced ISG activation (Nie *et al.* 2014). Simultaneously, the suppressive mechanism of SOCS1 in fish was uncovered in studies from Atlantic salmon, in which SOCS1 can directly interact with TYK2 and STAT1, and inhibit type I IFN-induced nuclear localization of STAT1, thus strongly impairing type I IFN signalling and antiviral defense (Skjesol *et al.* 2014; Sobhkhez *et al.* 2017a). It has recently been revealed that gibel carp USP18 can dramatically down-regulate the induced expression of ISGs triggered by type I IFNs (Chen *et al.* 2015a), but the molecular mechanism of this inhibitory process is yet to be determined.

Similarly, not all the members of IRF family are transcriptional factors to prime the expression of type I IFNs, and some of them can even inhibit the activation of type I IFN response. For example, zebrafish IRF10 can interact with STING to block the activation of IFN1 and IFN3, and can also inhibit the induction of IFN1 and IFN3 by binding to the ISRE motif in their promoters (Li *et al.* 2014a). Analogously, grass carp IRF2 binds to the promoter of IFN1, thus down-regulating the expression of IFN1 and ISGs (Gu *et al.* 2015). On the other hand, some transcriptional factors from other gene families may play antagonistic roles in the IRF3/7-induced type I IFN activation. For instance, zebrafish forkhead box O (FOXO) 3 can interact directly with IRF3 and IRF7, thereby suppressing IRF3/7-mediated type I IFN response (Liu *et al.* 2016). Consistently, the higher expression of IFN1 and ISGs were observed in FOXO3-null zebrafish larvae, which display stronger antiviral capability when compared with wild-type larvae (Liu *et al.* 2016).

As stated above, fish RLR signalling plays vital roles in the induction of type I IFN expression, which is strictly controlled by several negative regulatory molecules. As

revealed in zebrafish model, the spliced variants of some RLR members and key downstream molecules were identified as the negative regulators of type I IFN response. For instance, the wild-type LGP2 in zebrafish has dual functions in the regulation of type I IFN signalling, but its two spliced isoforms LGP2v1 and LGP2v2 only retain the inhibitory role (Zhang *et al.* 2018). Similarly, the overexpression of zebrafish MAVS_tv2, a spliced variant of wild-type MAVS, suppressed the activation of IFN1 promoter and the enhancement of IFN1 and ISG production triggered by IRF7 (Chen *et al.* 2015d). By contrast, the negative regulation of fish TBK1 may be much more complicated, which is related with the two spliced variants of zebrafish TBK1, namely TBK1_tv1 and TBK1_tv2, and a paralogous gene, termed TBK1L (Zhang *et al.* 2016; Hu *et al.* 2018b). Mechanistically, TBK1_tv1 and TBK1_tv2 can competitively interact with TBK1 and IRF3 to disturb TBK1-mediated IRF3 phosphorylation, thereby suppressing the activation of type I IFN signalling, and TBK1L has a similar inhibitory effect on TBK1-mediated type I IFN response (Zhang *et al.* 2016; Hu *et al.* 2018b). Similarly, zebrafish STAT6 can become associated with TBK1, thereby impeding the phosphorylation of IRF3 and the expression of IFN1 (Li *et al.* 2017). In addition, zebrafish protein inhibitor of activated STAT4a (PIAS4a) and von Hippel–Lindau (VHL) may be involved also in the repression of RLR signalling, which is supported by the fact that the co-overexpression of PIAS4a and MAVS in embryos reduced dramatically the MAVS-induced IFN1 expression (Xiong *et al.* 2012), and that VHL induced the degradation of MAVS *in vitro* and inhibited type I IFN-mediated antiviral response *in vivo* (Du *et al.* 2015). Apart from the zebrafish model, studies from orange-spotted grouper have shown that several members of tripartite motif (TRIM) protein family, including TRIM13, TRIM16L, TRIM35 and TRIM62, may negatively regulate type I IFN response by suppressing the RLR signalling (Huang *et al.* 2016a, 2017b; Yang *et al.* 2016b; Yu *et al.* 2016b).

Indeed, fish type I IFN signalling is tightly controlled at virtually every step of the signal process, including ligand production, signalling adaptors and enzymes, and transcriptional factors. However, the inhibitory mechanisms of type I IFN signalling at the level of receptors are still poorly understood in fish and require further investigation. Interestingly, a more recent study from gibel carp has shown that the type I IFN promoter activity could be suppressed by diverse immunoglobulin domain-containing proteins (DICPs) (Gao *et al.* 2018), which belong to a fish-specific gene family (Rodriguez-Nunez *et al.* 2016). It seems likely that fish may have some lineage-specific mechanisms in type I IFN signalling.

Expression characters of type I IFNs

Based on the current knowledge in mammals and birds, most type I IFNs are ubiquitously expressed at a relatively low level in unstimulated organs/tissues compared with housekeeping gene, and are highly induced by poly(I:C) stimulation or viral infection (Pestka *et al.* 2004; Teijaro 2016; Secombes & Zou 2017). This basic rule may be also true for most type I IFNs in fish, but extensive studies have revealed a diversity in the pattern of type I IFN expression in fish (Altmann *et al.* 2003; Lutfalla *et al.* 2003; Robertsen *et al.* 2003; Long *et al.* 2006, 2004; Gonzalez *et al.* 2007; Zou *et al.* 2007, 2014a; Casani *et al.* 2009; Chang *et al.* 2009; Kitao *et al.* 2009; Purcell *et al.* 2009; Li *et al.* 2012; Wan *et al.* 2012; Chen *et al.* 2014; Parhi *et al.* 2014; Pereira *et al.* 2014; Valero *et al.* 2015; Huang *et al.* 2015b, 2018; Ding *et al.* 2016, 2017; Kuo *et al.* 2016; Liao *et al.* 2016; Maekawa *et al.* 2016; Wei *et al.* 2016; Yan *et al.* 2016, 2017b; Hu *et al.* 2017; Laghari *et al.* 2018; Milne *et al.* 2018; Wu *et al.* 2018; Xia *et al.* 2018). Firstly, considerable divergence may exist in the expression of group I and II type I IFNs, and expression difference may even exist among the members in each groups. For instance, an initial study in rainbow trout indicated that IFN1 (also known as IFN α 1, belonging to group I IFNs) was not detected in all analysed tissues from unstimulated fish. By comparison, IFN2 (IFN α 2, group I IFNs) was detected in all examined tissues, and IFN3 (IFN β 1, group II IFNs) was mainly expressed in ovary and testis (Zou *et al.* 2007). Secondly, distinct expression is observed among fish type I IFNs in response to different immunological stimuli. Subtypes A of type I IFNs in Atlantic salmon were expressed differently upon the *in vivo* treatment of poly(I:C) (a ligand for TLR3/19/22 in fish) or S-27609 (a putative ligand for TLR7) (Sun *et al.* 2009). Following the stimulation of poly(I:C), IFN α was strongly induced during the first 24 h followed by decline, whereas IFN β was only slightly induced at 14 h, and IFN γ level persistently increased during the experimental period. In contrast, the treatment with S-27609 led to a persistent up-regulation of IFN α , a transient but very strong induction of IFN β , and a short induction of IFN γ (Sun *et al.* 2009). Thirdly, different patterns in the expression may be observed in some type I IFN members at different life stages of fish (Aggad *et al.* 2009). IFN2 in zebrafish is not expressed and not induced by viral infection in larvae, but it is expressed at relatively high level and well induced by viral infection in adults (Aggad *et al.* 2009).

In mammals, nearly all nucleated cells can secrete type I IFNs, but the capacity in producing the cytokines may vary in different cell types (Liu 2005). Plasmacytoid dendritic cells (pDCs) are considered as a predominant source of type I IFNs, which can rapidly produce a large amount of IFN α and IFN β by TLR7/TLR8/TLR9–MyD88 signalling

pathway upon viral infection (Liu 2005). Intriguingly, cell type-dependent expression of type I IFNs may also exist in fish (Zou *et al.* 2007; Svingerud *et al.* 2012; Palha *et al.* 2013). The expression of type I IFNs in different cell types were initially compared in a study from rainbow trout, and all examined type I IFNs, including IFN1 and IFN2 (group I IFNs) and IFN3 (group II IFNs), were expressed in primary head kidney leukocytes following poly(I:C) stimulation, but only IFN1 and IFN2 were expressed in RTG-2 cells, a kind of fibroblast-like cell line (Zou *et al.* 2007). More clearly, as revealed in a transgenic reporter zebrafish, hepatocytes and neutrophils were identified as the two main cell types expressing IFN1 in larvae upon chikungunya virus (CHIKV) infection (Palha *et al.* 2013). In addition, a fluorescence *in situ* hybridization (FISH) study in Atlantic salmon has shown that IFN α and IFN γ were expressed in a minority of cells in head kidney, spleen, gills, liver and heart from poly(I:C)-stimulated fish, with some cells co-expressing these two IFNs (Svingerud *et al.* 2012). Importantly, IFN β and IFN γ were co-expressed in a few cells of head kidney and spleen from R848-stimulated fish (Svingerud *et al.* 2012), and these high IFN β /IFN γ -producing cells in the lymphoid tissues may be similar to mammalian pDCs, functioning as specialized high type I IFN producers in fish since R848 is a putative ligand for TLR7. Despite the scarcity in understanding type I IFN-producing cells in fish, it seems true, as observed in mammals, that different members of type I IFNs may be expressed in certain types of cells, which may need to be further clarified.

Functional properties of type I IFNs

The well-demonstrated function of type I IFNs is their antiviral activity. Type I IFNs can establish an antiviral state in cells through the induction of antiviral ISGs. Similar to mammals, fish possess a large repertoire of ISGs, which is supported by a recent study in which over 2000 genes were up-regulated by recombinant salmon IFN α in TO cells (Xu *et al.* 2015). A small subset of ISGs have been shown to possess direct antiviral effect in fish, such as Mx, viperin, ISG15, PKR and PKZ (Larsen *et al.* 2004; Liu *et al.* 2011, 2018; Schoggins & Rice 2011; Langevin *et al.* 2013b; Wang *et al.* 2014). Obviously, the antiviral activity of fish type I IFNs is closely related to the antiviral activity of their induced ISGs.

Currently, the antiviral activity of type I IFNs has been clearly defined in a wide range of species, including cyprinids, salmonids and perciform fish. The understanding of antiviral activity of type I IFNs in cyprinid fish is mainly derived from the studies in zebrafish (Altmann *et al.* 2003; Wang *et al.* 2006; Levraud *et al.* 2007; Aggad *et al.* 2009; Lopez-Munoz *et al.* 2009, 2010; Li *et al.* 2010; Palha *et al.* 2013). Among the four type I IFN members in zebrafish,

IFN2 and IFN3 (group II IFN, subgroup c) can elicit a rapid and transient expression of antiviral ISGs to protect fish against SVCV and infectious haematopoietic necrosis virus (IHNV) (Aggad *et al.* 2009; Lopez-Munoz *et al.* 2009, 2010). By contrast, IFN1 (group I IFN, subgroup a) induces a delayed but higher level of antiviral ISGs, thus providing an efficient protection against different viruses including SVCV, IHNV, snakehead rhabdovirus (SHRV), infectious spleen and kidney necrosis virus (ISKNV), and CHIKV *in vitro* and *in vivo* (Altmann *et al.* 2003; Wang *et al.* 2006; Levraud *et al.* 2007; Aggad *et al.* 2009; Li *et al.* 2010; Lopez-Munoz *et al.* 2009, 2010; Palha *et al.* 2013). It is indicated that the two type I IFN groups may have complementary antiviral roles at different stages of virus infection. Nevertheless, zebrafish IFN4 (group I IFN, subgroup d) could not protect embryos against IHNV infection, which may be resulted from a much weaker up-regulation of antiviral ISGs (Aggad *et al.* 2009). In addition, the ISG-induced activity and antiviral activity of type I IFNs have been characterized also in other cyprinids, including common carp, grass carp, black carp, gibel carp, and red crucian carp (Kitao *et al.* 2009; Li *et al.* 2013; Huang *et al.* 2015b; Yan *et al.* 2016, 2017b; Wu *et al.* 2018; Xia *et al.* 2018). Consistently, all these type I IFNs belong to subgroups a or c (Kitao *et al.* 2009; Li *et al.* 2013; Huang *et al.* 2015b; Yan *et al.* 2016, 2017b; Wu *et al.* 2018; Xia *et al.* 2018), and to date no IFNd in cyprinids has been proven to have antiviral activity.

As described above, salmonids appear to have the most complexed type I IFN repertoire containing subgroups a, b, c, d, e and f (Sun *et al.* 2009; Svingerud *et al.* 2012; Zou *et al.* 2014a), which is a remarkable model to study functionalization of type I IFNs in fish. Most research regarding antiviral activity of type I IFNs in salmonids has been conducted in Atlantic salmon (Ooi *et al.* 2008; Xu *et al.* 2010a; Svingerud *et al.* 2012, 2013; Chang *et al.* 2014, 2016), with discovery of considerable divergence in antiviral effects among different type I IFN subgroups. *In vitro* experiments have shown that salmon IFNa and IFNc display potent antiviral activity against infectious pancreatic necrosis virus (IPNV) and salmonid alphavirus (SAV) 3 (Ooi *et al.* 2008; Xu *et al.* 2010a; Svingerud *et al.* 2012; Chang *et al.* 2016), but only IFNa can protect fish cells against infectious salmon anaemia virus (ISAV) infection (Svingerud *et al.* 2013). In contrast, IFNb induces lower antiviral effect against IPNV than IFNa and IFNc *in vitro* (Svingerud *et al.* 2012). Nevertheless, distinct results have been obtained when testing antiviral activity of IFNs *in vivo* by injecting fish intramuscularly with expression plasmids expressing IFNa, IFNb and IFNc, respectively (Chang *et al.* 2014). All three IFNs induced similarly the expression of antiviral ISGs at the muscle injection site, but only IFNb and IFNc up-regulated the expression of antiviral ISGs in head

kidney, liver, heart and pancreas of injected fish (Chang *et al.* 2014), suggesting that IFNa is active mainly at the injection site, whereas IFNb and IFNc can be systemically active. In addition, IFNc plasmid induced stronger Mx protein expression in heart and liver than IFNb plasmid (Chang *et al.* 2014). Being consistent to the antiviral ISG-induced activity of these IFNs, the treatment of fish with IFNc plasmids before ISAV or SAV3 infection gave a strong protection, but IFNb a weak protection, and IFNa no protection (Chang *et al.* 2014, 2016). Interestingly, salmon IFNd showed no ISG-inducing ability and antiviral activity against IPNV in fish cells (Svingerud *et al.* 2012), and the antiviral activity of IFNe and IFNf in salmonids remains to be assessed.

Apart from cyprinids and salmonids, type I IFNs have been characterized in a few species of perciform fish, which appear to harbour subgroups c, d and h (Casani *et al.* 2009; Wan *et al.* 2012; Chen *et al.* 2014; Valero *et al.* 2015; Ding *et al.* 2016, 2017; Kuo *et al.* 2016; Laghari *et al.* 2018; Milne *et al.* 2018). Recently, it has been shown that IFNc and IFNh in perciform fish can trigger a strong expression of antiviral ISGs to have an efficient protection against viral infection (Ding *et al.* 2017; Laghari *et al.* 2018). By contrast, conflicting results have been reported on the function of IFNd in this lineage of fish. IFNd has been proven to have ISG-inducing activity and antiviral activity in several perciform fish including rock bream, orange-spotted grouper, seven band grouper and large yellow croaker (Wan *et al.* 2012; Chen *et al.* 2014; Ding *et al.* 2016; Kuo *et al.* 2016), and in some species from other orders, such as channel catfish and medaka (Long *et al.* 2006; Maekawa *et al.* 2016). However, a most recent study showed that mandarin fish IFNd has no effect on the phosphorylation of STAT1 and STAT2, nor on the expression of antiviral ISGs, thus providing no protection against viral infection (Laghari *et al.* 2018), which is similar to IFNd in zebrafish and Atlantic salmon as mentioned above (Aggad *et al.* 2009; Svingerud *et al.* 2012). It is interesting to investigate the mechanism underlying the functional divergence of IFNd in these species of fish.

Extensive studies have shown that recombinant type I IFNs or type I IFN expression plasmids when injected into fish can provide *in vivo* efficient protection against viral infection (Levraud *et al.* 2007; Aggad *et al.* 2009; Lopez-Munoz *et al.* 2009, 2010; Li *et al.* 2010, 2013; Palha *et al.* 2013; Chang *et al.* 2014, 2016; Pereiro *et al.* 2014; Kuo *et al.* 2016; Xia *et al.* 2018), indicating that fish type I IFNs are promising candidates for potential antiviral agents in the prevention of infectious viral diseases in aquaculture. More intriguingly, fish type I IFNs seem to be able to trigger adaptive immune response, and hence have the potential as novel adjuvants in antiviral vaccines. This notion is supported by recent studies from Atlantic salmon, in which

the fish were immunized with the injection of a plasmid encoding the haemagglutinin-esterase (HE) gene of ISAV alone or together with a plasmid expressing either IFN α , IFN β or IFN γ (Chang *et al.* 2015; Robertsen *et al.* 2016; Sobhkhez *et al.* 2017b). Remarkably, HE plasmid alone only provided a weak protection against ISAV infection, with the induction of low IgM antibody titre, whereas all three IFN plasmids together with HE plasmid enhanced markedly the protection with a strong and prolonged antibody response against ISAV (Chang *et al.* 2015; Robertsen *et al.* 2016). The mechanism behind the adjuvant effect of fish type I IFNs is not yet fully understandable, but may be related to the increasing expression of chemokines induced by type I IFNs, which may lead to the attraction of lymphocytes at the muscle injection site (Chang *et al.* 2015; Sobhkhez *et al.* 2017b).

Type II IFNs in fish

Discovery of type II IFNs

IFN- γ , belonging to the type II IFN family, is encoded by a single copy gene with the genomic structure of four exons and three introns in mammals (Savan *et al.* 2009). Human *IFNG* is located on chromosome 12 containing another two members of class II cytokine family, IL-22 and IL-26. Mammalian IFN- γ is a multi-functional cytokine. For example, IFN- γ can modulate CD4⁺ T helper 1 (Th1) cell response and is pivotal for the clearance of intracellular pathogens (Boehm *et al.* 1997). On the other hand, IFN- γ is a key regulator for macrophage activation (Green *et al.* 2017) and for promoting antigen presentation process to inhibit viral replication and tumour growth by the induction of immune-related ISGs (Schoenborn & Wilson 2007).

The first identified type II IFN in fish was the IFN- γ gene in fugu by gene synteny analysis (Zou *et al.* 2004). In fact, type II IFN loci have conserved synteny in vertebrates from fish to mammals (Savan *et al.* 2009; Chen *et al.* 2013). However, a unique gene as the new member, named as IFN- γ 1 (also known as IFN- γ related gene, IFN- γ rel, see below), of type II IFN family was identified at the up-stream of IFN- γ 2 gene (the orthologous gene of fugu IFN- γ) in zebrafish and pufferfish (Igawa *et al.* 2006). Indeed, the teleost IFN- γ rel (or the so-called IFN- γ 1) contains IFN- γ signature motif and has moderate sequence similarity with IFN- γ (IFN- γ 2) which possesses more conserved vertebrate IFN- γ sequence features, such as nuclear localization signal (NLS) motif at C-terminal region (Igawa *et al.* 2006; Zou & Secombes 2011; Chen *et al.* 2013). The NLS sequence of IFN- γ is conserved from fish to human, with usually several continuously encoded lysines or arginines (for example, RRKR and KRKR in trout and human respectively), which have been proven to be essential for the function of fish and mammal IFN- γ (Subramaniam *et al.* 1999; Zou *et al.* 2005), although

ginbuna crucian carp (*C. auratus langsdorfi*) IFN- γ rel1 (one of IFN- γ rel isoforms) also has functional NLS sequence (KHHHR) at C-terminal region (Shibasaki *et al.* 2014). On the other hand, more than two copies of fish type II IFN gene are present in several species of fish (Zou & Secombes 2011). To avoid any confusion, it is suggested that fish IFN- γ 1 and IFN- γ 2 gene were renamed as IFN- γ related molecule (IFN- γ rel) and IFN- γ , respectively (Zou & Secombes 2011), and this nomenclature has been generally accepted among fish immunologists.

To date, IFN- γ genes have been identified in many different species of fish, such as goldfish (*C. auratus*), ginbuna crucian carp, grass carp, Atlantic cod (*Gadus morhua*), Atlantic salmon, Atlantic halibut (*Hippoglossus hippoglossus*), black seabream (*Acanthopagrus schlegelii*), channel catfish, common carp, medaka, large yellow croaker, rainbow trout, orange-spotted grouper, Japanese flounder, turbot, Nile tilapia (*Oreochromis niloticus*), European seabass and barramundi perch (*Lates calcarifer*) (Table 2) (Zou *et al.* 2005; Milev-Milovanovic *et al.* 2006; Robertsen 2006; Stolte *et al.* 2008; Furnes *et al.* 2009; Grayfer & Belosevic 2009a; Matsuyama *et al.* 2009; Yabu *et al.* 2011; Overgard *et al.* 2012a; Yang *et al.* 2013; Kuznetsova *et al.* 2014; Nuñez Ortiz *et al.* 2014; Chen *et al.* 2015c; Mohapatra *et al.* 2015; Pereiro *et al.* 2016; Velazquez *et al.* 2017; Xiang *et al.* 2017; Zhong *et al.* 2017; Peng *et al.* 2018). Similar to other vertebrate counterparts, teleost IFN- γ has a conserved signal peptide, IFN- γ signature sequence and NLS motif (continuous lysines or arginines). Among the species with known IFN- γ gene structure, four exons and three introns are observed, except in medaka which has two splicing forms of IFN- γ gene (designated as IFN- γ 2a and 2b in the original publication) derived from eight exons and six exons, respectively, by comparing to the genomic sequences (Mohapatra *et al.* 2015). It seems likely that the two isoforms are chimeric sequences based on the *IFNG* locus data from medaka RNA-seq exon coverage (NCBI *Oryzias latipes* Annotation Release 103) and genomic assembly (ASM223467v1). Although it is unclear whether all teleost IFN- γ genes possess conserved four-exon gene organization, the splicing of IFN- γ gene transcripts is likely to have effect on biological activities of IFN- γ .

Except for the presence in zebrafish, fugu and spotted green pufferfish, IFN- γ rel genes were also discovered in channel catfish, common carp, goldfish, ginbuna crucian carp, grass carp, rohu and orange-spotted grouper (Table 2) (Chen *et al.* 2010; Grayfer *et al.* 2010; Milev-Milovanovic *et al.* 2006; Peng *et al.* 2018; Shibasaki *et al.* 2014; Stolte *et al.* 2008; Swain *et al.* 2015; Yabu *et al.* 2011). These teleost IFN- γ rel molecules contain conserved signal peptide and IFN- γ signature sequences, but without NLS motif of continuous lysines or arginines. In addition, it was failed to identify IFN- γ rel gene in other different

Table 2 Currently reported type II IFNs in fish

Order	Species	Gene	GenBank accession number	References
Cypriniformes	Zebrafish (<i>D. rerio</i>)	IFN- γ 1 (IFN- γ rel)	NM_001020793	Igawa <i>et al.</i> (2006)
		IFN- γ 2 (IFN- γ)	NM_212864	Igawa <i>et al.</i> (2006)
	Grass carp (<i>C. idella</i>)	IFN- γ rel	FJ695519	Chen <i>et al.</i> (2010)
		IFN- γ 2 (IFN- γ)	JX196701.1	Yang <i>et al.</i> (2013)
	Common carp (<i>C. carpio</i>)	IFN- γ 1 (IFN- γ rel)	AM261214	Stolte <i>et al.</i> (2008)
		IFN- γ 2 (IFN- γ)	AM168523	Stolte <i>et al.</i> (2008)
	Crucian carp (<i>C. auratus</i>)	IFN- γ 1 (IFN- γ rel)	GQ149696	Grayfer and Belosevic (2009a)
		IFN- γ 2 (IFN- γ)	EU909368	Grayfer and Belosevic (2009b)
	Ginbuna crucian carp (<i>C. auratus langsdorfii</i>)	IFN- γ rel1	AB570433	Yabu <i>et al.</i> (2011)
		IFN- γ rel2	AB614642	Shibasaki <i>et al.</i> (2014)
		IFN- γ 1 (IFN- γ)	AB570431	Yabu <i>et al.</i> (2011)
		IFN- γ 2 (IFN- γ)	AB570432	Yabu <i>et al.</i> (2011)
	Rohu (<i>L. rohita</i>)	IFN- γ rel	KJ874352	Swain <i>et al.</i> (2015)
Channel catfish (<i>I. punctatus</i>)		IFN- γ 1 (IFN- γ rel)	DQ124249	Milev-Milovanovic <i>et al.</i> (2006)
	IFN- γ 2a (IFN- γ)	DQ124250	Milev-Milovanovic <i>et al.</i> (2006)	
	IFN- γ 2b (IFN- γ)	DQ124251	Milev-Milovanovic <i>et al.</i> (2006)	
Salmoniformes	Rainbow trout (<i>O. mykiss</i>)	IFN- γ	AJ616215	Zou <i>et al.</i> (2005)
	Atlantic salmon (<i>S. salar</i>)	IFN- γ	AY795563	Robertsen (2006)
Gadiformes	Atlantic cod (<i>G. morhua</i>)	IFN- γ	FJ356235	Furnes <i>et al.</i> (2009)
Cichliformes	Nile tilapia (<i>O. niloticus</i>)	IFN- γ	NM_001287402.1	(Velazquez <i>et al.</i> 2017; Zhong <i>et al.</i> 2017)
Perciformes	Large yellow croaker (<i>L. crocea</i>)	IFN- γ	KM501500.2	Chen <i>et al.</i> (2015a,b,c,d,e)
	Orange-spotted grouper (<i>E. coioides</i>)	IFN- γ 1 (IFN- γ rel)	-	Peng <i>et al.</i> (2018)
		IFN- γ	JX013936	–
	European sea bass (<i>D. labrax</i>)	IFN- γ	KJ818329	Nuñez Ortiz <i>et al.</i> (2014)
	Barramundi perch (<i>L. calcarifer</i>)	IFN- γ	NM_001360734.1	Kuznetsova <i>et al.</i> (2014)
Spariformes	Black seabream (<i>A. schlegelii</i>)	IFN- γ	KY921614	Xiang <i>et al.</i> (2017)
Pleuronectiformes	Turbot (<i>S. maximus</i>)	IFN- γ	KX360748	Pereiro <i>et al.</i> (2016)
	Japanese flounder (<i>P. olivaceus</i>)	IFN- γ	AB435093	Matsuyama <i>et al.</i> (2009)
Tetraodontiformes	Atlantic halibut (<i>H. hippoglossus</i>)	IFN- γ	GU985450	Overgard <i>et al.</i> (2012)
	Spotted green pufferfish (<i>T. nigroviridis</i>)	IFN- γ 1 (IFN- γ rel)	KJ524454	Yi <i>et al.</i> (2014)
		IFN- γ 2 (IFN- γ)	KJ524455	Yi <i>et al.</i> (2014)
Fugu (<i>T. rubripes</i>)	IFN- γ	AJ616216.2	Zou <i>et al.</i> (2004)	

vertebrates, including elephant shark, tropical clawed frog, green anole (*Anolis carolinensis*), chicken (*Gallus gallus*) and human, (Chen *et al.* 2013; Qi & Nie 2008; Savan *et al.* 2009; Venkatesh *et al.* 2014). However, all these vertebrates have IFN- γ gene, and a scenario is accepted to explain the occurrence of IFN- γ rel in teleost fish that the teleost-specific IFN- γ rel gene is likely derived from the duplication of IFN- γ gene (Igawa *et al.* 2006; Savan *et al.* 2009). However, much more effort is needed to understand how and when the duplication event might have occurred.

Structure of type II IFNs

The recombinant human IFN- γ contains six helices (A-F) and is crystallized in the form of homodimer (Ealick *et al.* 1991). Similar to human, homologous dimerization of recombinant or native IFN- γ was observed in zebrafish, goldfish, ginbuna crucian carp and Japanese flounder (Grayfer & Belosevic 2009a,b; Yabu *et al.* 2011; Yoon *et al.*

2016; Zahradnik *et al.* 2018). The crystal structure of the flounder IFN- γ is also in general similar to that of other known IFN- γ , despite of some significant differences, including additional helix G at C-terminal region and a different angle between helices C and D, in the fold of flounder IFN- γ when compared with mammals (Zahradnik *et al.* 2018). However, there are converse opinions on the dimerization of IFN- γ rel. Recombinant goldfish IFN- γ rel is capable to form homodimer (Grayfer & Belosevic 2009a,b), while monomers of IFN- γ rels possess biological activity in ginbuna crucian carp (Shibasaki *et al.* 2014). To date, little is known in relation with the crystal structure of IFN- γ rel and type II IFN receptors in teleost fish.

Receptors and signalling of type II IFNs

In mammals, the homodimer of IFN- γ activates the canonical JAK-STAT signalling pathway through the tetrameric IFN- γ receptor (IFN- γ R) complex, which includes two

subunits of IFN- γ R1 and IFN- γ R2, respectively (Boehm *et al.* 1997; Green *et al.* 2017; Ivashkiv 2018; Stark *et al.* 1998). Human IFN- γ R1 is encoded by a single copy gene and is also called ligand binding chain due to its high affinity binding to IFN- γ , and its intracellular region has two key anchor sites (L₂₆₆PKS₂₆₉ and Y₄₄₀DKPH₄₄₄) for docking and activation of JAK1 and STAT1, respectively (Bach *et al.* 1997; Stark *et al.* 1998). In fact, the phosphorylation of JAK1 and STAT1 needs first the activation of JAK2, which binds to intracellular region of IFN- γ R2 (Stark *et al.* 1998). Activated STAT1 molecules become dimerized and can bind to the DNA sequences called gamma-interferon-activated site (GAS) to initiate the transcription of ISGs in the nucleus (Green *et al.* 2017). Furthermore, IFN- γ is also able to activate other transcriptional factors, such as STAT3 and the complex of STAT1 and IRF9 (Green *et al.* 2017).

Unlike in mammals, IFN- γ R1 gene in teleost was duplicated as two copies in consideration of phylogenetic and syntenic analyses, which were designated as CRFB13/IFNGR1-2/IFNGR1-like and CRFB17/IFNGR1-1/IFNGR1 respectively (Grayfer & Belosevic 2009a,b; Chen *et al.* 2013; Lu *et al.* 2014; Liao *et al.* 2016; Zahradnik *et al.* 2018), although four copies of IFN- γ R1 gene are present in some salmonid fish, such as Atlantic salmon, due to the separate duplication of IFNGR1-1 and IFNGR1-2 (Zahradnik *et al.* 2018). Actually, the two teleost IFN- γ R1 genes are separated on different loci, which all show conserved synteny when compared with other vertebrates: CRFB13/IFNGR1-2 gene is located adjacent to oligodendrocyte transcription factor 3 (OLIG3), and CRFB17/IFNGR1-1 gene is linked with LTV1 ribosome biogenesis factor (LTV1), respectively (Grayfer & Belosevic 2009a,b; Chen *et al.* 2013; Lu *et al.* 2014). Expectedly, many studies have shown that both CRFB13/IFNGR1-2/IFNGR1-like and CRFB17/IFNGR1-1/IFNGR1 are main subunits of type II IFN receptor complex based on experimental evidence revealed by various approaches, such as ligand-receptor cross-linking, receptor-specific knockdown/knockout, expression of receptors in xenogenic cells, ligand-receptor immunoprecipitation and affinity analysis (Gao *et al.* 2009; Grayfer & Belosevic 2009a,b; Aggad *et al.* 2010; Yabu *et al.* 2011; Lu *et al.* 2014; Sawamiphak *et al.* 2014; Zahradnik *et al.* 2018). However, the IFN- γ R1 model of teleost type II IFN receptor complex seems to be not very clear, or there is no universal model but species-specific ligand-receptor binding. For example, goldfish IFN- γ 2 (IFN- γ) and IFN- γ 1 (IFN- γ rel) can be cross-linked with IFNGR1-2 (CRFB13) and IFNGR1-1 (CRFB17) respectively *in vitro*, while no binding activity has been detected after the exchange of receptors (Grayfer & Belosevic 2009a,b). But in zebrafish, the limitation of CRFB17 translation has significant effect on signalling and function of both IFN- γ 2 (IFN- γ) and IFN- γ 1 (IFN- γ rel), and simultaneously IFN- γ 2 (IFN- γ) signalling also needs CRFB13 (Aggad *et al.* 2010).

Similarly, both IFNGR1-1 (CRFB17) and IFNGR1-2 (CRFB13) are likely to be associated with IFN- γ in pufferfish, and IFNGR1-1 is the main component of IFN- γ rel receptor complex (Lu *et al.* 2014). In addition, IFN- γ 1 (but this is not IFN- γ rel) and IFN- γ 2, the two isoforms of IFN- γ , use different IFN- γ R1s, IFNGR1-2 (IFNGR1-like/CRFB13) and IFNGR1-1 (IFNGR1/CRFB17), respectively, in gibel carp (Yabu *et al.* 2011). Anyway, the two teleost IFN- γ R1 genes may be originated from one ancestral gene, with similar protein features in having a signal peptide, an extracellular region with FNIII domains, a transmembrane region and an intracellular region. It is implied that cross-talk may occur possibly in the IFNGR1-1 or IFNGR1-2 mediated signal pathway for teleost type II IFNs.

In fact, the potential STAT1 binding sites (consensus sequences: YDxxP[K/H]) are found in intracellular region of both CRFB13/IFNGR1-2 and CRFB17/IFNGR1-1 (Grayfer & Belosevic 2009a,b; Aggad *et al.* 2010; Zou & Secombes 2011). Both IFN- γ and IFN- γ rel are able to induce STAT1 expression and its phosphorylation in goldfish, zebrafish etc. (Grayfer *et al.* 2010; Skjesol *et al.* 2010; Lopez-Munoz *et al.* 2011; Collins *et al.* 2014; Ruan *et al.* 2017). Simultaneously, mutants of GAS sequences on promoters or intron of teleost ISGs led to dramatically depressed activation of ISG luciferase reporter plasmids mediated by IFN- γ or STAT1 (Shi *et al.* 2013; Cheng *et al.* 2014; Ruan *et al.* 2017; Liu *et al.* 2018). Moreover, IFN- γ can also activate STAT2 and STAT3 (Sawamiphak *et al.* 2014; Sobhkhaz *et al.* 2014). The ISREs on promoters of some ISGs are also essential for response to recombinant IFN- γ (Castro *et al.* 2008). It is thus revealed that STAT1/3-involved signal transduction of type II IFN is likely to be conserved from teleost to human. However, the role of JAK1 in signalling of teleost type II IFNs is still unclear, although the intracellular region of IFN- γ R1s contains a conserved binding motif (consensus sequences: PxxL) for JAK1 (Aggad *et al.* 2010; Zou & Secombes 2011).

The fish IFN- γ R2 (also known as CRFB6) has been identified in zebrafish, grass carp, rainbow trout and Dabry's sturgeon, and it possesses a signal peptide, an extracellular region, a transmembrane region and an intracellular region (Gao *et al.* 2009; Aggad *et al.* 2010; Liao *et al.* 2016; Luo *et al.* 2018). The IFN- γ R2, as in human, is a single copy gene and links with transmembrane protein 50B (TMEM50B) gene on a conserved locus of vertebrate class II cytokine receptor, except in zebrafish (Stein *et al.* 2007; Chen *et al.* 2013; Lutfalla *et al.* 2003; Zahradnik *et al.* 2018). Unsurprisingly, IFN- γ R2 is a component of IFN- γ receptor complex (Gao *et al.* 2009; Aggad *et al.* 2010). However, the knockdown of all known class II cytokine receptor (CRFB1-2, 4-8 and 12-17), including CRFB6 (IFN- γ R2), could not block the signalling mediated by IFN- γ 1 (IFN- γ rel), except that the knockdown of CRFB17

(IFNGR1-1) in zebrafish led to the failure of IFN- γ 1 (IFN- γ rel) signalling (Aggad *et al.* 2010). It is bewildering that there might be another unknown IFN- γ 2-like receptor in teleost genome, or that CRFB17 (IFNGR1-1) alone is sufficient for the signal transduction of IFN- γ rel (Aggad *et al.* 2010), although IFN- γ 2 is essential for immune-related function of IFN- γ in mammals (Lu *et al.* 1998). In addition, zebrafish JAK2a, rather than JAK2b, plays a role in signalling transmission of IFN- γ (Aggad *et al.* 2010).

Functional properties of type II IFNs

Antiviral activity of type II IFNs

To date, it has been observed that the expression of teleost IFN- γ or IFN- γ rel genes was significantly induced by many kinds of viruses, including SVCV, GCRV, VHSV, red RGNNV, IHNV, crucian carp haematopoietic necrosis virus (CHNV) or synthetic mimics of virus, such as poly(I:C) in different species of fish (Zou *et al.* 2005; Igawa *et al.* 2006; Furnes *et al.* 2009; Grayfer & Belosevic 2009a,b; Purcell *et al.* 2009; Aggad *et al.* 2010; Chen *et al.* 2010; Xu *et al.* 2010b; Overgard *et al.* 2012b; Du *et al.* 2013; Wang *et al.* 2013; Shibasaki *et al.* 2014; Somamoto *et al.* 2015; Swain *et al.* 2015; Pereiro *et al.* 2016; Velazquez *et al.* 2017; Xiang *et al.* 2017; Yang *et al.* 2017), although the induction of IFN- γ and IFN- γ rel is tissue-specific or developmental stage-specific in some species of fish (Aggad *et al.* 2010; Lopez-Munoz *et al.* 2010; Velazquez *et al.* 2017). Meanwhile, type II IFNs are able to promote antiviral genes, including Mx, ISG15, RSAD2 etc., and can inhibit viral replication *in vitro* (Lopez-Munoz *et al.* 2009; Sun *et al.* 2011a; Shibasaki *et al.* 2014; Kwak *et al.* 2017; Xiang *et al.* 2017), despite that a minor inhibitory effect was observed for Atlantic salmon IFN- γ on the replication of virus in TO cell (Xu *et al.* 2010a). Furthermore, the antiviral activity of IFN- γ is partially dependent on the induction of type I IFN in TO cells (Sun *et al.* 2011a). These data suggest that teleost type II IFNs are involved in antiviral immune responses. However, the *in vivo* overexpression of IFN- γ or IFN- γ rel could not protect zebrafish larvae against SVCV infection but surprisingly led to a higher mortality rate than that infected with SVCV only, although the overexpression of these two genes significantly increased the expression of antiviral genes (Lopez-Munoz *et al.* 2011). A possible explanation is that type II IFNs activate phagocytes and induce excessive inflammation due to strongly up-regulated expression of pro-inflammatory genes (Lopez-Munoz *et al.* 2011). In fact, it is still unclear whether viral proliferation is repressed *in vivo* by type II IFNs. However, some results seem to indicate viral inhibition led by IFN- γ *in vivo*: cumulative mortality of Japanese flounder was decreased when the fish were infected with recombinant VHSV which expressed the flounder IFN- γ comparing to the control

groups by other recombinant or wild-type VHSVs (Kwak *et al.* 2017).

The role of type II IFNs in antibacterial immunity

Teleost type II IFNs may have antibacterial activity. First, IFN- γ and IFN- γ rel are capable of inducing the expression of nitric oxide synthase (iNOS) gene, enhancing nitric oxide (NO) production in phagocytes and increasing phagocytic activity of immune-related cells (Grayfer & Belosevic 2009a,b; Arts *et al.* 2010; Grayfer *et al.* 2010, 2018; Yang *et al.* 2013, 2017; Biswas *et al.* 2016; Peng *et al.* 2016, 2018; Wiegertjes *et al.* 2016). NO is essential for antimicrobial activity of macrophages in mammals (MacMicking *et al.* 1997). Second, IFN- γ and IFN- γ rel are able to induce cytochrome b-245 beta chain (CYBB, also known as gp91^{phox} and NOX2, a subunit of the NADPH oxidase), neutrophil cytosolic factor 1 (NCF1, also known as p47^{phox} and NOXO2) and respiratory burst, including the production of reactive oxygen species (ROS), in phagocytes (Zou *et al.* 2005; Grayfer & Belosevic 2009a,b; Grayfer *et al.* 2010, 2018; Wiegertjes *et al.* 2016; Yang *et al.* 2017; Peng *et al.* 2018), which are involved in killing bacteria in mammals (Schroder *et al.* 2004). Third, the knockdown of type II IFNs (IFN- γ 1/IFN- γ rel and IFN- γ 2/IFN- γ) and their receptors, including CRFB13, CRFB17 and CRFB6, resulted in significantly increased susceptibility to the infection of *Yersinia ruckeri* in zebrafish larvae (Sieger *et al.* 2009; Aggad *et al.* 2010). On the other hand, excessive IFN- γ and IFN- γ rel in zebrafish may lead to higher mortality rate than in wild-type fish, and it seems possible that the expression level of IFN- γ and IFN- γ rel is strictly controlled in response to bacteria (Sieger *et al.* 2009), although both IFN- γ and IFN- γ rel are up-regulated by a variety of bacteria (Raida & Buchmann 2008; Furnes *et al.* 2009; Metzger *et al.* 2010; Gjessing *et al.* 2011; Evenhuis & Cleveland 2012; Hodgkinson *et al.* 2012; Kondo *et al.* 2014; Chen *et al.* 2015c; Swain *et al.* 2015; Pereiro *et al.* 2016; Takizawa *et al.* 2016; Velazquez *et al.* 2017). Moreover, intraperitoneal injection of recombinant IFN- γ 2 (IFN- γ) and IFN- γ 1 (IFN- γ rel) in pufferfish could not provide protection against the infection of *Vibrio parahaemolyticus*; simultaneously, recombinant IFN- γ 1 (IFN- γ rel) reduced survival rate and induced stronger inflammatory response than IFN- γ 2 (IFN- γ) (Peng *et al.* 2016). Nevertheless, the survival rate of Japanese flounder with recombinant IFN- γ injected intraperitoneally was increased when the fish was infected with *Edwardsiella tarda*, with the expression of inflammatory factors up-regulated remarkably by recombinant IFN- γ (Jung *et al.* 2012). Similarly, recombinant ginbuna crucian carp IFN- γ also protected fish against *Nocardia seriolae* infection (Nayak *et al.* 2014). The contradictory results may be impossible to explain at present, but may be associated with IFN- γ dose or different pathogens, as *E. tarda*

and *N. seriolae* are intracellular bacteria and are able to replicate in fish cells (Tan *et al.* 2005; Nayak *et al.* 2014; Xie *et al.* 2014; Matsumoto *et al.* 2017). However, *in vitro* pre-treatment of monocytes/macrophages with recombinant IFN- γ enhanced the invasion of another intracellular bacterium (*Francisella noatunensis* subsp. *noatunensis*) in Atlantic cod, which may have been resulted from macrophage activation as an early step of host defense, because the internalization of *F. noatunensis* subsp. *noatunensis* is likely mediated by phagocytosis (Bakkemo *et al.* 2016). Furthermore, Atlantic salmon IFN- γ plays a role in the TLR9-mediated innate immune response *in vitro* (Skjaeveland *et al.* 2008; Iliev *et al.* 2013). It is thus indicated that teleost type II IFNs may exhibit some functional complexity in antibacterial immunity.

The role of type II IFNs in cell proliferation and development

Teleost type II IFN signalling may have an effect on cell proliferation and development. In zebrafish embryos, type II IFN signalling is able to enhance the formation of haematopoietic stem and progenitor cells (HSPCs) in the aorta-gonad-mesonephros (AGM) region (Li *et al.* 2014b). Indeed, IFN- γ signalling is involved in the development of haematopoietic stem cells (HSCs), and endothelial-to-HSC conversion is positively regulated by IFN- γ and CRFB17 (IFNGR1-1), whose expression is controlled by Notch signalling and blood flow in the AGM region (Sawamiphak *et al.* 2014). In fact, the transcription of type II IFN receptor genes, including CRFB13 (IFNGR1-2) and CRFB17, is also maintained by RNA polymerase II (Pol II) pausing, whose deficiency may result in the defective HSPC formation and decreased chromatin accessibility on promoters of type II IFN receptor genes, thus suggesting that Pol II pausing is required for HSC emergence mediated by type II IFN signalling in zebrafish embryos (Yang *et al.* 2016a). Furthermore, IFN- γ signalling participated in developmental biliary defects in zebrafish (Matthews *et al.* 2011; Cui *et al.* 2013). Additionally, the expression of medaka IFN- γ 2b (one of IFN- γ 2 splicing forms) was modulated by steroid and was detected only and predominantly in female germ cells whose proliferation is possibly associated with IFN- γ 2b (Mohapatra *et al.* 2015).

Other functions of type II IFNs

As in mammals, type II IFNs in teleost are multi-functional molecules that play roles in many other biological processes, including allograft rejection, antigen presentation etc. For instance, the treatment of recombinant IFN- γ rel1 (an isoform of IFN- γ rel) enhances the scale allograft rejection and increases the number of CD4⁺, CD8⁺ and sIgM⁺ cells in grafted scales of gibel carp (Shibasaki *et al.* 2016). Teleost type II IFNs can strongly induce antigen-presentation-associated genes, including proteasome

subunit beta 9 (PSMB9, also known as LMP2), transporter 2 (ATP binding cassette subfamily B member, TAP2), beta-2-microglobulin (B2M), major histocompatibility complex (MHC) molecules etc. (Zou *et al.* 2005; Martin *et al.* 2007a,b,c; Castro *et al.* 2008; Lopez-Munoz *et al.* 2011). However, type II IFN mediated antigen presentation is not well understood in fish. Moreover, the expression of IFN- γ was reduced in those zebrafish less responsive to social stimuli (Kirsten *et al.* 2018), and zebrafish IFN- γ promotes anti-pathogen response in brain, which may be associated with social/aggregation behaviour (Filiano *et al.* 2016).

MicroRNAs (miRNAs) are small non-coding RNAs that target messenger RNA (mRNA) to regulate post-transcription of genes to impact many biological processes (Ambros 2004; Bushati & Cohen 2007; Krol *et al.* 2010). It has been reported that teleost miRNAs are regulated by type II IFNs and are involved in biological functions mediated by type II IFNs. In green spotted pufferfish, the expression of over 1500 miRNAs has changed in spleen cells after the treatment of recombinant type II IFNs (Yi *et al.* 2014). Orange-spotted grouper IFN- γ 2 (IFN- γ) is likely to be involved in anti-inflammatory response through the induction of microRNA-146a (miR-146a) which possibly inhibits TRAF6 (Peng *et al.* 2018). In addition, the deletion of miR-142-3p in zebrafish causes impaired neutrophil development mediated by the abnormally activated IFN- γ signalling (Fan *et al.* 2014). Zebrafish IFN- γ is likely to inhibit the over-inflammation induced by miR-223 deficiency, although this requires more experimental evidence (Zhou *et al.* 2018).

Conclusions and future perspectives

From an evolutionary point of view, fish possess a unique type I IFN repertoire, which contains only intron-containing type I IFNs (Zou & Secombes 2011; Langevin *et al.* 2013a; Boudinot *et al.* 2016), and fish may represent an important model to study the origin and evolution of type I IFNs in vertebrates. Different from other lineages of vertebrates, teleost fish have two members of type II IFNs, i.e. IFN- γ or IFN- γ 2 and IFN- γ rel or IFN- γ 1, which use different receptor systems to activate JAK-STAT signalling. Although the understanding on the fish type I IFN and type II systems has been significantly increased in recent years, some significant issues remain to be resolved.

- (1) RNA virus-induced type I IFN response in RIG-I-absent fish species. As RIG-I seems to be lost in certain groups of fish species, such as in Acanthopterygii (Zou *et al.* 2009; Chen *et al.* 2017b), and also RIG-I is absent in chicken, which is probably a key factor for the high susceptibility to RNA viruses, such as zoonotic influenza viruses (Barber *et al.* 2010; Santhakumar *et al.* 2017), whether RIG-I-absent fish species are more susceptible to RNA viruses, and whether these species of

fish may have some undiscovered mechanisms to compensate the absence of RIG-I remains to be investigated. Hence, one important future direction is to gain a deeper understanding of type I IFN-inducing mechanism in RIG-I-absent fish species upon RNA virus infection.

- (2) The cell type-specific expression of type I IFNs in fish. In mammals, type I IFNs are mainly produced by pDCs through TLR7/TLR8/TLR9–MyD88 signalling after viral infection (Liu 2005). Whether fish possess a cell type specific expression pattern similar to mammalian pDCs remains as a mystery. It is necessary to analyse cell surface markers of type I IFN-producing cell populations in fish, and to elucidate the molecular mechanism in which these cell populations can produce large amount of type I IFNs.
- (3) The functional property of IFN δ in fish. Unlike other IFN subtypes, IFN δ has been proven to have antiviral activity in some fish species, but not in some other fish species, such as zebrafish, Atlantic salmon and mandarin fish (Aggad *et al.* 2009; Svingerud *et al.* 2012; Laghari *et al.* 2018). The factors or mechanisms related with the functional or dysfunctional IFN δ in different species of fish is of value for further research.
- (4) The ligand-receptor models for type II IFNs in fish. There are three possible models for IFN- γ (IFN- γ 2), homodimer of IFN- γ binds to the receptor complex with $2 \times$ IFN- γ R2 (CRFB6) subunits + $2 \times$ IFN-

γ R1-2 (CRFB13) subunits, $2 \times$ IFN- γ R2 + $2 \times$ IFN- γ R1-1 (CRFB17) and $2 \times$ IFN- γ R2 + heterodimer of IFN- γ R1-1 and IFN- γ R1-2 (Fig. 3). It would be necessary to investigate if these ligand-receptor relationships are species-specific or coexist in some cell-types of a species. On the other hand, for IFN- γ rel (IFN- γ 1), four cases of ligand-receptor complex may be possibly included as the followings: homodimer of IFN- γ rel binds to $2 \times$ unknown molecules + $2 \times$ IFN- γ R1-1 molecules or binds to homodimer of IFN- γ R1-1 only (Aggad *et al.* 2010); monomer of IFN- γ rel binds to homodimer or monomer of IFN- γ R1-1 (Fig. 3). All these receptor models need to be experimentally tested or verified in future (Aggad *et al.* 2010).

- (5) The possible signalling and functional difference mediated by fish IFN- γ and IFN- γ rel. It has been reported that the signalling pathway or function of IFN- γ and IFN- γ rel are quite different (Grayfer *et al.* 2010; Lopez-Munoz *et al.* 2011; Peng *et al.* 2016; Shibasaki *et al.* 2016), and that IFN- γ and IFN- γ rel may target possibly different cell-types. It would be a novel breakthrough to understand the functional similarity or divergence, if any, of the two members of type II IFNs in teleost.

The most challenging question would be “Does type III IFN exist in fish”? To date, type I and II IFNs have been identified in elephant shark, a cartilaginous fish, but not type III IFN (Venkatesh *et al.* 2014). In fact, it is reported that the elephant shark contains IFNL1 gene, which




Figure 3 Signalling and ligand-receptor models for type II IFNs in fish. The homodimer of IFN- γ likely binds to the receptor complex with $2 \times$ IFN- γ R2 (CRFB6) subunits plus $2 \times$ IFN- γ R1-2 (CRFB13) subunits, $2 \times$ IFN- γ R2 plus $2 \times$ IFN- γ R1-1 (CRFB17) or $2 \times$ IFN- γ R2 plus heterodimer of IFN- γ R1-1 and IFN- γ R1-2, while the receptors of IFN- γ rel contain at least IFN- γ R1-1 (CRFB17). IFN- γ and IFN- γ rel regulate the expression of ISGs (IRF1, PSMB9 and etc.) via JAK-STAT signalling. In addition, IRF1, as the secondary transcription factor, is likely involved in signal transduction mediated by type II IFNs.

encodes the specific receptor chain of type III IFNs (Hemann *et al.* 2017), with conserved syntenic in comparison with tetrapods (Venkatesh *et al.* 2014; Secombes & Zou 2017). It is worthwhile to search for type III IFN in cartilaginous fish on the basis of ligand-receptor coevolution hypothesis (Moyle *et al.* 1994).

Acknowledgements

This work was financially supported by grants (Nos. 31320103913 and 31702369) from the National Natural Science Foundation of China (NSFC), and also by China Agriculture Research System (CARS-46), and by a special talent programme “One Thing One Decision (Yishi Yiyi)” and “First class fishery discipline” programme in Shandong Province, China.

References

- Aggad D, Mazel M, Boudinot P, Mogensen KE, Hamming OJ, Hartmann R *et al.* (2009) The two groups of zebrafish virus-induced interferons signal via distinct receptors with specific and shared chains. *Journal of Immunology* **183**: 3924–3931.
- Aggad D, Stein C, Sieger D, Mazel M, Boudinot P, Herbolme P *et al.* (2010) *In vivo* analysis of Ifn- γ 1 and Ifn- γ 2 signaling in zebrafish. *Journal of Immunology* **185**: 6774–6782.
- Ai K, Luo K, Xia L, Gao W, Hu W, Qi Z *et al.* (2018) Functional characterization of interferon regulatory factor 5 and its role in the innate antiviral immune response. *Fish and Shellfish Immunology* **72**: 31–36.
- Altmann SM, Mellon MT, Distel DL, Kim CH (2003) Molecular and functional analysis of an interferon gene from the zebrafish, *Danio rerio*. *Journal of Virology* **77**: 1992–2002.
- Ambros V (2004) The functions of animal microRNAs. *Nature* **431**: 350–355.
- Aoki T, Hikima J, Hwang SD, Jung TS (2013) Innate immunity of finfish: primordial conservation and function of viral RNA sensors in teleosts. *Fish and Shellfish Immunology* **35**: 1689–1702.
- Arts JA, Tijhaar EJ, Chadzinska M, Savelkoul HF, Verburg-van Kemenade BM (2010) Functional analysis of carp interferon- γ : evolutionary conservation of classical phagocyte activation. *Fish and Shellfish Immunology* **29**: 793–802.
- Bach EA, Aguet M, Schreiber RD (1997) The IFN γ receptor: a paradigm for cytokine receptor signaling. *Annual Review of Immunology* **15**: 563–591.
- Bakkemo KR, Mikkelsen H, Johansen A, Robertsen B, Seppola M (2016) *Francisella noatunensis* subsp. *noatunensis* invades, survives and replicates in Atlantic cod cells. *Diseases of Aquatic Organisms* **121**: 149–159.
- Barber MR, Aldridge JR Jr, Webster RG, Magor KE (2010) Association of RIG-I with innate immunity of ducks to influenza. *Proceedings of the National Academy of Sciences of the United States of America* **107**: 5913–5918.
- Bergan V, Kileng O, Sun B, Robertsen B (2010) Regulation and function of interferon regulatory factors of Atlantic salmon. *Molecular Immunology* **47**: 2005–2014.
- Biacchesi S, LeBerre M, Lamoureux A, Louise Y, Lauret E, Boudinot P *et al.* (2009) Mitochondrial antiviral signaling protein plays a major role in induction of the fish innate immune response against RNA and DNA viruses. *Journal of Virology* **83**: 7815–7827.
- Biacchesi S, Merour E, Lamoureux A, Bernard J, Bremont M (2012) Both STING and MAVS fish orthologs contribute to the induction of interferon mediated by RIG-I. *PLoS ONE* **7**: e47737.
- Biswas G, Nagamine R, Hikima J, Sakai M, Kono T (2016) Inductive immune responses in the Japanese pufferfish (*Takifugu rubripes*) treated with recombinant IFN- γ , IFN- γ marel, IL-4/13A and IL-4/13B. *International Immunopharmacology* **31**: 50–56.
- Boehm U, Klamp T, Groot M, Howard JC (1997) Cellular responses to interferon- γ . *Annual Review of Immunology* **15**: 749–795.
- Boudinot P, Langevin C, Secombes CJ, Levraud JP (2016) The peculiar characteristics of fish type I interferons. *Viruses* **8**: 298.
- Bruns AM, Horvath CM (2014) Antiviral RNA recognition and assembly by RLR family innate immune sensors. *Cytokine and Growth Factor Reviews* **25**: 507–512.
- Bushati N, Cohen SM (2007) MicroRNA functions. *Annual Review of Cell and Developmental Biology* **23**: 175–205.
- Caipang CM, Hirono I, Aoki T (2005) Induction of antiviral state in fish cells by Japanese flounder, *Paralichthys olivaceus*, interferon regulatory factor-1. *Fish and Shellfish Immunology* **19**: 79–91.
- Cao Y, Xu L, LaPatra SE, Zhao J, Liu M, Liu H *et al.* (2016) The kinetics and protection of the antiviral state induced by recombinant iIFN1 α in rainbow trout against infectious hematopoietic necrosis virus. *Molecular Immunology* **76**: 55–61.
- Casani D, Randelli E, Costantini S, Facchiano AM, Zou J, Martin S *et al.* (2009) Molecular characterisation and structural analysis of an interferon homologue in sea bass (*Dicentrarchus labrax* L.). *Molecular Immunology* **46**: 943–952.
- Castro R, Martin SA, Bird S, Lamas J, Secombes CJ (2008) Characterisation of gamma-interferon responsive promoters in fish. *Molecular Immunology* **45**: 3454–3462.
- Chang M, Nie P, Collet B, Secombes CJ, Zou J (2009) Identification of an additional two-cysteine containing type I interferon in rainbow trout *Oncorhynchus mykiss* provides evidence of a major gene duplication event within this gene family in teleosts. *Immunogenetics* **61**: 315–325.
- Chang M, Collet B, Nie P, Lester K, Campbell S, Secombes CJ *et al.* (2011) Expression and functional characterization of the RIG-I-like receptors MDA5 and LGP2 in rainbow trout (*Oncorhynchus mykiss*). *Journal of Virology* **85**: 8403–8412.
- Chang MX, Zou J, Nie P, Huang B, Yu Z, Collet B *et al.* (2013) Intracellular interferons in fish: a unique means to combat viral infection. *PLoS Pathogens* **9**: e1003736.

- Chang CJ, Robertsen C, Sun B, Robertsen B (2014) Protection of Atlantic salmon against virus infection by intramuscular injection of IFN γ expression plasmid. *Vaccine* **32**: 4695–4702.
- Chang CJ, Sun B, Robertsen B (2015) Adjuvant activity of fish type I interferon shown in a virus DNA vaccination model. *Vaccine* **33**: 2442–2448.
- Chang CJ, Jenssen I, Robertsen B (2016) Protection of Atlantic salmon against salmonid alphavirus infection by type I interferons IFN α , IFN β and IFN γ . *Fish and Shellfish Immunology* **57**: 35–40.
- Chen WQ, Xu QQ, Chang MX, Zou J, Secombes CJ, Peng KM *et al.* (2010) Molecular characterization and expression analysis of the IFN- γ related gene (IFN- γ rel) in grass carp *Ctenopharyngodon idella*. *Veterinary Immunology and Immunopathology* **134**: 199–207.
- Chen SN, Huang B, Zhang XW, Li Y, Zhao LJ, Li N *et al.* (2013) IFN- γ and its receptors in a reptile reveal the evolutionary conservation of type II IFNs in vertebrates. *Developmental and Comparative Immunology* **41**: 587–596.
- Chen YM, Kuo CE, Chen GR, Kao YT, Zou J, Secombes CJ *et al.* (2014) Functional analysis of an orange-spotted grouper (*Epinephelus coioides*) interferon gene and characterisation of its expression in response to nodavirus infection. *Developmental and Comparative Immunology* **46**: 117–128.
- Chen C, Zhang YB, Gui JF (2015a) Expression characterization, genomic structure and function analysis of fish ubiquitin-specific protease 18 (USP18) genes. *Developmental and Comparative Immunology* **52**: 112–122.
- Chen H, Liu W, Wang B, Mao H, Sun Z, Hou Q *et al.* (2015b) Cloning, identification of the two cytokine receptor family B subunits CRFB1 and CRFB5 from grass carp (*Ctenopharyngodon idella*). *Fish and Shellfish Immunology* **45**: 211–220.
- Chen RN, Su YQ, Wang J, Liu M, Qiao Y, Mao Y *et al.* (2015c) Molecular characterization and expression analysis of interferon- γ in the large yellow croaker *Larimichthys crocea*. *Fish and Shellfish Immunology* **46**: 596–602.
- Chen WQ, Hu YW, Zou PF, Ren SS, Nie P, Chang MX (2015d) MAVS splicing variants contribute to the induction of interferon and interferon-stimulated genes mediated by RIG-I-like receptors. *Developmental and Comparative Immunology* **49**: 19–30.
- Chen X, Yang C, Su J, Rao Y, Gu T (2015e) LGP2 plays extensive roles in modulating innate immune responses in *Ctenopharyngodon idella* kidney (CIK) cells. *Developmental and Comparative Immunology* **49**: 138–148.
- Chen Q, Sun L, Chen ZJ (2016) Regulation and function of the cGAS-STING pathway of cytosolic DNA sensing. *Nature Immunology* **17**: 1142–1149.
- Chen H, Xiao J, Li J, Liu J, Wang C, Feng C *et al.* (2017a) TRAF2 of black carp upregulates MAVS-mediated antiviral signaling during innate immune response. *Fish and Shellfish Immunology* **71**: 1–9.
- Chen SN, Zou PF, Nie P (2017b) Retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs) in fish: current knowledge and future perspectives. *Immunology* **151**: 16–25.
- Cheng CH, Chou CM, Chu CY, Chen GD, Lien HW, Hwang PP *et al.* (2014) Differential regulation of *Tetraodon nigroviridis* Mx gene promoter activity by constitutively-active forms of STAT1, STAT2, and IRF9. *Fish and Shellfish Immunology* **38**: 230–243.
- Collet B (2014) Innate immune responses of salmonid fish to viral infections. *Developmental and Comparative Immunology* **43**: 160–173.
- Collins C, Ganne G, Collet B (2014) Isolation and activity of the promoters for STAT1 and 2 in Atlantic salmon *Salmo salar*. *Fish and Shellfish Immunology* **40**: 644–647.
- Collins C, Lorenzen N, Collet B (2018) DNA vaccination for fin-fish aquaculture. *Fish and Shellfish Immunology* **49**: 138–148.
- Cui H, Yan Y, Wei J, Huang X, Huang Y, Ouyang Z *et al.* (2011) Identification and functional characterization of an interferon regulatory factor 7-like (IRF7-like) gene from orange-spotted grouper, *Epinephelus coioides*. *Developmental and Comparative Immunology* **35**: 672–684.
- Cui S, Eaucraire SF, Matthews RP (2013) Interferon- γ directly mediates developmental biliary defects. *Zebrafish* **10**: 177–183.
- Ding Y, Ao J, Huang X, Chen X (2016) Identification of two subgroups of type I IFNs in perciforme fish large yellow croaker *Larimichthys crocea* provides novel insights into function and regulation of fish type I IFNs. *Frontiers in Immunology* **7**: 343.
- Ding Y, Guan Y, Ao J, Chen X (2017) Induction of type I interferons in response to bacterial stimuli in large yellow croaker *Larimichthys crocea*. *Fish and Shellfish Immunology* **62**: 349–355.
- Du F, Su J, Huang R, Liao L, Zhu Z, Wang Y (2013) Cloning and preliminary functional studies of the JAM-A gene in grass carp (*Ctenopharyngodon idellus*). *Fish and Shellfish Immunology* **34**: 1476–1484.
- Du J, Zhang D, Zhang W, Ouyang G, Wang J, Liu X *et al.* (2015) pVHL negatively regulates antiviral signaling by targeting MAVS for proteasomal degradation. *Journal of Immunology* **195**: 1782–1790.
- Ealick SE, Cook WJ, Vijay-Kumar S, Carson M, Nagabhushan TL, Trotta PP *et al.* (1991) Three-dimensional structure of recombinant human interferon- γ . *Science* **252**: 698–702.
- Evenhuis JP, Cleveland BM (2012) Modulation of rainbow trout (*Oncorhynchus mykiss*) intestinal immune gene expression following bacterial challenge. *Veterinary Immunology and Immunopathology* **146**: 8–17.
- Evensen O, Leong JA (2013) DNA vaccines against viral diseases of farmed fish. *Fish and Shellfish Immunology* **35**: 1751–1758.
- Fan HB, Liu YJ, Wang L, Du TT, Dong M, Gao L *et al.* (2014) miR-142-3p acts as an essential modulator of neutrophil development in zebrafish. *Blood* **124**: 1320–1330.
- Feng X, Su J, Yang C, Yan N, Rao Y, Chen X (2014a) Molecular characterizations of grass carp (*Ctenopharyngodon idella*) TBK1 gene and its roles in regulating IFN-I pathway. *Developmental and Comparative Immunology* **45**: 278–290.
- Feng X, Yang C, Zhang Y, Peng L, Chen X, Rao Y *et al.* (2014b) Identification, characterization and immunological response

- analysis of stimulator of interferon gene (STING) from grass carp *Ctenopharyngodon idella*. *Developmental and Comparative Immunology* **45**: 163–176.
- Feng H, Zhang YB, Zhang QM, Li Z, Zhang QY, Gui JF (2015a) Zebrafish IRF1 regulates IFN antiviral response through binding to IFN ϕ 1 and IFN ϕ 3 promoters downstream of MyD88 signaling. *Journal of Immunology* **194**: 1225–1238.
- Feng X, Zhang Y, Yang C, Liao L, Wang Y, Su J (2015b) Functional characterizations of IPS-1 in CIK cells: potential roles in regulating IFN-I response dependent on IRF7 but not IRF3. *Developmental and Comparative Immunology* **53**: 23–32.
- Feng H, Zhang QM, Zhang YB, Li Z, Zhang J, Xiong YW *et al.* (2016) Zebrafish IRF1, IRF3, and IRF7 differentially regulate IFN ϕ 1 and IFN ϕ 3 expression through assembly of homo- or heteroprotein complexes. *Journal of Immunology* **197**: 1893–1904.
- Filiano AJ, Xu Y, Tustison NJ, Marsh RL, Baker W, Smirnov I *et al.* (2016) Unexpected role of interferon-gamma in regulating neuronal connectivity and social behaviour. *Nature* **535**: 425–429.
- Funami K, Matsumoto M, Oshiumi H, Akazawa T, Yamamoto A, Seta T (2004) The cytoplasmic ‘linker region’ in Toll-like receptor 3 controls receptor localization and signaling. *International Immunology* **16**: 1143–1154.
- Furnes C, Seppola M, Robertsen B (2009) Molecular characterization and expression analysis of interferon gamma in Atlantic cod (*Gadus morhua*). *Fish and Shellfish Immunology* **26**: 285–292.
- Gabor KA, Charette JR, Pietraszewski MJ, Wingfield DJ, Shim JS, Millard PJ *et al.* (2015) A DN-mda5 transgenic zebrafish model demonstrates that Mda5 plays an important role in snakehead rhabdovirus resistance. *Developmental and Comparative Immunology* **51**: 298–304.
- Gan Z, Chen SN, Huang B, Hou J, Nie P (2017) Intronless and intron-containing type I IFN genes coexist in amphibian *Xenopus tropicalis*: insights into the origin and evolution of type I IFNs in vertebrates. *Developmental and Comparative Immunology* **67**: 166–176.
- Gan Z, Yang YC, Chen SN, Hou J, Laghari ZA, Huang B *et al.* (2018) Unique composition of intronless and intron-containing type I IFNs in the Tibetan frog *Nanorana parkeri* provides new evidence to support independent retroposition hypothesis for type I IFN genes in amphibians. *Journal of Immunology* **201**: 3329–3342.
- Gao Q, Nie P, Thompson KD, Adams A, Wang T, Secombes CJ *et al.* (2009) The search for the IFN-gamma receptor in fish: functional and expression analysis of putative binding and signalling chains in rainbow trout *Oncorhynchus mykiss*. *Developmental and Comparative Immunology* **33**: 920–931.
- Gao FX, Lu WJ, Wang Y, Zhang QY, Zhang YB, Mou CY *et al.* (2018) Differential expression and functional diversification of diverse immunoglobulin domain-containing protein (DICP) family in three gynogenetic clones of gibel carp. *Developmental and Comparative Immunology* **84**: 396–407.
- Ge R, Zhou Y, Peng R, Wang R, Li M, Zhang Y *et al.* (2015) Conservation of the STING-mediated cytosolic DNA sensing pathway in zebrafish. *Journal of Virology* **89**: 7696–7706.
- Gjessing MC, Inami M, Weli SC, Ellingsen T, Falk K, Koppang EO *et al.* (2011) Presence and interaction of inflammatory cells in the spleen of Atlantic cod, *Gadus morhua* L., infected with *Francisella noatunensis*. *Journal of Fish Diseases* **34**: 687–699.
- Gonzalez SF, Chatziandreou N, Nielsen ME, Li W, Rogers J, Taylor R *et al.* (2007) Cutaneous immune responses in the common carp detected using transcript analysis. *Molecular Immunology* **44**: 1664–1679.
- Grayfer L, Belosevic M (2009a) Molecular characterization, expression and functional analysis of goldfish (*Carassius auratus* L.) interferon gamma. *Developmental and Comparative Immunology* **33**: 235–246.
- Grayfer L, Belosevic M (2009b) Molecular characterization of novel interferon gamma receptor 1 isoforms in zebrafish (*Danio rerio*) and goldfish (*Carassius auratus* L.). *Molecular Immunology* **46**: 3050–3059.
- Grayfer L, Garcia EG, Belosevic M (2010) Comparison of macrophage antimicrobial responses induced by type II interferons of the goldfish (*Carassius auratus* L.). *Journal of Biological Chemistry* **285**: 23537–23547.
- Grayfer L, Kerimoglu B, Yaparla A, Hodgkinson JW, Xie J, Belosevic M (2018) Mechanisms of fish macrophage antimicrobial immunity. *Frontiers in Immunology* **9**: 1105.
- Green DS, Young HA, Valencia JC (2017) Current prospects of type II interferon gamma signaling and autoimmunity. *Journal of Biological Chemistry* **292**: 13925–13933.
- Gu M, Lin G, Lai Q, Zhong B, Liu Y, Mi Y *et al.* (2015) *Ctenopharyngodon idella* IRF2 plays an antagonistic role to IRF1 in transcriptional regulation of IFN and ISG genes. *Developmental and Comparative Immunology* **49**: 103–112.
- Guo CJ, Zhang YF, Yang LS, Yang XB, Wu YY, Liu D *et al.* (2009) The JAK and STAT family members of the mandarin fish *Siniperca chuatsi*: molecular cloning, tissues distribution and immunobiological activity. *Fish and Shellfish Immunology* **27**: 349–359.
- Hamming OJ, Lutfalla G, Levraud JP, Hartmann R (2011) Crystal structure of zebrafish interferons I and II reveals conservation of type I interferon structure in vertebrates. *Journal of Virology* **85**: 8181–8187.
- Hemann EA, Gale M Jr, Savan R (2017) Interferon lambda genetics and biology in regulation of viral control. *Frontiers in Immunology* **8**: 1707.
- Hertzog PJ, Williams BR (2013) Fine tuning type I interferon responses. *Cytokine and Growth Factor Reviews* **24**: 217–225.
- Hodgkinson JW, Ge JQ, Grayfer L, Stafford J, Belosevic M (2012) Analysis of the immune response in infections of the goldfish (*Carassius auratus* L.) with *Mycobacterium marinum*. *Developmental and Comparative Immunology* **38**: 456–465.
- Holland JW, Bird S, Williamson B, Woudstra C, Mustafa A, Wang T *et al.* (2008) Molecular characterization of IRF3 and IRF7 in rainbow trout, *Oncorhynchus mykiss*: functional

- analysis and transcriptional modulation. *Molecular Immunology* **46**: 269–285.
- Honda K, Yanai H, Negishi H, Asagiri M, Sato M, Mizutani T *et al.* (2005) IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* **434**: 772–777.
- Hou Q, Gong R, Liu X, Mao H, Xu X, Liu D *et al.* (2017) Poly I: C facilitates the phosphorylation of Ctenopharyngodon idellus type I IFN receptor subunits and JAK kinase. *Fish and Shellfish Immunology* **60**: 13–20.
- Hu G, Yin X, Lou H, Xia J, Dong X, Zhang J *et al.* (2011) Interferon regulatory factor 3 (IRF-3) in Japanese flounder, *Paralichthys olivaceus*: sequencing, limited tissue distribution, inducible expression and induction of fish type I interferon promoter. *Developmental and Comparative Immunology* **35**: 164–173.
- Hu Y, Yoshikawa T, Chung S, Hirono I, Kondo H (2017) Identification of 2 novel type I IFN genes in Japanese flounder, *Paralichthys olivaceus*. *Fish and Shellfish Immunology* **67**: 7–10.
- Hu Y, Huang Y, Liu J, Zhang J, Qin Q, Huang X (2018a) TBK1 from orange-spotted grouper exerts antiviral activity against fish viruses and regulates interferon response. *Fish and Shellfish Immunology* **73**: 92–99.
- Hu YW, Zhang J, Wu XM, Cao L, Nie P, Chang MX (2018b) TANK-binding kinase 1 (TBK1) isoforms negatively regulate type I interferon induction by inhibiting TBK1-IRF3 interaction and IRF3 phosphorylation. *Frontiers in Immunology* **9**: 84.
- Huang B, Qi ZT, Xu Z, Nie P (2010) Global characterization of interferon regulatory factor (IRF) genes in vertebrates: glimpse of the diversification in evolution. *BMC Immunology* **11**: 22.
- Huang Y, Ouyang Z, Wang W, Yu Y, Li P, Zhou S *et al.* (2015a) Antiviral role of grouper STING against iridovirus infection. *Fish and Shellfish Immunology* **47**: 157–167.
- Huang Z, Chen S, Liu J, Xiao J, Yan J, Feng H (2015b) IFN α of black carp is an antiviral cytokine modified with N-linked glycosylation. *Fish and Shellfish Immunology* **46**: 477–485.
- Huang Y, Yang M, Yu Y, Yang Y, Zhou L, Huang X *et al.* (2016a) Grouper TRIM13 exerts negative regulation of antiviral immune response against nodavirus. *Fish and Shellfish Immunology* **55**: 106–115.
- Huang Y, Yu Y, Yang Y, Yang M, Zhou L, Huang X *et al.* (2016b) Antiviral function of grouper MDA5 against iridovirus and nodavirus. *Fish and Shellfish Immunology* **54**: 188–196.
- Huang CJ, Chou CM, Lien HW, Chu CY, Ho JY, Wu Y *et al.* (2017a) IRF9-Stat2 fusion protein as an innate immune inducer to activate Mx and interferon-stimulated gene expression in zebrafish larvae. *Marine Biotechnology* **19**: 310–319.
- Huang Y, Zhang J, Liu J, Hu Y, Ni S, Yang Y *et al.* (2017b) Fish TRIM35 negatively regulates the interferon signaling pathway in response to grouper nodavirus infection. *Fish and Shellfish Immunology* **69**: 142–152.
- Huang B, Wang ZX, Liang Y, Zhai SW, Huang WS, Nie P (2018a) Identification of four type I IFNs from Japanese eel with differential expression properties and Mx promoter inducibility. *Developmental and Comparative Immunology* **91**: 62–71.
- Huang Y, Zhang J, Ouyang Z, Liu J, Zhang Y, Hu Y *et al.* (2018b) Grouper MAVS functions as a crucial antiviral molecule against nervous necrosis virus infection. *Fish and Shellfish Immunology* **72**: 14–22.
- Igawa D, Sakai M, Savan R (2006) An unexpected discovery of two interferon gamma-like genes along with interleukin (IL)-22 and -26 from teleost: IL-22 and -26 genes have been described for the first time outside mammals. *Molecular Immunology* **43**: 999–1009.
- Ikushima H, Negishi H, Taniguchi T (2013) The IRF family transcription factors at the interface of innate and adaptive immune responses. *Cold Spring Harbor Symposia on Quantitative Biology* **78**: 105–116.
- Iliev DB, Sobhkhaz M, Fremmerlid K, Jorgensen JB (2011) MyD88 interacts with interferon regulatory factor (IRF) 3 and IRF7 in Atlantic salmon (*Salmo salar*): transgenic SsMyD88 modulates the IRF-induced type I interferon response and accumulates in aggregates. *Journal of Biological Chemistry* **286**: 42715–42724.
- Iliev DB, Skjaeveland I, Jorgensen JB (2013) CpG oligonucleotides bind TLR9 and RRM-containing proteins in Atlantic salmon (*Salmo salar*). *BMC Immunology* **14**: 12.
- Ivashkiv LB (2018) IFN γ : signalling, epigenetics and roles in immunity, metabolism, disease and cancer immunotherapy. *Nature Reviews Immunology* **18**: 545–558.
- Ji J, Rao Y, Wan Q, Liao Z, Su J (2018) Teleost-specific TLR19 localizes to endosome, recognizes dsRNA, recruits TRIF, triggers both IFN and NF- κ B pathways, and protects cells from grass carp reovirus infection. *Journal of Immunology* **200**: 573–585.
- Jia P, Jia K, Chen L, Le Y, Jin Y, Zhang J *et al.* (2016a) Identification and characterization of the melanoma differentiation-associated gene 5 in sea perch, *Lateolabrax japonicus*. *Developmental and Comparative Immunology* **61**: 161–168.
- Jia P, Jin Y, Chen L, Zhang J, Jia K, Yi M (2016b) Molecular characterization and expression analysis of mitochondrial antiviral signaling protein gene in sea perch, *Lateolabrax japonicus*. *Developmental and Comparative Immunology* **55**: 188–193.
- Jin Y, Zhou T, Li N, Liu S, Xu X, Pan Y *et al.* (2018) JAK and STAT members in channel catfish: identification, phylogenetic analysis and expression profiling after *Edwardsiella ictaluri* infection. *Developmental and Comparative Immunology* **81**: 334–341.
- Jung CY, Hikima J, Ohtani M, Jang HB, del Castillo CS, Nho SW *et al.* (2012) Recombinant interferon-gamma activates immune responses against *Edwardsiella tarda* infection in the olive flounder, *Paralichthys olivaceus*. *Fish and Shellfish Immunology* **33**: 197–203.
- Kasthuri SR, Wan Q, Whang I, Lim BS, Yeo SY, Choi CY *et al.* (2014) Functional characterization of the evolutionarily preserved mitochondrial antiviral signaling protein (MAVS) from rock bream, *Oplegnathus fasciatus*. *Fish and Shellfish Immunology* **40**: 399–406.
- Kawai T, Akira S (2010) The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nature Immunology* **11**: 373–384.

- Kawai T, Akira S (2011) Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity* **34**: 637–650.
- Kileng O, Bergan V, Workenhe ST, Robertsen B (2009) Structural and functional studies of an IRF-7-like gene from Atlantic salmon. *Developmental and Comparative Immunology* **33**: 18–27.
- Kirsten K, Fior D, Kreutz LC, Barcellos LJG (2018) First description of behavior and immune system relationship in fish. *Scientific Reports* **8**: 846.
- Kitao Y, Kono T, Korenaga H, Iizasa T, Nakamura K, Savan R *et al.* (2009) Characterization and expression analysis of type I interferon in common carp *Cyprinus carpio* L. *Molecular Immunology* **46**: 2548–2556.
- Kondo H, Kawana Y, Suzuki Y, Hirono I (2014) Comprehensive gene expression profiling in Japanese flounder kidney after injection with two different formalin-killed pathogenic bacteria. *Fish and Shellfish Immunology* **41**: 437–440.
- Kretschmer S, Lee-Kirsch MA (2017) Type I interferon-mediated autoinflammation and autoimmunity. *Current Opinion in Immunology* **49**: 96–102.
- Krishnan RKJ, Mushtaq Z, Shyam KU, Kurcheti PP (2018) Antiviral activity of transiently expressed mitochondrial antiviral signaling adapter, MAVS orthologue from Asian seabass. *Fish and Shellfish Immunology* **76**: 183–186.
- Krol J, Loedige I, Filipowicz W (2010) The widespread regulation of microRNA biogenesis, function and decay. *Nature Reviews Genetics* **11**: 597–610.
- Kung HC, Evensen O, Hong JR, Kuo CY, Tso CH, Ngou FH *et al.* (2014) Interferon regulatory factor-1 (IRF-1) is involved in the induction of phosphatidylserine receptor (PSR) in response to dsRNA virus infection and contributes to apoptotic cell clearance in CHSE-214 cell. *International Journal of Molecular Sciences* **15**: 19281–19306.
- Kuo HP, Chung CL, Hung YF, Lai YS, Chiou PP, Lu MW *et al.* (2016) Comparison of the responses of different recombinant fish type I interferons against betanodavirus infection in grouper. *Fish and Shellfish Immunology* **49**: 143–153.
- Kuznetsova IS, Thevasagayam NM, Sridatta PS, Komissarov AS, Saju JM, Ngoh SY *et al.* (2014) Primary analysis of repeat elements of the Asian seabass (*Lates calcarifer*) transcriptome and genome. *Frontiers in Genetics* **5**: 223.
- Kwak JS, Kim MS, Kim KH (2017) Generation of a recombinant viral hemorrhagic septicemia virus (VHSV) expressing olive flounder (*Paralichthys olivaceus*) interferon-gamma and its effects on type I interferon response and virulence. *Fish and Shellfish Immunology* **68**: 530–535.
- Laghari ZA, Chen SN, Li L, Huang B, Gan Z, Zhou Y *et al.* (2018) Functional, signalling and transcriptional differences of three distinct type I IFNs in a perciform fish, the mandarin fish *Siniperca chuatsi*. *Developmental and Comparative Immunology* **84**: 94–108.
- Lai Q, Lin G, Ma M, Huang S, Li W, Li D *et al.* (2013) IRF-1 acts as a positive regulator in the transcription of grass carp (*Ctenopharyngodon idella*) IFN gene. *Fish and Shellfish Immunology* **34**: 1432–1438.
- Langevin C, Aleksejeva E, Passoni G, Palha N, Levraud JP, Boudinot P (2013a) The antiviral innate immune response in fish: evolution and conservation of the IFN system. *Journal of Molecular Biology* **425**: 4904–4920.
- Langevin C, van der Aa LM, Houel A, Torhy C, Briolat V, Lunazzi A *et al.* (2013b) Zebrafish ISG15 exerts a strong antiviral activity against RNA and DNA viruses and regulates the interferon response. *Journal of Virology* **87**: 10025–10036.
- Larsen R, Rokenes TP, Robertsen B (2004) Inhibition of infectious pancreatic necrosis virus replication by Atlantic salmon Mx1 protein. *Journal of Virology* **78**: 7938–7944.
- Lauksund S, Svingerud T, Bergan V, Robertsen B (2009) Atlantic salmon IPS-1 mediates induction of IFN α 1 and activation of NF-kappaB and localizes to mitochondria. *Developmental and Comparative Immunology* **33**: 1196–1204.
- Lazarte JMS, Kim YR, Lee JS, Im SP, Kim SW, Jung JW *et al.* (2017) Enhancement of glycoprotein-based DNA vaccine for viral hemorrhagic septicemia virus (VHSV) via addition of the molecular adjuvant, DDX41. *Fish and Shellfish Immunology* **62**: 356–365.
- Lazear HM, Nice TJ, Diamond MS (2015) Interferon-lambda: immune functions at barrier surfaces and beyond. *Immunity* **43**: 15–28.
- Levraud JP, Boudinot P, Colin I, Benmansour A, Peyrieras N, Herbomel P *et al.* (2007) Identification of the zebrafish IFN receptor: implications for the origin of the vertebrate IFN system. *Journal of Immunology* **178**: 4385–4394.
- Li Z, Xu X, Huang L, Wu J, Lu Q, Xiang Z *et al.* (2010) Administration of recombinant IFN1 protects zebrafish (*Danio rerio*) from ISKNV infection. *Fish and Shellfish Immunology* **29**: 399–406.
- Li D, Tan W, Ma M, Yu X, Lai Q, Wu Z *et al.* (2012) Molecular characterization and transcription regulation analysis of type I IFN gene in grass carp (*Ctenopharyngodon idella*). *Gene* **504**: 31–40.
- Li D, Lin G, Yu X, Huang S, Lai Q, Liu Y *et al.* (2013) Immunoprotection of grass carp (*Ctenopharyngodon idella*) with recombinant interferon (rCiIFN) against GCHV infection. *Aquaculture* **388**: 42–48.
- Li S, Lu LF, Feng H, Wu N, Chen DD, Zhang YB *et al.* (2014a) IFN regulatory factor 10 is a negative regulator of the IFN responses in fish. *Journal of Immunology* **193**: 1100–1109.
- Li Y, Esain V, Teng L, Xu J, Kwan W, Frost IM *et al.* (2014b) Inflammatory signaling regulates embryonic hematopoietic stem and progenitor cell production. *Genes and Development* **28**: 2597–2612.
- Li S, Lu LF, Wang ZX, Chen DD, Zhang YA (2016) Fish IRF6 is a positive regulator of IFN expression and involved in both of the MyD88 and TBK1 pathways. *Fish and Shellfish Immunology* **57**: 262–268.
- Li S, Lu LF, LaPatra SE, Chen DD, Zhang YA (2017) Zebrafish STAT6 negatively regulates IFN α 1 production by attenuating the kinase activity of TANK-binding kinase 1. *Developmental and Comparative Immunology* **67**: 189–201.

- Liao Z, Wan Q, Su J (2016) Bioinformatics analysis of organizational and expressional characterizations of the IFNs, IRFs and CRFBs in grass carp *Ctenopharyngodon idella*. *Developmental and Comparative Immunology* **61**: 97–106.
- Liongue C, O'Sullivan LA, Trengove MC, Ward AC (2012) Evolution of JAK-STAT pathway components: mechanisms and role in immune system development. *PLoS ONE* **7**: e32777.
- Liu YJ (2005) IPC: professional type 1 interferon-producing cells and plasmacytoid dendritic cell precursors. *Annual Review of Immunology* **23**: 275–306.
- Liu TK, Zhang YB, Liu Y, Sun F, Gui JF (2011) Cooperative roles of fish protein kinase containing Z-DNA binding domains and double-stranded RNA-dependent protein kinase in interferon-mediated antiviral response. *Journal of Virology* **85**: 12769–12780.
- Liu X, Cai X, Zhang D, Xu C, Xiao W (2016) Zebrafish foxo3b negatively regulates antiviral response through suppressing the transactivity of IRF3 and IRF7. *Journal of Immunology* **197**: 4736–4749.
- Liu J, Li J, Xiao J, Chen H, Lu L, Wang X *et al.* (2017) The antiviral signaling mediated by black carp MDA5 is positively regulated by LGP2. *Fish and Shellfish Immunology* **66**: 360–371.
- Liu W, Xiang Y, Zhang W, Jia P, Yi M, Jia K (2018) Expression pattern, antiviral role and regulation analysis of interferon-stimulated gene 15 in black seabream, *Acanthopagrus schlegelii*. *Fish and Shellfish Immunology* **82**: 60–67.
- Long S, Wilson M, Bengten E, Bryan L, Clem LW, Miller NW *et al.* (2004) Identification of a cDNA encoding channel catfish interferon. *Developmental and Comparative Immunology* **28**: 97–111.
- Long S, Milev-Milovanovic I, Wilson M, Bengten E, Clem LW, Miller NW *et al.* (2006) Identification and expression analysis of cDNAs encoding channel catfish type I interferons. *Fish and Shellfish Immunology* **21**: 42–59.
- Loo YM, Gale M Jr (2011) Immune signaling by RIG-I-like receptors. *Immunity* **34**: 680–692.
- Lopez-Munoz A, Roca FJ, Meseguer J, Mulero V (2009) New insights into the evolution of IFNs: zebrafish group II IFNs induce a rapid and transient expression of IFN-dependent genes and display powerful antiviral activities. *Journal of Immunology* **182**: 3440–3449.
- Lopez-Munoz A, Roca FJ, Sepulcre MP, Meseguer J, Mulero V (2010) Zebrafish larvae are unable to mount a protective antiviral response against waterborne infection by spring viremia of carp virus. *Developmental and Comparative Immunology* **34**: 546–552.
- Lopez-Munoz A, Sepulcre MP, Roca FJ, Figueras A, Meseguer J, Mulero V (2011) Evolutionary conserved pro-inflammatory and antigen presentation functions of zebrafish IFN γ revealed by transcriptomic and functional analysis. *Molecular Immunology* **48**: 1073–1083.
- Lu B, Ebensperger C, Dembic Z, Wang Y, Kvatyuk M, Lu T *et al.* (1998) Targeted disruption of the interferon-gamma receptor 2 gene results in severe immune defects in mice. *Proceedings of the National Academy of Sciences of the United States of America* **95**: 8233–8238.
- Lu DQ, Leng TT, Ding X, Peng W, Yao M, Li SS *et al.* (2014) Two IFNGR1 homologues in *Tetraodon nigroviridis*: origin, expression analysis and ligand-binding preference. *Developmental and Comparative Immunology* **44**: 270–279.
- Lu LF, Li S, Lu XB, Zhang YA (2015) Functions of the two zebrafish MAVS variants are opposite in the induction of IFN1 by targeting IRF7. *Fish and Shellfish Immunology* **45**: 574–582.
- Lu L, Wang X, Wu S, Song X, Zou Z, Xie X *et al.* (2017) Black carp STING functions importantly in innate immune defense against RNA virus. *Fish and Shellfish Immunology* **70**: 13–24.
- Luo K, Zhang S, Tang D, Xia L, Gao W, Tian G *et al.* (2018) Analysis of the expression patterns of the cytokine receptor family B (CRFB) and interferon gamma receptor (IFNGR) in Dabry's sturgeon (*Acipenser dabryanus*). *Developmental and Comparative Immunology* **84**: 420–426.
- Lutfalla G, Roest Crollius H, Stange-Thomann N, Jaillon O, Mogensen K, Monneron D (2003) Comparative genomic analysis reveals independent expansion of a lineage-specific gene family in vertebrates: the class II cytokine receptors and their ligands in mammals and fish. *BMC Genomics* **4**: 29.
- Ma JX, Li JY, Fan DD, Feng W, Lin AF, Xiang LX *et al.* (2018) Identification of DEAD-Box RNA helicase DDX41 as a trafficking protein that involves in multiple innate immune signaling pathways in a zebrafish model. *Frontiers in Immunology* **9**: 1327.
- MacMicking J, Xie QW, Nathan C (1997) Nitric oxide and macrophage function. *Annual Review of Immunology* **15**: 323–350.
- Maekawa S, Chiang YA, Hikima J, Sakai M, Lo CF, Wang HC *et al.* (2016) Expression and biological activity of two types of interferon genes in medaka (*Oryzias latipes*). *Fish and Shellfish Immunology* **48**: 20–29.
- Marie I, Durbin JE, Levy DE (1998) Differential viral induction of distinct interferon-alpha genes by positive feedback through interferon regulatory factor-7. *EMBO Journal* **17**: 6660–6669.
- Martin SA, Mohanty BP, Cash P, Houlihan DF, Secombes CJ (2007a) Proteome analysis of the Atlantic salmon (*Salmo salar*) cell line SHK-1 following recombinant IFN-gamma stimulation. *Proteomics* **7**: 2275–2286.
- Martin SA, Taggart JB, Seear P, Bron JE, Talbot R, Teale AJ *et al.* (2007b) Interferon type I and type II responses in an Atlantic salmon (*Salmo salar*) SHK-1 cell line by the salmon TRAITS/SGP microarray. *Physiological Genomics* **32**: 33–44.
- Martin SA, Zou J, Houlihan DF, Secombes CJ (2007c) Directional responses following recombinant cytokine stimulation of rainbow trout (*Oncorhynchus mykiss*) RTS-11 macrophage cells as revealed by transcriptome profiling. *BMC Genomics* **8**: 150.
- Matsumoto M, Araki K, Hayashi K, Takeuchi Y, Shiozaki K, Suetake H *et al.* (2017) Adjuvant effect of recombinant interleukin-12 in the Nocardiosis formalin-killed vaccine of the amberjack *Seriola dumerili*. *Fish and Shellfish Immunology* **67**: 263–269.

- Matsuo A, Oshiumi H, Tsujita T, Mitani H, Kasai H, Yoshimizu M *et al.* (2008) Teleost TLR22 recognizes RNA duplex to induce IFN and protect cells from birnaviruses. *Journal of Immunology* **181**: 3474–3485.
- Matsuyama T, Fujiwara A, Sakai T, Nakayasu C (2009) Molecular cloning and expression analysis of interferon gamma gene in Japanese flounder *Paralichthys olivaceus*. *Fisheries Science* **75**: 253–255.
- Matthews RP, Eauclore SF, Mugnier M, Lorent K, Cui S, Ross MM *et al.* (2011) DNA hypomethylation causes bile duct defects in zebrafish and is a distinguishing feature of infantile biliary atresia. *Hepatology* **53**: 905–914.
- Metzger DC, Elliott DG, Wargo A, Park LK, Purcell MK (2010) Pathological and immunological responses associated with differential survival of Chinook salmon following *Renibacterium salmoninarum* challenge. *Diseases of Aquatic Organisms* **90**: 31–41.
- Milev-Milovanovic I, Long S, Wilson M, Bengten E, Miller NW, Chinchar VG (2006) Identification and expression analysis of interferon gamma genes in channel catfish. *Immunogenetics* **58**: 70–80.
- Milne DJ, Campoverde C, Andree KB, Chen X, Zou J, Secombes CJ (2018) The discovery and comparative expression analysis of three distinct type I interferons in the perciform fish, meagre (*Argyrosomus regius*). *Developmental and Comparative Immunology* **84**: 123–132.
- Mohapatra S, Chakraborty T, Miyagawa S, Zhou L, Ohta K, Iguchi T *et al.* (2015) Steroid responsive regulation of IFN γ alternative splicing and its possible role in germ cell proliferation in medaka. *Molecular and Cellular Endocrinology* **400**: 61–70.
- Moyle WR, Campbell RK, Myers RV, Bernard MP, Han Y, Wang X (1994) Co-evolution of ligand-receptor pairs. *Nature* **368**: 251–255.
- Nayak SK, Shibasaki Y, Nakanishi T (2014) Immune responses to live and inactivated *Nocardia seriolae* and protective effect of recombinant interferon gamma (rIFN γ) against nocardiosis in ginbuna crucian carp, *Carassius auratus langsdorfii*. *Fish and Shellfish Immunology* **39**: 354–364.
- Nie L, Xiong R, Zhang YS, Zhu LY, Shao JZ, Xiang LX (2014) Conserved inhibitory role of teleost SOCS-1s in IFN signaling pathways. *Developmental and Comparative Immunology* **43**: 23–29.
- Núñez Ortiz N, Gerdol M, Stocchi V, Marozzi C, Randelli E, Bernini C *et al.* (2014) T cell transcripts and T cell activities in the gills of the teleost fish sea bass (*Dicentrarchus labrax*). *Developmental and Comparative Immunology* **47**: 309–318.
- Oates AC, Wollberg P, Pratt SJ, Paw BH, Johnson SL, Ho RK *et al.* (1999) Zebrafish stat3 is expressed in restricted tissues during embryogenesis and stat1 rescues cytokine signaling in a STAT1-deficient human cell line. *Developmental Dynamics* **215**: 352–370.
- Ohtani M, Hikima J, Kondo H, Hirono I, Jung TS, Aoki T (2010) Evolutional conservation of molecular structure and antiviral function of a viral RNA receptor, LGP2, in Japanese flounder, *Paralichthys olivaceus*. *Journal of Immunology* **185**: 7507–7517.
- Ohtani M, Hikima J, Kondo H, Hirono I, Jung TS, Aoki T (2011) Characterization and antiviral function of a cytosolic sensor gene, MDA5, in Japanese flounder, *Paralichthys olivaceus*. *Developmental and Comparative Immunology* **35**: 554–562.
- Oidtman B, Dixon P, Way K, Joiner C, Bayley AE (2018) Risk of waterborne virus spread - review of survival of relevant fish and crustacean viruses in the aquatic environment and implications for control measures. *Reviews in Aquaculture* **10**: 641–669.
- Ooi EL, Verjan N, Hirono I, Nochi T, Kondo H, Aoki T *et al.* (2008) Biological characterisation of a recombinant Atlantic salmon type I interferon synthesized in *Escherichia coli*. *Fish and Shellfish Immunology* **24**: 506–513.
- Overgard AC, Nepstad I, Nerland AH, Patel S (2012a) Characterisation and expression analysis of the Atlantic halibut (*Hippoglossus hippoglossus* L.) cytokines: IL-1 β , IL-6, IL-11, IL-12 β and IFN γ . *Molecular Biology Reports* **39**: 2201–2213.
- Overgard AC, Nerland AH, Fiksdal IU, Patel S (2012b) Atlantic halibut experimentally infected with nodavirus shows increased levels of T-cell marker and IFN γ transcripts. *Developmental and Comparative Immunology* **37**: 139–150.
- Palha N, Guivel-Benhassine F, Briolat V, Lutfalla G, Sourisseau M, Ellett F *et al.* (2013) Real-time whole-body visualization of Chikungunya virus infection and host interferon response in zebrafish. *PLoS Pathogens* **9**: e1003619.
- Parhi J, Mukherjee SC, Saxena G, Sahoo L, Makesh M (2014) Molecular characterization and expression of type-I interferon gene in *Labeo rohita*. *Molecular Biology Reports* **41**: 2979–2987.
- Peng W, Lu DQ, Li GF, Zhang X, Yao M, Zhang Y *et al.* (2016) Two distinct interferon-gamma genes in *Tetraodon nigroviridis*: functional analysis during *Vibrio parahaemolyticus* infection. *Molecular Immunology* **70**: 34–46.
- Peng W, Sun Y, Li GF, He LG, Li RZ, Liang YS *et al.* (2018) Two distinct interferon-gamma in the orange-spotted grouper (*Epinephelus coioides*): molecular cloning, functional characterization, and regulation in Toll-like receptor pathway by induction of miR-146a. *Frontiers in Endocrinology* **9**: 41.
- Pereiro P, Costa MM, Diaz-Rosales P, Dios S, Figueras A, Novoa B (2014) The first characterization of two type I interferons in turbot (*Scophthalmus maximus*) reveals their differential role, expression pattern and gene induction. *Developmental and Comparative Immunology* **45**: 233–244.
- Pereiro P, Forn-Cuni G, Figueras A, Novoa B (2016) Pathogen-dependent role of turbot (*Scophthalmus maximus*) interferon-gamma. *Fish and Shellfish Immunology* **59**: 25–35.
- Pestka S, Krause CD, Walter MR (2004) Interferons, interferon-like cytokines, and their receptors. *Immunological Reviews* **202**: 8–32.
- Piehl J, Thomas C, Garcia KC, Schreiber G (2012) Structural and dynamic determinants of type I interferon receptor assembly and their functional interpretation. *Immunological Reviews* **250**: 317–334.

- Pietretti D, Wiegertjes GF (2014) Ligand specificities of Toll-like receptors in fish: indications from infection studies. *Developmental and Comparative Immunology* **43**: 205–222.
- Platanias LC (2005) Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nature Reviews Immunology* **5**: 375–386.
- Poynter S, Lisser G, Monjo A, DeWitte-Orr S (2015) Sensors of infection: viral nucleic acid PRRs in fish. *Biology* **4**: 460–493.
- Purcell MK, Laing KJ, Woodson JC, Thorgaard GH, Hansen JD (2009) Characterization of the interferon genes in homozygous rainbow trout reveals two novel genes, alternate splicing and differential regulation of duplicated genes. *Fish and Shellfish Immunology* **26**: 293–304.
- Qi ZT, Nie P (2008) Comparative study and expression analysis of the interferon gamma gene locus cytokines in *Xenopus tropicalis*. *Immunogenetics* **60**: 699–710.
- Quynh NT, Hikima J, Kim YR, Fagutao FF, Kim MS, Aoki T *et al.* (2015) The cytosolic sensor, DDX41, activates antiviral and inflammatory immunity in response to stimulation with double-stranded DNA adherent cells of the olive flounder, *Paralichthys olivaceus*. *Fish and Shellfish Immunology* **44**: 576–583.
- Raida MK, Buchmann K (2008) Development of adaptive immunity in rainbow trout, *Oncorhynchus mykiss* (Walbaum) surviving an infection with *Yersinia ruckeri*. *Fish and Shellfish Immunology* **25**: 533–541.
- Ran Y, Shu HB, Wang YY (2014) MITA/STING: a central and multifaceted mediator in innate immune response. *Cytokine and Growth Factor Reviews* **25**: 631–639.
- Rao Y, Wan Q, Yang C, Su J (2017) Grass carp Laboratory of genetics and physiology 2 serves as a negative regulator in retinoic acid-inducible gene I- and melanoma differentiation-associated gene 5-mediated antiviral signaling in resting state and early stage of grass carp reovirus infection. *Frontiers in Immunology* **8**: 352.
- Rebl A, Goldammer T (2018) Under control: the innate immunity of fish from the inhibitors' perspective. *Fish and Shellfish Immunology* **77**: 328–349.
- Robertson B (2006) The interferon system of teleost fish. *Fish and Shellfish Immunology* **20**: 172–191.
- Robertson B (2018) The role of type I interferons in innate and adaptive immunity against viruses in Atlantic salmon. *Developmental and Comparative Immunology* **80**: 41–52.
- Robertson B, Bergan V, Rokenes T, Larsen R, Albuquerque A (2003) Atlantic salmon interferon genes: cloning, sequence analysis, expression, and biological activity. *Journal of Interferon and Cytokine Research* **23**: 601–612.
- Robertson B, Chang CJ, Bratland L (2016) IFN-adjuvanted DNA vaccine against infectious salmon anemia virus: antibody kinetics and longevity of IFN expression. *Fish and Shellfish Immunology* **54**: 328–332.
- Rodriguez-Nunez I, Wcisel DJ, Litman RT, Litman GW, Yoder JA (2016) The identification of additional zebrafish DICP genes reveals haplotype variation and linkage to MHC class I genes. *Immunogenetics* **68**: 295–312.
- Rothenfusser S, Goutagny N, DiPerna G, Gong M, Monks BG, Schoenemeyer A *et al.* (2005) The RNA helicase LGP2 inhibits TLR-independent sensing of viral replication by retinoic acid-inducible gene-I. *Journal of Immunology* **175**: 5260–5268.
- Ruan BY, Chen SN, Hou J, Huang B, Laghari ZA, Li L *et al.* (2017) Two type II IFN members, IFN-gamma and IFN-gamma related (rel), regulate differentially IRF1 and IRF11 in zebrafish. *Fish and Shellfish Immunology* **65**: 103–110.
- Santhakumar D, Rubbenstroth D, Martinez-Sobrido L, Munir M (2017) Avian interferons and their antiviral effectors. *Frontiers in Immunology* **8**: 49.
- Sato M, Suemori H, Hata N, Asagiri M, Ogasawara K, Nakao K *et al.* (2000) Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN-alpha/beta gene induction. *Immunity* **13**: 539–548.
- Sato T, Kato H, Kumagai Y, Yoneyama M, Sato S, Matsushita K *et al.* (2010) LGP2 is a positive regulator of RIG-I- and MDA5-mediated antiviral responses. *Proceedings of the National Academy of Sciences of the United States of America* **107**: 1512–1517.
- Savan R, Ravichandran S, Collins JR, Sakai M, Young HA (2009) Structural conservation of interferon gamma among vertebrates. *Cytokine and Growth Factor Reviews* **20**: 115–124.
- Sawamiphak S, Kontarakis Z, Stainier DY (2014) Interferon gamma signaling positively regulates hematopoietic stem cell emergence. *Developmental Cell* **31**: 640–653.
- Schoenborn JR, Wilson CB (2007) Regulation of interferon-gamma during innate and adaptive immune responses. *Advances in Immunology* **96**: 41–101.
- Schoggins JW, Rice CM (2011) Interferon-stimulated genes and their antiviral effector functions. *Current Opinion in Virology* **1**: 519–525.
- Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P *et al.* (2011) A diverse range of gene products are effectors of the type I interferon antiviral response. *Nature* **472**: 481–485.
- Schroder K, Hertzog PJ, Ravasi T, Hume DA (2004) Interferon-gamma: an overview of signals, mechanisms and functions. *Journal of Leukocyte Biology* **75**: 163–189.
- Secombes CJ, Zou J (2017) Evolution of interferons and interferon receptors. *Frontiers in Immunology* **8**: 209.
- Shi J, Zhang YB, Liu TK, Sun F, Gui JF (2012) Subcellular localization and functional characterization of a fish IRF9 from crucian carp *Carassius auratus*. *Fish and Shellfish Immunology* **33**: 258–266.
- Shi J, Zhang YB, Zhang JS, Gui JF (2013) Expression regulation of zebrafish interferon regulatory factor 9 by promoter analysis. *Developmental and Comparative Immunology* **41**: 534–543.
- Shibasaki Y, Yabu T, Araki K, Mano N, Shiba H, Moritomo T *et al.* (2014) Peculiar monomeric interferon gammas, IFN γ mamrel 1 and IFN γ mamrel 2, in ginbuna crucian carp. *FEBS Journal* **281**: 1046–1056.

- Shibasaki Y, Hatanaka C, Matsuura Y, Miyazawa R, Yabu T, Moritomo T *et al.* (2016) Effects of IFN γ administration on allograft rejection in ginbuna crucian carp. *Developmental and Comparative Immunology* **62**: 108–115.
- Sieger D, Stein C, Neifer D, van der Sar AM, Leptin M (2009) The role of gamma interferon in innate immunity in the zebrafish embryo. *Disease Models and Mechanisms* **2**: 571–581.
- Simora RM, Ohtani M, Hikima J, Kondo H, Hirono I, Jung TS *et al.* (2010) Molecular cloning and antiviral activity of IFN-beta promoter stimulator-1 (IPS-1) gene in Japanese flounder, *Paralichthys olivaceus*. *Fish and Shellfish Immunology* **29**: 979–986.
- Skjaeveland I, Iliev DB, Zou J, Jorgensen T, Jorgensen JB (2008) A TLR9 homolog that is up-regulated by IFN-gamma in Atlantic salmon (*Salmo salar*). *Developmental and Comparative Immunology* **32**: 603–607.
- Skjesol A, Hansen T, Shi CY, Thim HL, Jorgensen JB (2010) Structural and functional studies of STAT1 from Atlantic salmon (*Salmo salar*). *BMC Immunology* **11**: 17.
- Skjesol A, Liebe T, Iliev DB, Thomassen EI, Tollersrud LG, Sobhkhez M *et al.* (2014) Functional conservation of suppressors of cytokine signaling proteins between teleosts and mammals: atlantic salmon SOCS1 binds to JAK/STAT family members and suppresses type I and II IFN signaling. *Developmental and Comparative Immunology* **45**: 177–189.
- Sobhkhez M, Hansen T, Iliev DB, Skjesol A, Jorgensen JB (2013) The Atlantic salmon protein tyrosine kinase Tyk2: molecular cloning, modulation of expression and function. *Developmental and Comparative Immunology* **41**: 553–563.
- Sobhkhez M, Skjesol A, Thomassen E, Tollersrud LG, Iliev DB, Sun B *et al.* (2014) Structural and functional characterization of salmon STAT1, STAT2 and IRF9 homologs sheds light on interferon signaling in teleosts. *FEBS Open Bio* **4**: 858–871.
- Sobhkhez M, Joensen LL, Tollersrud LG, Strandskog G, Thim HL, Jorgensen JB (2017a) A conserved inhibitory role of suppressor of cytokine signaling 1 (SOCS1) in salmon antiviral immunity. *Developmental and Comparative Immunology* **67**: 66–76.
- Sobhkhez M, Krasnov A, Chang CJ, Robertsen B (2017b) Transcriptome analysis of plasmid-induced genes sheds light on the role of type I IFN as adjuvant in DNA vaccine against infectious salmon anemia virus. *PLoS ONE* **12**: e0188456.
- Somamoto T, Miura Y, Nakanishi T, Nakao M (2015) Local and systemic adaptive immune responses toward viral infection via gills in ginbuna crucian carp. *Developmental and Comparative Immunology* **52**: 81–87.
- Song H, Yan YL, Titus T, He X, Postlethwait JH (2011) The role of STAT1b in zebrafish hematopoiesis. *Mechanisms of Development* **128**: 442–456.
- Stark GR, Darnell JE Jr (2012) The JAK-STAT pathway at twenty. *Immunity* **36**: 503–514.
- Stark GR, Kerr IM, Williams BRG, Silverman RH, Schreiber RD (1998) How cells respond to interferons. *Annual Review of Biochemistry* **67**: 227–264.
- Stein C, Caccamo M, Laird G, Leptin M (2007) Conservation and divergence of gene families encoding components of innate immune response systems in zebrafish. *Genome Biology* **8**: R251.
- Stolte EH, Savelkoul HFJ, Wiegertjes G, Flik G, Verburg-van Kemenade BML (2008) Differential expression of two interferon-gamma genes in common carp (*Cyprinus carpio* L.). *Developmental and Comparative Immunology* **32**: 1467–1481.
- Subramaniam PS, Mujtaba MG, Paddy MR, Johnson HM (1999) The carboxyl terminus of interferon-gamma contains a functional polybasic nuclear localization sequence. *Journal of Biological Chemistry* **274**: 403–407.
- Sullivan C, Postlethwait JH, Lage CR, Millard PJ, Kim CH (2007) Evidence for evolving Toll-IL-1 receptor-containing adaptor molecule function in vertebrates. *Journal of Immunology* **178**: 4517–4527.
- Sun B, Robertsen B, Wang Z, Liu B (2009) Identification of an Atlantic salmon IFN multigene cluster encoding three IFN subtypes with very different expression properties. *Developmental and Comparative Immunology* **33**: 547–558.
- Sun F, Zhang YB, Liu TK, Gan L, Yu FF, Liu Y *et al.* (2010) Characterization of fish IRF3 as an IFN-inducible protein reveals evolving regulation of IFN response in vertebrates. *Journal of Immunology* **185**: 7573–7582.
- Sun B, Skjaeveland I, Svingerud T, Zou J, Jorgensen J, Robertsen B (2011a) Antiviral activity of salmonid gamma interferon against infectious pancreatic necrosis virus and salmonid alphavirus and its dependency on type I interferon. *Journal of Virology* **85**: 9188–9198.
- Sun F, Zhang YB, Liu TK, Shi J, Wang B, Gui JF (2011b) Fish MITA serves as a mediator for distinct fish IFN gene activation dependent on IRF3 or IRF7. *Journal of Immunology* **187**: 2531–2539.
- Sun B, Greiner-Tollersrud L, Koop BF, Robertsen B (2014) Atlantic salmon possesses two clusters of type I interferon receptor genes on different chromosomes, which allows for a larger repertoire of interferon receptors than in zebrafish and mammals. *Developmental and Comparative Immunology* **47**: 275–286.
- Svingerud T, Solstad T, Sun B, Nyrud ML, Kileng O, Greiner-Tollersrud L *et al.* (2012) Atlantic salmon type I IFN subtypes show differences in antiviral activity and cell-dependent expression: evidence for high IFNb/IFNc-producing cells in fish lymphoid tissues. *Journal of Immunology* **189**: 5912–5923.
- Svingerud T, Holand JK, Robertsen B (2013) Infectious salmon anemia virus (ISAV) replication is transiently inhibited by Atlantic salmon type I interferon in cell culture. *Virus Research* **177**: 163–170.
- Swain B, Basu M, Lenka SS, Das S, Jayasankar P, Samanta M (2015) Characterization and inductive expression analysis of interferon gamma-related gene in the Indian major carp, rohu (*Labeo rohita*). *DNA and Cell Biology* **34**: 367–378.
- Takizawa F, Magadan S, Parra D, Xu Z, Korytar T, Boudinot P *et al.* (2016) Novel teleost CD4-bearing cell populations provide insights into the evolutionary origins and primordial roles of CD4⁺ lymphocytes and CD4⁺ macrophages. *Journal of Immunology* **196**: 4522–4535.

- Tan YP, Zheng J, Tung SL, Rosenshine I, Leung KY (2005) Role of type III secretion in *Edwardsiella tarda* virulence. *Microbiology* **151**: 2301–2313.
- Teijaro JR (2016) Type I interferons in viral control and immune regulation. *Current Opinion in Virology* **16**: 31–40.
- Valero Y, Morcillo P, Meseguer J, Buonocore F, Esteban MA, Chaves-Pozo E et al. (2015) Characterization of the IFN pathway in the teleost fish gonad against vertically transmitted viral nervous necrosis virus. *Journal of General Virology* **96**: 2176–2187.
- Velazquez J, Acosta J, Herrera N, Morales A, Gonzalez O, Herrera F et al. (2017) Novel IFN γ homologue identified in Nile tilapia (*Oreochromis niloticus*) links with immune response in gills under different stimuli. *Fish and Shellfish Immunology* **71**: 275–285.
- Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB et al. (2014) Elephant shark genome provides unique insights into gnathostome evolution. *Nature* **505**: 174–179.
- Wan Q, Wicramaarachchi WD, Whang I, Lim BS, Oh MJ, Jung SJ et al. (2012) Molecular cloning and functional characterization of two duplicated two-cysteine containing type I interferon genes in rock bream *Oplegnathus fasciatus*. *Fish and Shellfish Immunology* **33**: 886–898.
- Wan Q, Yang C, Rao Y, Liao Z, Su J (2017) MDA5 induces a stronger interferon response than RIG-I to GCRV infection through a mechanism involving the phosphorylation and dimerization of IRF3 and IRF7 in CIK Cells. *Frontiers in Immunology* **8**: 189.
- Wang L, Wang L, Zhang HX, Zhang JH, Chen WH, Ruan XF et al. (2006) *In vitro* effects of recombinant zebrafish IFN on spring viremia of carp virus and infectious hematopoietic necrosis virus. *Journal of Interferon and Cytokine Research* **26**: 256–259.
- Wang L, Shang N, Feng H, Guo Q, Dai H (2013) Molecular cloning of grass carp (*Ctenopharyngodon idellus*) T-bet and GATA-3, and their expression profiles with IFN- γ in response to grass carp reovirus (GCRV) infection. *Fish Physiology and Biochemistry* **39**: 793–805.
- Wang B, Zhang YB, Liu TK, Shi J, Sun F, Gui JF (2014) Fish viperin exerts a conserved antiviral function through RLR-triggered IFN signaling pathway. *Developmental and Comparative Immunology* **47**: 140–149.
- Wang R, Li Y, Zhou Z, Liu Q, Zeng L, Xiao T (2018) Involvement of interferon regulatory factor 3 from the barbel chub *Squaliobarbus curriculus* in the immune response against grass carp reovirus. *Gene* **648**: 5–11.
- Wei X, Li XZ, Zheng X, Jia P, Wang J, Yang X et al. (2016) Toll-like receptors and interferon associated immune factors response to spring viraemia of carp virus infection in common carp (*Cyprinus carpio*). *Fish and Shellfish Immunology* **55**: 568–576.
- Wickramaarachchi WD, Wan Q, Lim BS, Jung HB, De Zoysa M, Park MA et al. (2014) Genomic characterization of interferon regulatory factor 5 from rock bream (*Oplegnathus fasciatus*) and its role in antiviral defense. *Fish and Shellfish Immunology* **37**: 256–267.
- Wiegertjes GF, Wentzel AS, Spaink HP, Elks PM, Fink IR (2016) Polarization of immune responses in fish: the ‘macrophages first’ point of view. *Molecular Immunology* **69**: 146–156.
- Workenhe ST, Rise ML, Kibenge MJ, Kibenge FS (2010) The fight between the teleost fish immune response and aquatic viruses. *Molecular Immunology* **47**: 2525–2536.
- Wu J, Chen ZJ (2014) Innate immune sensing and signaling of cytosolic nucleic acids. *Annual Review of Immunology* **32**: 461–488.
- Wu Z, Wang L, Xu X, Lin G, Mao H, Ran X et al. (2017) Interaction of IRF9 and STAT2 synergistically up-regulates IFN and PKR transcription in *Ctenopharyngodon idella*. *Molecular Immunology* **85**: 273–282.
- Wu H, Liu L, Wu S, Wang C, Feng C, Xiao J et al. (2018) IFN β of black carp functions importantly in host innate immune response as an antiviral cytokine. *Fish and Shellfish Immunology* **74**: 1–9.
- Xia S, Wang H, Hong X, Lu J, Xu D, Jiang Y et al. (2018) Identification and characterization of a type I interferon induced by cyprinid herpesvirus 2 infection in crucian carp *Carassius auratus gibelio*. *Fish and Shellfish Immunology* **76**: 35–40.
- Xiang Z, Qi L, Chen W, Dong C, Liu Z, Liu D et al. (2011) Characterization of a TnMAVS protein from *Tetraodon nigroviridis*. *Developmental and Comparative Immunology* **35**: 1103–1115.
- Xiang Y, Liu W, Jia P, Li Y, Jin Y, Chen L et al. (2017) Molecular characterization and expression analysis of interferon- γ in black seabream *Acanthopagrus schlegelii*. *Fish and Shellfish Immunology* **70**: 140–148.
- Xiao J, Yan J, Chen H, Li J, Tian Y, Feng H (2016) LGP2 of black carp plays an important role in the innate immune response against SVCV and GCRV. *Fish and Shellfish Immunology* **57**: 127–135.
- Xiao J, Yan C, Zhou W, Li J, Wu H, Chen T et al. (2017) CARD and TM of MAVS of black carp play the key role in its self-association and antiviral ability. *Fish and Shellfish Immunology* **63**: 261–269.
- Xie HX, Lu JF, Rolhion N, Holden DW, Nie P, Zhou Y et al. (2014) *Edwardsiella tarda*-induced cytotoxicity depends on its type III secretion system and flagellin. *Infection and Immunity* **82**: 3436–3445.
- Xiong R, Nie L, Xiang LX, Shao JZ (2012) Characterization of a PIAS4 homologue from zebrafish: insights into its conserved negative regulatory mechanism in the TRIF, MAVS, and IFN signaling pathways during vertebrate evolution. *Journal of Immunology* **188**: 2653–2668.
- Xu C, Guo TC, Mutoloki S, Haugland O, Marjara IS, Evensen O (2010a) Alpha interferon and not gamma interferon inhibits salmonid alphavirus subtype 3 replication in vitro. *Journal of Virology* **84**: 8903–8912.
- Xu X, Huang L, Weng S, Wang J, Lin T, Tang J et al. (2010b) *Tetraodon nigroviridis* as a nonlethal model of infectious spleen and kidney necrosis virus (ISKNV) infection. *Virology* **406**: 167–175.
- Xu C, Evensen O, Munang’andu HM (2015) De novo assembly and transcriptome analysis of Atlantic salmon macrophage/

- dendritic-like TO cells following type I IFN treatment and salmonid alphavirus subtype-3 infection. *BMC Genomics* **16**: 96.
- Xu C, Evensen O, Munang'andu HM (2016) A de novo transcriptome analysis shows that modulation of the JAK-STAT signaling pathway by salmonid alphavirus subtype 3 favors virus replication in macrophage/dendritic-like TO-cells. *BMC Genomics* **17**: 390.
- Xu X, Li M, Wu Z, Wang H, Wang L, Huang K *et al.* (2017) Endoplasmic reticulum transmembrane proteins ZDHHC1 and STING both act as direct adaptors for IRF3 activation in teleost. *Journal of Immunology* **199**: 3623–3633.
- Yabu T, Toda H, Shibasaki Y, Araki K, Yamashita M, Anzai H *et al.* (2011) Antiviral protection mechanisms mediated by ginbuna crucian carp interferon gamma isoforms 1 and 2 through two distinct interferon gamma-receptors. *Journal of Biochemistry* **150**: 635–648.
- Yan J, Peng L, Li Y, Fan H, Tian Y, Liu S *et al.* (2016) IFN α of triploid hybrid of gold fish and allotetraploid is an antiviral cytokine against SVCV and GCRV. *Fish and Shellfish Immunology* **54**: 529–536.
- Yan C, Xiao J, Li J, Chen H, Liu J, Wang C *et al.* (2017a) TBK1 of black carp plays an important role in host innate immune response against SVCV and GCRV. *Fish and Shellfish Immunology* **69**: 108–118.
- Yan J, Peng L, Chi M, Xiao J, Li J, Liu S *et al.* (2017b) IFN α 2 of triploid hybrid of gold fish and allotetraploid is an intracellular antiviral cytokine against SVCV and GCRV. *Fish and Shellfish Immunology* **62**: 238–246.
- Yang K, Zhang S, Chen D, Zhang A, Wang X, Zhou H (2013) IFN-gamma-activated lymphocytes boost nitric oxide production in grass carp monocytes/macrophages. *Fish and Shellfish Immunology* **35**: 1635–1641.
- Yang Q, Liu X, Zhou T, Cook J, Nguyen K, Bai X (2016a) RNA polymerase II pausing modulates hematopoietic stem cell emergence in zebrafish. *Blood* **128**: 1701–1710.
- Yang Y, Huang Y, Yu Y, Zhou S, Wang S, Yang M *et al.* (2016b) Negative regulation of the innate antiviral immune response by TRIM62 from orange spotted grouper. *Fish and Shellfish Immunology* **57**: 68–78.
- Yang S, Li Q, Mu Y, Ao J, Chen X (2017) Functional activities of interferon gamma in large yellow croaker *Larimichthys crocea*. *Fish and Shellfish Immunology* **70**: 545–552.
- Yeh DW, Liu YL, Lo YC, Yuh CH, Yu GY, Lo JF *et al.* (2013) Toll-like receptor 9 and 21 have different ligand recognition profiles and cooperatively mediate activity of CpG-oligodeoxynucleotides in zebrafish. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 20711–20716.
- Yi S, Lu D, Peng W, Wang T, Zhang Y, Lin H (2014) Differential expression profiling of spleen microRNAs in response to two distinct type II interferons in *Tetraodon nigroviridis*. *PLoS ONE* **9**: e96336.
- Yim HY, Park C, Lee YD, Arimoto K, Jeon R, Baek SH *et al.* (2016) Elevated response to type I IFN enhances RANKL-mediated osteoclastogenesis in Usp18-knockout mice. *Journal of Immunology* **196**: 3887–3895.
- Yoon S, Alnabulsi A, Wang TY, Lee PT, Chen TY, Bird S *et al.* (2016) Analysis of interferon gamma protein expression in zebrafish (*Danio rerio*). *Fish and Shellfish Immunology* **57**: 79–86.
- Yu FF, Zhang YB, Liu TK, Liu Y, Sun F, Jiang J *et al.* (2010) Fish virus-induced interferon exerts antiviral function through STAT1 pathway. *Molecular Immunology* **47**: 2330–2341.
- Yu Y, Huang X, Zhang J, Liu J, Hu Y, Yang Y *et al.* (2016a) Fish TRIM16L exerts negative regulation on antiviral immune response against grouper iridoviruses. *Fish and Shellfish Immunology* **59**: 256–267.
- Yu Y, Huang Y, Yang Y, Wang S, Yang M, Huang X *et al.* (2016b) Negative regulation of the antiviral response by grouper LGP2 against fish viruses. *Fish and Shellfish Immunology* **56**: 358–366.
- Yu CF, Peng WM, Schlee M, Barchet W, Eis-Hubinger AM, Kolanus W *et al.* (2018a) SOCS1 and SOCS3 target IRF7 degradation to suppress TLR7-mediated type I IFN production of human plasmacytoid dendritic cells. *Journal of Immunology* **200**: 4024–4035.
- Yu N, Xu X, Qi G, Liu D, Chen X, Ran X *et al.* (2018b) *Ctenopharyngodon idella* TBK1 activates innate immune response via IRF7. *Fish and Shellfish Immunology* **80**: 521–527.
- Zahradnik J, Kolarova L, Parizkova H, Kolenko P, Schneider B (2018) Interferons type II and their receptors R1 and R2 in fish species: evolution, structure, and function. *Fish and Shellfish Immunology* **79**: 140–152.
- Zhang YB, Gui JF (2012) Molecular regulation of interferon antiviral response in fish. *Developmental and Comparative Immunology* **38**: 193–202.
- Zhang J, Zhang YB, Wu M, Wang B, Chen C, Gui JF (2014) Fish MAVS is involved in RLR pathway-mediated IFN response. *Fish and Shellfish Immunology* **41**: 222–230.
- Zhang L, Chen WQ, Hu YW, Wu XM, Nie P, Chang MX (2016) TBK1-like transcript negatively regulates the production of IFN and IFN-stimulated genes through RLRs-MAVS-TBK1 pathway. *Fish and Shellfish Immunology* **54**: 135–143.
- Zhang J, Huang X, Ni S, Liu J, Hu Y, Yang Y *et al.* (2017) Grouper STAT1 α is involved in antiviral immune response against iridovirus and nodavirus infection. *Fish and Shellfish Immunology* **70**: 351–360.
- Zhang QM, Zhao X, Li Z, Wu M, Gui JF, Zhang YB (2018) Alternative splicing transcripts of zebrafish LGP2 gene differentially contribute to IFN antiviral response. *Journal of Immunology* **200**: 688–703.
- Zhao X, Wang R, Li Y, Xiao T (2017) Molecular cloning and functional characterization of interferon regulatory factor 7 of the barbel chub, *Squaliobarbus curriculus*. *Fish and Shellfish Immunology* **69**: 185–194.
- Zhong X, Chen M, Ding M, Zhong M, Li B, Wang Y *et al.* (2017) Clr and Cls from Nile tilapia (*Oreochromis niloticus*): molecular characterization, transcriptional profiling upon bacterial and

- IFN-gamma inductions and potential role in response to bacterial infection. *Fish and Shellfish Immunology* **70**: 240–251.
- Zhou W, Zhou J, Lv Y, Qu Y, Chi M, Li J et al. (2015) Identification and characterization of MAVS from black carp *Mylopharyngodon piceus*. *Fish and Shellfish Immunology* **43**: 460–468.
- Zhou W, Pal AS, Hsu AY, Gurol T, Zhu X, Wirbisky-Hershberger SE et al. (2018) MicroRNA-223 suppresses the canonical NF-kappaB pathway in basal keratinocytes to dampen neutrophilic inflammation. *Cell Reports* **22**: 1810–1823.
- Zou J, Secombes CJ (2011) Teleost fish interferons and their role in immunity. *Developmental and Comparative Immunology* **35**: 1376–1387.
- Zou J, Yasutoshi YB, Dijkstra JM, Sakai M, Ototake M, Secombes C (2004) Identification of an interferon gamma homologue in fugu, *Takifugu rubripes*. *Fish and Shellfish Immunology* **17**: 403–409.
- Zou J, Carrington A, Collet B, Dijkstra JM, Yoshiura Y, Bols N et al. (2005) Identification and bioactivities of IFN-gamma in rainbow trout *Oncorhynchus mykiss*: the first Th1-type cytokine characterized functionally in fish. *Journal of Immunology* **175**: 2484–2494.
- Zou J, Tafalla C, Truckle J, Secombes CJ (2007) Identification of a second group of type I IFNs in fish sheds light on IFN evolution in vertebrates. *Journal of Immunology* **179**: 3859–3871.
- Zou J, Chang M, Nie P, Secombes CJ (2009) Origin and evolution of the RIG-I like RNA helicase gene family. *BMC Evolutionary Biology* **9**: 85.
- Zou J, Gorgoglione B, Taylor NG, Summated T, Lee PT, Panigrahi A et al. (2014a) Salmonids have an extraordinary complex type I IFN system: characterization of the IFN locus in rainbow trout *Oncorhynchus mykiss* reveals two novel IFN subgroups. *Journal of Immunology* **193**: 2273–2286.
- Zou PF, Chang MX, Xue NN, Liu XQ, Li JH, Fu JP et al. (2014b) Melanoma differentiation-associated gene 5 in zebrafish provoking higher interferon-promoter activity through signalling enhancing of its shorter splicing variant. *Immunology* **141**: 192–202.
- Zou PF, Chang MX, Li Y, Huan Zhang S, Fu JP, Chen SN et al. (2015) Higher antiviral response of RIG-I through enhancing RIG-I/MAVS-mediated signaling by its long insertion variant in zebrafish. *Fish and Shellfish Immunology* **43**: 13–24.