
Being injective removes one of Direct's handicaps, and not the larger one
deg 5, spread 0.75 (4.1 values of h fit each y)
deg 3, spread 0.75 (2.8 values of h fit each y)
deg 6, spread 0.75 (5.1 values of h fit each y)
z^4 (2.0 values of h fit each y)
deg 4, spread 0.25 (2.8 values of h fit each y)
deg 4, spread 0.75 (3.8 values of h fit each y)
deg 6, spread 0.25 (3.3 values of h fit each y)
z^6 (2.0 values of h fit each y)
deg 6, spread 0.5 (4.4 values of h fit each y)
deg 4, spread 0.5 (3.4 values of h fit each y)
deg 5, spread 1 (4.5 values of h fit each y)
deg 4, spread 1 (3.9 values of h fit each y)
deg 3, spread 0.5 (2.5 values of h fit each y)
deg 6, spread 1 (5.5 values of h fit each y)
deg 3, spread 1 (3.0 values of h fit each y)
deg 5, spread 0.25 (2.3 values of h fit each y)
deg 5, spread 0.5 (3.4 values of h fit each y)
deg 3, spread 0.25 (1.9 values of h fit each y)
z^5 (INJECTIVE)
z^3 (INJECTIVE)
hyperparameter estimation alone, measured on the linear maps
no bar: exactly zero
no bar: exactly zero
0.0
0.1
0.2
0.3
0.4
how far Direct's prediction of the objective is from Composite's
(Kolmogorov distance between the two posteriors; bigger = Direct is more wrong)
everything that is NOT information
(hyperparameters + model shape)
Direct never sees the intermediates h
(vanishes exactly when g is injective)