

Colosense output example


Case 4,

2024 data

The output represent only 1 scenario of guideline retrieval (based on my most frequently go-to guideline. RAG may retrieve different relevant guidelines and give outputs with subtle revision.)


Category 1, Q1.1

Question	Output template	Output
1.1 How should the severity of this patient be classified using validated scoring systems?	Remission, mild, moderate, severe	The patient is in remission because total Mayo score was 1. (partial Mayo score 0, MES 1).




Category 1, Q1.2

Question	Output template	Output
1.2 What is the remission status of the patient?	Clinical remission, bio-chemical remission, endoscopic remission, histologic remission.	The patient has achieved clinical remission (pMayo=0), bio-chemical remission (CRP 0.02, FC 25), endoscopic remission (MES 1), and histologic remission (Nancy 1).




Category 1, Q1.3

Question	Output template	Output
1.3 Does this patient have any poor prognostic factor?	Yes, specify which. No.	The patient has the below poor prognostic factors: young onset (at 11 years old) and extensive colitis.




Category 2, Q2.1

Question	Output template	Output
2.1 Has the patient achieved treatment target, according to the treat-to-target strategy	The patient has achieved short / intermediate / and/or long term treatment target.	Yes the patient had achieved long-term treatment target (endoscopic remission).



Category 2, Q2.2


Question	Output template	Output
2.2 Based on the patient's current status, should the medication be adjusted?	Yes, according to treat-to-target strategy, the current medication should be adjusted. No.	No.



Category 3, Q3.1

Extensive colitis.
MES 1, Nancy 1
No PSC
No family history
Disease duration: 18 years


Question	Output template	Output
3.1 Is the risk for the patient getting colorectal cancer high? When should the patient be screened for colorectal cancer?	Since the patient belongs to low / intermediate / high risk group, the next surveillance colonoscopy should be in ___ years.	[Tier 1] Since the patient belongs to low risk group, the next surveillance colonoscopy should be in 5 years. [ECCO, 2023]



Category 3, Q3.2

29 years old female
No PSC
Active medication:
Mezavant XL 4800 mg

Question	Output template	Output
3.2 What are the other types of cancer that the patient should be aware of? How and when should the patient be screened?	Based on the patient's sex, age, underlying disease, and medication history, the patient should receive screening for ___ cancer with ___, every ___ year.	[Tier 1] Based on the patient's sex, age, underlying disease, and medication history, the patient should receive screening for cervical cancer with cervical cytology, every 3 year. [AGA, 2025]




Category 4, Q4.1

In remission
Active medication:
Mezavant XL 4800 mg since
2024/7/16

Last CRP =0.02 (2024/4/23)
Last FC=25 (2024/11/28)


Question	Output template	Output
4.1 What should the next non-invasive monitoring exams be scheduled to assess disease activities?	Based on the patient's current status, the following exams ____ should be arranged at ____.	[Tier 1] Based on the patient's current status, the interval biomarker monitoring may be performed every 6-12 months. [AGA, 2023] Assessment of subclinical inflammation with biomarkers (FC) and intestinal ultrasound every 6-12 months should be considered. [ECCO-ESGAR-ESP-IBUS, 2025]



Category 4, Q4.2

**Active medication:
Mezavant XL 4800 mg
(not advanced therapy
or immunomodulator)**

Question	Output template	Output
4.2 Should therapeutic drug monitoring be performed for this patient's current therapy?	Yes, proactive / reactive TDM is recommended, with target drug level____. No.	No current evidence support TDM for the patient.




Category 4, Q4.3

**Active medication:
Mezavant XL 4800 mg**

**Treatment duration:
Mezavant for 4 years
(since 2022/3/22)**


Question	Output template	Output
4.3 Based on the medication history, is there any medication-specific monitor during the treatment?	For patients under ____ medication, ____ should be checked every ____ months.	[Tier 1] For patients under Mezavant XL, renal function should be monitored periodically. [AGA, 2019]



Category 4, Q4.4

29 years old female
No PSC
Active medication:
Mezavant XL 4800 mg
Past medication:
Mezavant XL 2400 mg

Question	Output template	Output
4.4 What opportunistic infection risk and vaccinations should be given according to the patient's status?	Screening for ____ and ____ vaccinations prior to treatment initiation are recommended.	[Tier 1] Annual influenza vaccine and hepatitis B vaccine (if not immune) should be performed. Herpes zoster vaccines should be given prior to starting immune-modifying therapy. [ACG, 2025]



Category 5, Q5.1

Question	Output template	Output
5.1 Is there any dietary recommendation for the patient?	This patient is encouraged to have more ____ intake and less ____.	[Tier 1] Mediterranean diet could be considered for maintenance of remission. This patient is encouraged to have less red and processed meat intake. [ECCO, 2025]

29 years old female
In remission
Active medication:
Mezavant XL 4800 mg

Category 5, Q5.2

Question	Output template	Output
5.2 Does this patient require nutritional supplementation or screening for deficiency based on the history?	Yes, the patient is recommended to be screened for ____ deficiency. No.	[Tier 1] Nutritional assessment is recommended at diagnosis, then annually. The patient is recommended to be monitored for hemoglobin, iron, folate, vitamin D, vitamin B12, and zinc. [ECCO, 2025]

29 years old female
In remission
Active medication:
Mezavant XL 4800 mg

Category 5, Q5.3

Question	Output template	Output
5.3 What lifestyle modifications are suggested for the patient?	The patient should quit ____ and enhance ____.	[Tier 3] The patient may increase physical activity and mindfulness-based therapies. [Rozich, 2025]

29 years old female
In remission
Active medication:
Mezavant XL 4800 mg
No smoking.

Category 6, Q6.1

Question	Output template	Output
6.1 Is the patient's current medication safe to continue during pregnancy, lactation, or while trying to conceive?	These ____ medications were safe to be continued. These ____ medication should be stopped ____ months before conception.	[Tier 1] Mezavant XL is safe during pregnancy and is recommended to continue. [AGA, 2025]

29 years old female
In remission
Active medication:
Mezavant XL 4800 mg

Category 6, Q6.2

Question	Output template	Output
6.2 What are the risks to the patient associated with the patient's disease activity and current medications?	Maternally, the risk of _____ is increased / comparable to the non-IBD patients.	[Tier 1] Maternally, the risk of relapse or worsening disease is increased. [ECCO, 2023] Controlling disease activity during pregnancy is critical to reduce adverse outcomes. [AGA, 2025]

29 years old female
In remission
Active medication:
Mezavant XL 4800 mg

Category 6, Q6.3

Question	Output template	Output
6.3 What are the risks to the fetus associated with the patient's disease activity and current medications?	Neonatally, the risk of ____ is increased / comparable to the mothers of non-IBD patients.	[Tier 1] Neonatally, the risks of low birth weight and preterm delivery are increased to the mothers of non-IBD patients. [AGA, 2025] Controlling disease activity during pregnancy is critical to reduce adverse outcomes. [AGA, 2025]

29 years old female
In remission
Active medication:
Mezavant XL 4800 mg