

CSD2301 Practice Solutions

15. **Angular Momentum**

LIN QINJIE

Practice Question 1

A woman with mass 50 kg is standing on the rim of a large disk that is rotating at 0.50 rev/s about an axis through its center. The disk has a mass 110 kg and radius 4.0 m. Calculate the magnitude of the total angular momentum of the woman-plus-disk system. Assume that you can treat the person as a point in your calculation.

IDENTIFY: $L = I\omega$ and $I = I_{\text{disk}} + I_{\text{woman}}$.

SET UP: $\omega = 0.50 \text{ rev/s} = 3.14 \text{ rad/s}$. $I_{\text{disk}} = \frac{1}{2}m_{\text{disk}}R^2$ and $I_{\text{woman}} = m_{\text{woman}}R^2$.

EXECUTE: $I = (55 \text{ kg} + 50.0 \text{ kg})(4.0 \text{ m})^2 = 1680 \text{ kg} \cdot \text{m}^2$. $L = (1680 \text{ kg} \cdot \text{m}^2)(3.14 \text{ rad/s}) = 5.28 \times 10^3 \text{ kg} \cdot \text{m}^2/\text{s}$

EVALUATE: The disk and the woman have similar values of I , even though the disk has twice the mass.

$\frac{1}{2}m_{\text{disk}}$

m_{woman}

Practice Question 2

Find the magnitude of the angular momentum of the second hand on a clock about an axis through the center of the clock face. The clock hand has a length of 15.0 cm and a mass of 6.00 g. Take the second hand to be a slender rod rotating with constant angular velocity about one end.

IDENTIFY and SET UP: Use $L = I\omega$

EXECUTE: The second hand makes 1 revolution in 1 minute, so

$$\omega = (1.00 \text{ rev/min})(2\pi \text{ rad/1 rev})(1 \text{ min/60 s}) = 0.1047 \text{ rad/s}$$

For a slender rod, with the axis about one end,

$$I = \frac{1}{3}ML^2 = \frac{1}{3}(6.00 \times 10^{-3} \text{ kg})(0.150 \text{ m})^2 = 4.50 \times 10^{-5} \text{ kg} \cdot \text{m}^2$$

$$\text{Then } L = I\omega = (4.50 \times 10^{-5} \text{ kg} \cdot \text{m}^2)(0.1047 \text{ rad/s}) = 4.71 \times 10^{-6} \text{ kg} \cdot \text{m}^2/\text{s}.$$

EVALUATE: \vec{L} is clockwise.

Solution takes clockwise to be +ve.
So if answer of L is +ve, means it
is clockwise

Practice Question 3

A diver jumps off a board with arms straight up and legs straight down, giving her a moment of inertia about her rotation axis of $18 \text{ kg}\cdot\text{m}^2$. She then tucks into a small ball, decreasing this moment of inertia to $3.6 \text{ kg}\cdot\text{m}^2$. While tucked, she makes two complete revolutions in 1.0 s . If she hadn't tucked at all, how many revolutions would she have made in the 1.5 s from board to water?

IDENTIFY: Apply conservation of angular momentum to the diver. $I_i \omega_i = I_f \omega_f$

SET UP: The number of revolutions she makes in a certain time is proportional to her angular velocity. The ratio of her untucked to tucked angular velocity $(\frac{\omega_i}{\omega_f})$ is $(\frac{I_f}{I_i} = \frac{3.6 \text{ kg}\cdot\text{m}^2}{18 \text{ kg}\cdot\text{m}^2})$.

EXECUTE: If she had untucked, she would have made $(2 \text{ rev} \times \frac{3.6 \text{ kg}\cdot\text{m}^2}{18 \text{ kg}\cdot\text{m}^2} = 0.40 \text{ rev})$ in the last 1.0 s , so she would have made $(0.40 \text{ rev})(1.5/1.0) = 0.60 \text{ rev}$ in the total 1.5 s .

Practice Question 4

A hollow thin walled sphere of mass 12.0 kg and diameter 48.0 cm is rotating about an axle through its center. The angle (in radians) through which it turns as a function of time (in seconds) is given by $\theta(t) = At^2 + Bt^4$, where A has a numerical value of 1.50 and B has a numerical value of 1.10. (a) What are the units of the constants A and B? (b) At the time 3.00 s find (i) the angular momentum of the sphere and (ii) the net torque on the sphere.

IDENTIFY: $\omega_z = d\theta/dt$. $L_z = I\omega_z$ and $\tau_z = dL_z/dt$.

SET UP: For a hollow, thin-walled sphere rolling about an axis through its center, $I = \frac{2}{3}MR^2$. $R = 0.240$ m .

EXECUTE: (a) $A = 1.50 \text{ rad/s}^2$ and $B = 1.10 \text{ rad/s}^4$, so that $\theta(t)$ will have units of radians.

(b) (i) $\omega_z = \frac{d\theta}{dt} = 2At + 4Bt^3$. At $t = 3.00$ s , $\omega_z = 2(1.50 \text{ rad/s}^2)(3.00 \text{ s}) + 4(1.10 \text{ rad/s}^4)(3.00 \text{ s})^3 = 128 \text{ rad/s}$.

$L_z = (\frac{2}{3}MR^2)\omega_z = \frac{2}{3}(12.0 \text{ kg})(0.240 \text{ m})^2(128 \text{ rad/s}) = 59.0 \text{ kg} \cdot \text{m}^2/\text{s}$.

(ii) $\tau_z = \frac{dL_z}{dt} = I \frac{d\omega_z}{dt} = I(2A + 12Bt^2)$ and Differentiate ω to get α . Because $\tau = I\alpha$

$\tau_z = \frac{2}{3}(12.0 \text{ kg})(0.240 \text{ m})^2(2[1.50 \text{ rad/s}^2] + 12[1.10 \text{ rad/s}^4][3.00 \text{ s}]^2) = 56.1 \text{ N} \cdot \text{m}$.

Practice Question 5

A large wooden turntable in the shape of a flat uniform disk has a radius of 2.00 m and a total mass of 120 kg. The turntable is initially rotating at 3.00 rad/s about a vertical axis through its center. Suddenly, a 70.0 kg parachutist makes a soft landing on the turntable at a point near the outer edge. (a) Find the angular speed of the turntable after the parachutist lands. (Assume that you can treat the parachutist as a particle.) (b) Compute the kinetic energy of the system before and after the parachutist lands. Why are these kinetic energies not equal?

EXECUTE: (a) $L_1 = L_2$ gives $I_1\omega_1 = I_2\omega_2$, so $\omega_2 = (I_1/I_2)\omega_1$

$$I_1 = I_{\text{tt}} = \frac{1}{2}MR^2 = \frac{1}{2}(120 \text{ kg})(2.00 \text{ m})^2 = 240 \text{ kg} \cdot \text{m}^2$$

$$I_2 = I_{\text{tt}} + I_{\text{p}} = 240 \text{ kg} \cdot \text{m}^2 + mR^2 = 240 \text{ kg} \cdot \text{m}^2 + (70 \text{ kg})(2.00 \text{ m})^2 = 520 \text{ kg} \cdot \text{m}^2$$

$$\omega_2 = (I_1/I_2)\omega_1 = (240 \text{ kg} \cdot \text{m}^2 / 520 \text{ kg} \cdot \text{m}^2)(3.00 \text{ rad/s}) = 1.38 \text{ rad/s}$$

(b) $K_1 = \frac{1}{2}I_1\omega_1^2 = \frac{1}{2}(240 \text{ kg} \cdot \text{m}^2)(3.00 \text{ rad/s})^2 = 1080 \text{ J}$

$$K_2 = \frac{1}{2}I_2\omega_2^2 = \frac{1}{2}(520 \text{ kg} \cdot \text{m}^2)(1.38 \text{ rad/s})^2 = 495 \text{ J}$$

EVALUATE: The kinetic energy decreases because of the negative work done on the turntable and the parachutist by the friction force between these two objects.