

- 1 X is an impure sample of a Group 2 metal carbonate, MCO_3 . X contains 57% by mass of MCO_3 . The impurities in X do **not** react with hydrochloric acid.

7.4 g of X is reacted with an excess of dilute hydrochloric acid.

0.050 mol of the Group 2 metal chloride is produced.

What is the identity of the Group 2 metal?

- A** Mg **B** Ca **C** Sr **D** Ba

- 2 Which of these samples of gas contains the same number of atoms as 1 g of hydrogen gas?

A 22 g of carbon dioxide (M_r : CO_2 , 44)

B 8 g of methane (M_r : CH_4 , 16)

C 20 g of neon (M_r : Ne, 20)

D 8 g of ozone (M_r : O_3 , 48)

- 3 What is the total number of protons, neutrons and electrons present in an ammonium ion with a relative formula mass of 21?



	number of protons	number of neutrons	number of electrons
A	11	10	10
B	10	11	11
C	10	11	10
D	11	10	11

- 4 This question is about the first ionisation energies of magnesium and neon.

Which row is correct?

	first ionisation energy	type of electron removed	
		from Mg	from Ne
A	Mg > Ne	p	s
B	Mg > Ne	s	p
C	Ne > Mg	p	s
D	Ne > Mg	s	p

25 Which alkene shows geometric isomerism?



26 What is the correct name of the major product of the reaction of HBr with 3-ethylhex-3-ene?

A 3-bromo-3-ethylhexane

B 3-bromo-4-ethylhexane

C 4-bromo-3-ethylhexane

D 4-bromo-4-ethylhexane

27 The alkane $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)_2$ undergoes free radical substitution with chlorine. No C–C bonds are broken in this reaction.

How many isomeric products, including positional and optical isomers, of molecular formula $\text{C}_5\text{H}_{11}\text{Cl}$ can be formed?

A 4

B 5

C 6

D 7

28 What is involved in the mechanism of the reaction between aqueous NaOH and 1-bromobutane?

A attack by a nucleophile on a carbon atom with a partial positive charge

B heterolytic bond fission and attack by a nucleophile on a carbocation

C homolytic bond fission and attack by an electrophile on a carbanion

D homolytic bond fission and attack by a nucleophile on a carbocation

29 But-2-ene reacts with cold dilute acidified KMnO_4 to give product X.

But-2-ene reacts with an excess of hot concentrated acidified KMnO_4 to give product Y.

Which statement about X and Y is correct?

- A** Only one of X and Y reacts with 2,4-dinitrophenylhydrazine.
- B** X and Y both react with sodium hydroxide.
- C** X and Y both react with sodium metal.
- D** Y reacts with LiAlH_4 to give X.

30 When heated with KOH dissolved in ethanol, halogenoalkanes can undergo an elimination reaction to form alkenes.

What are the possible elimination products when 2-bromobutane is heated with KOH dissolved in ethanol?

- A** $\text{CH}_3\text{CH}=\text{CHCH}_3$ only
- B** $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ only
- C** $\text{CH}_3\text{CH}=\text{CHCH}_3$ and $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$
- D** $\text{CH}_3\text{CH}=\text{CHCH}_3$ and $\text{CH}_2=\text{CHCH}=\text{CH}_2$

31 Chloroethane can be used to make sodium propanoate.

chloroethane \rightarrow intermediate Q \rightarrow sodium propanoate

Intermediate Q is hydrolysed with boiling aqueous NaOH to give sodium propanoate.

Which reagent would produce intermediate Q from chloroethane?

- A** concentrated ammonia solution
- B** dilute sulfuric acid
- C** hydrogen cyanide in water
- D** potassium cyanide in ethanol

32 Four different alcohols are treated with alkaline $\text{I}_2(\text{aq})$.

Which row is correct?

	name of alcohol	formulae of products
A	butan-2-ol	CH_3COO^- and CH_3CI_3
B	propan-1-ol	CH_3COO^- and CHI_3
C	propan-2-ol	CH_3COO^- and CHI_3
D	butan-2-ol	$\text{CH}_3\text{CH}_2\text{COO}^-$ and CH_3I

33 The M_r of compound X is 88.

Compound X is heated under reflux with an excess of acidified potassium dichromate(VI) to produce compound Y.

Compound Y reacts with compound X under suitable conditions to produce compound Z. The M_r of compound Z is 172.

What is compound X?

- A** $\text{CH}_3\text{CH}_2\text{CHOHCH}_2\text{CH}_3$
- B** $(\text{CH}_3)_2\text{COHCH}_2\text{CH}_3$
- C** $(\text{CH}_3)_2\text{CHCHOHCH}_3$
- D** $(\text{CH}_3)_3\text{CCH}_2\text{OH}$

34 Butanedione, $\text{CH}_3\text{COCOCH}_3$, is a yellow liquid.

How does butanedione react with 2,4-dinitrophenylhydrazine reagent and Fehling's reagent?

	2,4-dinitrophenylhydrazine	Fehling's
A	positive	positive
B	positive	negative
C	negative	positive
D	negative	negative

35 Which substance reacts with ethanoic acid to give the organic product with the highest M_r ?

- A** lithium aluminium hydride
- B** magnesium
- C** potassium carbonate
- D** propan-2-ol

36 A sample of propyl ethanoate is hydrolysed by heating under reflux with aqueous NaOH. The two organic products of the hydrolysis are separated, purified and weighed.

Out of the total mass of products obtained, what is the percentage by mass of each product?

- A** 32.4% and 67.6%
- B** 38.3% and 61.7%
- C** 42.3% and 57.7%
- D** 50.0% and 50.0%

37 Which statement about PVC is correct?

- A** Combustion products of PVC are very alkaline and harmful to breathe in.
- B** The empirical formula of PVC is the same as the empirical formula of the monomer.
- C** Molecules of PVC are unsaturated.
- D** The repeat unit of PVC is $(\text{CH}_2\text{CCl}_2)$.

38 Compound Q reacts separately with HCN and NaBH_4 under suitable conditions.

Both reactions produce an organic product with a chiral centre.

What is compound Q?

- A** butanone
- B** ethanal
- C** propanal
- D** propanone

39 Compound X has the following properties.

- When 0.20 mol of X undergoes complete combustion, 14.4 dm^3 of carbon dioxide is produced, measured under room conditions.
- X reacts with 2,4-dinitrophenylhydrazine reagent to give an orange crystalline product.
- X does **not** give a yellow precipitate with alkaline $\text{I}_2(\text{aq})$.

What could be X?

- A** hexan-3-one
- B** propanal
- C** propan-1-ol
- D** propanone

BLANK PAGE

BLANK PAGE

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.


Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge

The Periodic Table of Elements

Group																					
1	2															13	14	15	16	17	18

40 Measles is an infectious disease caused by a virus.

The graph shows the number of cases of measles each year in a country before and after a vaccine was introduced.



What could have caused the decrease in the number of cases of measles after vaccination was introduced?

	vaccines provided artificial active immunity in people	vaccines provided artificial passive immunity in people	fewer people are able to act as hosts for the virus
A	✓	✗	✓
B	✗	✓	✓
C	✗	✓	✗
D	✓	✗	✗

key
✓ = yes
✗ = no

- 5** Arsenic forms a compound with fluorine. In this compound, the arsenic atom has no lone pair of electrons and there are no dative bonds.


Selenium also forms a compound with fluorine. In this compound, the selenium atom has no lone pair of electrons and there are no dative bonds.

In which compounds are there two different bond angles?

(In this question, 180° bond angles should be ignored.)

- A** both arsenic fluoride and selenium fluoride
- B** arsenic fluoride only
- C** selenium fluoride only
- D** neither arsenic fluoride nor selenium fluoride

- 6** A structure for borazole, $\text{N}_3\text{B}_3\text{H}_6$, is shown.




Which shape is borazole and how many π electrons are there in the structure?

	shape	number of π electrons
A	non-planar	3
B	non-planar	6
C	planar	3
D	planar	6

7 The diagram shows the apparatus used to find the relative molecular mass of a volatile liquid.

When 0.10 g of a volatile liquid is injected into the syringe, all of the volatile liquid evaporates and the volume increases by 85 cm³.

The heater maintains a temperature of 400 K and the experiment is carried out at a pressure of 101 300 Pa.



If the vapour of the volatile liquid behaves as an ideal gas, which expression can be used to calculate the relative molecular mass of the liquid?

- A** $M_r = (85 \times 101\,300) \div (0.10 \times 8.31 \times 400)$
- B** $M_r = (85 \times 101.3) \div (0.10 \times 8.31 \times 400)$
- C** $M_r = (0.10 \times 8.31 \times 400) \div (85 \times 10^{-6} \times 101\,300)$
- D** $M_r = (0.10 \times 8.31 \times 400) \div (85 \times 10^{-6} \times 101.3)$


8 The table shows physical properties of four substances, W, X, Y and Z.

	melting point/°C	boiling point/°C	electrical conductivity of solid	electrical conductivity of liquid	electrical conductivity in water
W	993	1695	poor	good	good
X	-119	39	poor	poor	insoluble
Y	1535	2750	good	good	insoluble
Z	1610	2230	poor	poor	insoluble

What are the identities of W, X, Y and Z?

	W	X	Y	Z
A	MgO	C ₂ H ₅ Br	Fe	Al ₂ O ₃
B	MgO	HCl	K	Al ₂ O ₃
C	NaF	C ₂ H ₅ Br	Fe	SiO ₂
D	NaF	HCl	K	SiO ₂

- 9 The apparatus used to determine a value for the enthalpy of combustion of butan-1-ol is shown. The mass of 1.00 cm^3 of water is 1.00 g .




initial mass of burner + butan-1-ol	58.34 g
initial temperature of water	$17.6\text{ }^{\circ}\text{C}$
final mass of burner + butan-1-ol	57.85 g
final temperature of water	$41.1\text{ }^{\circ}\text{C}$

butan-1-ol $M_r = 74$

Which value, to three significant figures, for the enthalpy of combustion of butan-1-ol can be calculated from these data?


- A -114 J mol^{-1}
- B -17.2 kJ mol^{-1}
- C -2600 kJ mol^{-1}
- D -4540 kJ mol^{-1}

- 10** In the high temperatures of car engines, nitrogen reacts with oxygen to produce nitrogen monoxide.



This reaction has activation energy E_a .


Which reaction pathway diagram correctly represents this reaction?



- 11** In which reaction does the oxidation number of chlorine change by the largest amount?

- A** $2\text{KClO}_3 \rightarrow 2\text{KCl} + 3\text{O}_2$
- B** $2\text{ClO}^- \rightarrow \text{Cl}^- + \text{ClO}_2^-$
- C** $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow \text{HCl} + \text{HClO}$
- D** $2\text{NaClO}_2 + \text{Cl}_2 \rightarrow 2\text{NaCl} + 2\text{ClO}_2$

12 Hydrogen is produced industrially from methane as shown in the equation.



Which conditions give the highest yield of hydrogen at equilibrium?

	pressure	temperature
A	low	high
B	high	low
C	high	high
D	low	low


13 W moles of HNO_2 undergoes a disproportionation reaction to produce U moles of HNO_3 and V moles of NO.

- No other nitrogen containing product is produced.
- Nitrogen is the only element oxidised or reduced.

What are the values of W, U and V?

	W	U	V
A	2	1	1
B	3	1	2
C	5	3	2
D	5	1	4

14 Gas X dissociates on heating to set up the following equilibrium.




A quantity of gas X is heated at constant pressure, p , at a certain temperature. The equilibrium partial pressure of gas X is found to be $\frac{1}{7}p$.

What is the equilibrium constant, K_p , at this temperature?

- A** $\frac{6}{7}p$ **B** $\frac{9}{7}p$ **C** $\frac{36}{7}p$ **D** $9p$

- 15** In the diagram, X is the Boltzmann distribution for the energies of the particles in a reaction and E_{A1} is the activation energy for that reaction.



Which statement is correct?

- A** E_{A2} is the activation energy at a higher temperature.
- B** E_{A2} is the activation energy at a lower temperature.
- C** Y is the Boltzmann distribution at a lower temperature.
- D** Z is the Boltzmann distribution at a higher temperature.

- 16** Magnesium, aluminium and silicon are elements in the Periodic Table. Each element forms an oxide.

Which row is correct?

	MgO	Al_2O_3	SiO_2
A	basic	amphoteric	amphoteric
B	giant ionic	simple molecular	giant ionic
C	high melting point	high melting point	low melting point
D	slight reaction with water	no reaction with water	no reaction with water

- 17** Which statement correctly describes what happens when silicon tetrachloride is added to water?

- A** The $SiCl_4$ dissolves to give a neutral solution only.
- B** The $SiCl_4$ reacts to give an acidic solution only.
- C** The $SiCl_4$ reacts to give a precipitate and an acidic solution.
- D** The $SiCl_4$ reacts to give a precipitate and a neutral solution.

18 X and Y are two elements from Period 3 of the Periodic Table.

Element X has a higher electrical conductivity than element Y. Element Y has a higher melting point than element X.

Which formula is a compound formed from element X and element Y?

- A** MgS **B** Mg₂Si **C** NaCl **D** SiCl₄

19 A sample consisting of 1.0 mol of anhydrous calcium nitrate is completely decomposed by strong heating.

What is the total amount of gas produced in this reaction?

- A** 1.0 mol **B** 2.0 mol **C** 2.5 mol **D** 3.0 mol

20 Steam is passed over heated magnesium to give compound J and hydrogen.


What is **not** a property of compound J?

- A** It has an M_r of 40.3.
B It is basic.
C It is a white solid.
D It is very soluble in water.


21 Which statement is correct?

- A** Hydrogen bromide reduces concentrated sulfuric acid to form sulfur dioxide gas.
B Hydrogen bromide decomposes at a higher temperature than hydrogen chloride.
C When hydrogen bromide gas is shaken with aqueous silver nitrate a yellow precipitate is formed.
D When hydrogen bromide gas is bubbled through aqueous iodine the solution becomes colourless.

22 ICl is made when Cl_2 and I_2 react together.



ICl reacts with water.



Which row is correct?

	oxidation number of I in ICl	reaction occurring when ICl reacts with H_2O
A	+1	the iodine atoms are oxidised to form I_2
B	+1	the iodine atoms are oxidised to form HIO_3
C	-1	the chlorine atoms are reduced to form HCl
D	-1	the iodine atoms are oxidised to form HIO_3

23 NH_4Cl reacts with NaOH in an aqueous solution.

Which statement is correct?

- A** The reaction gives rise to two different polar product molecules.
- B** The bond angle in the nitrogen-containing species remains unchanged.
- C** The ammonium ion acts as a base.
- D** The oxidation state of nitrogen increases in the reaction.

24 What is produced when 60 g of nitrogen monoxide reacts with an excess of carbon monoxide in a catalytic converter?

- A** 12 g of carbon and 92 g of nitrogen dioxide
- B** 24 g of carbon and 92 g of nitrogen dioxide
- C** 88 g of carbon dioxide and 28 g of nitrogen
- D** 88 g of carbon dioxide and 56 g of nitrogen