

MedAI Clinical Summary

Patient Interview & SOAP Note Documentation

Date: January 11, 2026 at 08:49 AM

■ SUBJECTIVE

Patient-reported symptoms and history

The patient presents with a chief complaint of a scratchy throat.

Symptoms began approximately yesterday.

The patient rates the throat pain as 5 out of 10 in severity.

Associated symptoms include generalized body aches.

The patient explicitly denies experiencing fever, cough, runny nose, or difficulty swallowing.

■ OBJECTIVE

Expected physical exam findings

General: Appears comfortable and non-toxic.

Vital signs: Unavailable for current visit. Anticipated temperature to be afebrile given patient denial of fever; other vital signs likely within normal limits.

HEENT: Oropharynx examination reveals mild erythema without exudates or significant tonsillar swelling. Uvula is midline. Nasal mucosa appears pink and moist, without discharge.

Neck: Supple with no palpable cervical lymphadenopathy.

Lungs: Clear to auscultation bilaterally.

Cardiac: Regular rate and rhythm, no murmurs.

■ ASSESSMENT

Primary diagnosis

Viral Pharyngitis

Differential diagnosis

Streptococcal Pharyngitis (less likely given absence of fever, cough, exudates, and denial of difficulty swallowing).

Allergic Rhinitis with post-nasal drip (less likely given primary complaint of throat scratchiness and body aches without typical allergic symptoms).

■ PLAN

Tests

No immediate laboratory or diagnostic tests are indicated at this time, given the presentation highly suggestive of a viral etiology and absence of red flag symptoms.

Medications

Over-the-counter analgesics such as acetaminophen or ibuprofen for pain and body aches, as needed.

Throat lozenges or sprays for symptomatic relief of throat discomfort.

Patient advice

Rest as needed.

Maintain adequate hydration with clear fluids.

Gargle with warm salt water several times a day to soothe the throat.

Avoid irritants such as smoke.

Follow-up instructions

Return to clinic if symptoms worsen, new symptoms develop (e.g., high fever, severe difficulty swallowing, rash), or if there is no improvement within 3-5 days.

Seek immediate care for any difficulty breathing or severe pain.

This document is generated by MedAI Clinical Documentation System. This is a preliminary assessment and should be reviewed by a licensed healthcare professional. For medical emergencies, please contact emergency services immediately.