

Matrix Assignment

Mohamed Hamdan

September 2022

Problem Statement - ABC is a triangle right angled at C. A line through the mid-point M of hypotenuse AB and parallel to BC intersects AC at D. Show that

1. D is the mid-point of AC
2. $MD \perp AC$
3. $CM = MA = \frac{1}{2}AB$

The magnitudes of CM and AM are therefore

$$\|C - M\| = \|q - p\| = (q - p)^T(q - p) \quad (11)$$

$$\|A - M\| = \|r - p\| = (r - p)^T(r - p) \quad (12)$$

$$(13)$$

Upon expanding the vector products, the terms $q^T p$ and $r^T p$ evaluate to 0 (from eq.5). From eq.2,

$$\|q\| = \|r\|$$

. Therefore,

$$\|C - M\| = \|q\| + \|p\| \quad (14)$$

$$\|A - M\| = \|r\| + \|p\| = \|q\| + \|p\| \quad (15)$$

Equating (14) and (15), we get

$$AM = CM$$

Since given that M is midpoint of AB,

$$AM = \frac{1}{2}AB$$

Equating (14) and (15), the result is proved.

Solution

Part 1

Given MN is \parallel to BC, hence

$$\frac{AD}{DC} = \frac{AM}{MB} \quad (1)$$

Since M is the mid-point of AB, $\frac{AM}{MB} = 1$. Substituting in (1), we get

$$AD = DC \quad (2)$$

Therefore, D is the midpoint of AC.

Part 2

Since ABC is right angled at C,

$$(C - A)^T(C - B) = 0 \quad (3)$$

Given that MD is parallel to BC, so

$$(C - B) = \lambda(M - D) \quad (4)$$

Substituting (4) in (3) and dividing by λ , we get

$$(C - A)^T(M - B) = 0 \quad (5)$$

From (5) it can be concluded that $MD \perp AC$.

Part 3

Let

$$M - D = p \quad (6)$$

$$C - D = q \quad (7)$$

$$A - D = r \quad (8)$$

Then vectors along CM and AM can be written as

$$C - M = q - p \quad (9)$$

$$A - M = r - p \quad (10)$$

Construction

The input parameters are the lengths a and c.

Symbol	Value	Description
a	4	BC
c	5	AC
k	1	$\frac{AM}{MB} = \frac{AD}{DC}$
θ	$\arctan(\frac{c}{a})$	$\angle B$
A	$\sqrt{a^2 + c^2} \begin{pmatrix} \cos\theta \\ \sin\theta \end{pmatrix}$	Point A