

# Conic section Assignment

Mohamed Hamdan

September 2022

**Problem Statement** - Find the area of the triangle formed by the lines joining the vertex of the parabola  $x^2 = 12y$  to the ends of its latus rectum

**Solution**




Figure 1: Triangle formed by vertex and ends of latus rectum of parabola  $x^2 = 12y$

The given equation of parabola  $x^2 = 12y$  can be written in the general quadratic form as

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (1)$$

where

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (2)$$

$$\mathbf{u} = \begin{pmatrix} 1 \\ -6 \end{pmatrix}, \quad (3)$$

$$f = 0 \quad (4)$$

The parabola in (1) can be expressed in standard form (center/vertex at origin, major-axis -  $x$  axis) as

$$\mathbf{y}^T \mathbf{D} \mathbf{y} = -2\eta \mathbf{e}_1^T \mathbf{y} \quad |\mathbf{V}| = 0 \quad (5)$$

where

$$\mathbf{x} = \mathbf{P} \mathbf{y} + \mathbf{c} \quad (\text{Affine Transformation}) \quad (6)$$

$$\mathbf{P}^T \mathbf{V} \mathbf{P} = \mathbf{D}. \quad (\text{Eigenvalue Decomposition}) \quad (7)$$

$$\mathbf{D} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad (8)$$

$$\mathbf{P} = (\mathbf{p}_1 \quad \mathbf{p}_2), \quad \mathbf{P}^T = \mathbf{P}^{-1}, \quad (9)$$

$$\eta = \mathbf{u}^T \mathbf{p}_1 \quad (10)$$

$$\mathbf{e}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (11)$$

To find  $\mathbf{c}$  which is the center of the parabola in (1), substitute (6) in (1)

$$(\mathbf{P} \mathbf{y} + \mathbf{c})^T \mathbf{V} (\mathbf{P} \mathbf{y} + \mathbf{c}) + 2\mathbf{u}^T (\mathbf{P} \mathbf{y} + \mathbf{c}) + f = 0, \quad (12)$$

yielding

$$\mathbf{y}^T \mathbf{P}^T \mathbf{V} \mathbf{P} \mathbf{y} + 2(\mathbf{V} \mathbf{c} + \mathbf{u})^T \mathbf{P} \mathbf{y} + \mathbf{c}^T \mathbf{V} \mathbf{c} + 2\mathbf{u}^T \mathbf{c} + f = 0 \quad (13)$$

From (13) and (7),

$$\mathbf{y}^T \mathbf{D} \mathbf{y} + 2(\mathbf{V} \mathbf{c} + \mathbf{u})^T \mathbf{P} \mathbf{y} + \mathbf{c}^T (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^T \mathbf{c} + f = 0 \quad (14)$$

For a parabola  $|\mathbf{V}| = 0, \lambda_1 = 0$  and

$$\mathbf{V} \mathbf{p}_1 = 0, \mathbf{V} \mathbf{p}_2 = \lambda_2 \mathbf{p}_2. \quad (15)$$

where  $\mathbf{p}_1, \mathbf{p}_2$  are the eigenvectors of  $\mathbf{V}$  such that (7)

$$\mathbf{P} = (\mathbf{p}_1 \quad \mathbf{p}_2), \quad (16)$$

Substituting (16) in (14),

$$\begin{aligned} & \mathbf{y}^T \mathbf{D} \mathbf{y} + 2(\mathbf{c}^T \mathbf{V} + \mathbf{u}^T) (\mathbf{p}_1 \quad \mathbf{p}_2) \mathbf{y} \\ & + \mathbf{c}^T (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^T \mathbf{c} + f = 0 \\ & \implies \mathbf{y}^T \mathbf{D} \mathbf{y} \\ & + 2((\mathbf{c}^T \mathbf{V} + \mathbf{u}^T) \mathbf{p}_1 (\mathbf{c}^T \mathbf{V} + \mathbf{u}^T) \mathbf{p}_2) \mathbf{y} \\ & + \mathbf{c}^T (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^T \mathbf{c} + f = 0 \\ & \implies \mathbf{y}^T \mathbf{D} \mathbf{y} \\ & + 2(\mathbf{u}^T \mathbf{p}_1 \quad (\lambda_2 \mathbf{c}^T + \mathbf{u}^T) \mathbf{p}_2) \mathbf{y} \\ & + \mathbf{c}^T (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^T \mathbf{c} + f = 0 \text{ from (15)} \end{aligned}$$

$$\begin{aligned} & \implies \lambda_2 y_2^2 + 2(\mathbf{u}^T \mathbf{p}_1) y_1 + 2y_2 (\lambda_2 \mathbf{c} + \mathbf{u})^T \mathbf{p}_2 \\ & + \mathbf{c}^T (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^T \mathbf{c} + f = 0 \end{aligned}$$

which is the equation of a parabola. Thus, (17) can be expressed as (5) by choosing

$$\eta = \mathbf{u}^T \mathbf{p}_1 \quad (17)$$

and  $\mathbf{c}$  in (14) such that

$$\mathbf{P}^T (\mathbf{V}\mathbf{c} + \mathbf{u}) = \eta \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (18)$$

$$\mathbf{c}^T (\mathbf{V}\mathbf{c} + \mathbf{u}) + \mathbf{u}^T \mathbf{c} + f = 0 \quad (19)$$

Multiplying (18) by  $\mathbf{P}$  yields

$$(\mathbf{V}\mathbf{c} + \mathbf{u}) = \eta \mathbf{p}_1, \quad (20)$$

which, upon substituting in (19) results in

$$\eta \mathbf{c}^T \mathbf{p}_1 + \mathbf{u}^T \mathbf{c} + f = 0 \quad (21)$$

(20) and (21) can be clubbed together to obtain (22).

$$\begin{pmatrix} \mathbf{u}^T + \eta \mathbf{p}_1^T \\ \mathbf{V} \end{pmatrix} \mathbf{c} = \begin{pmatrix} -f \\ \eta \mathbf{p}_1 - \mathbf{u} \end{pmatrix} \quad |V| = 0 \quad (22)$$