

Week 8, 2024

Fractional step and alternating direction methods

Formulation of the problem:

The 2D Heat equation:

$$\frac{\partial U}{\partial t} = \alpha^2 \left( \frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} \right), \quad x \in (0, 1), \quad y \in (0, 1)$$

With the following boundary conditions:

$$U(x = 0, 0 < y < 0.7) = 0$$


$$U(x = 0, 0.7 < y < 1) = 1$$

$$U(x = 1, 0 < y < 0.3) = 1$$

$$U(x = 1, 0.3 < y < 1) = 0$$

$$U(x, y = 0) = 0$$

$$U(x, y = 1) = 0$$



Initial condition:  $U(t = 0, x, y) = 0$

**Numerical method:** Fractional step and alternating direction methods

These two implicit methods are very similar to each other because the main idea of this methods based on fractioning time internal on 2 or 3 parts for each own dimension so problem is converted into one-dimensional problem, and we can solve it easily as we solved in first 3 week doing simple Thomas algorithm. The approximation error of methods  $O(\Delta t^2, \Delta x^2, \Delta y^2)$ .

P.S. I hope you know how to solve the Thomas algorithm and how to find coefficients alpha and betta.

a) Implicit method FSM (Fractional step method)

First step:

$$\frac{U_{ij}^{n+1/2} - U_{ij}^n}{\Delta t} = \frac{1}{2}(\Lambda_1 U^{n+1/2} + \Lambda_1 U^n) + \Lambda_2 U^n$$

Second step:

$$\frac{U_{ij}^{n+1} - U_{ij}^{n+1/2}}{\Delta t} = \frac{1}{2}(\Lambda_2 U^{n+1} - \Lambda_2 U^n)$$

Here is the  $\Lambda_1$  and  $\Lambda_2$  operators have the following form:

$$\Lambda_1 = \alpha^2 \frac{\partial^2}{\partial x^2}, \quad \Lambda_2 = \alpha^2 \frac{\partial^2}{\partial y^2}$$

First step:

$$\frac{U_{ij}^{n+1/2} - U_{ij}^n}{\Delta t} = \frac{1}{2} \left( \alpha^2 \frac{U_{i+1j}^{n+1/2} - 2U_{ij}^{n+1/2} + U_{i-1j}^{n+1/2}}{\Delta x^2} + \alpha^2 \frac{U_{i+1j}^n - 2U_{ij}^n + U_{i-1j}^n}{\Delta x^2} \right) + \alpha^2 \frac{U_{ij+1}^n - 2U_{ij}^n + U_{ij-1}^n}{\Delta y^2}$$

The equation is converted in 1D, and we can solve it by the Thomas algorithm.

$$A_{ij} U_{i+1j}^{n+1/2} + B_{ij} U_{ij}^{n+1/2} + C_{ij} U_{i-1j}^{n+1/2} = D_{ij}, \quad i = \overline{1, N-1}, j \text{ is fixed}$$

$$A_{ij} = -\frac{\alpha^2}{2\Delta x^2}, \quad B_{ij} = \frac{1}{\Delta t} + \frac{\alpha^2}{\Delta x^2}, \quad C_{ij} = -\frac{\alpha^2}{2\Delta x^2}$$

$$D_{ij} = \frac{U_{ij}^n}{\Delta t} + \frac{1}{2} \left( \alpha^2 \frac{U_{i+1j}^n - 2U_{ij}^n + U_{i-1j}^n}{\Delta x^2} \right) + \alpha^2 \frac{U_{ij+1}^n - 2U_{ij}^n + U_{ij-1}^n}{\Delta y^2}$$

Second step:

$$\frac{U_{ij}^{n+1} - U_{ij}^{n+1/2}}{\Delta t} = \frac{1}{2} \left( \alpha^2 \frac{U_{ij+1}^{n+1} - 2U_{ij}^{n+1} + U_{ij-1}^{n+1}}{\Delta y^2} - \alpha^2 \frac{U_{ij+1}^n - 2U_{ij}^n + U_{ij-1}^n}{\Delta y^2} \right)$$

Thomas algorithm for y:

$$A_{ij} U_{ij+1}^{n+1} + B_{ij} U_{ij}^{n+1} + C_{ij} U_{ij-1}^{n+1} = D_{ij}, \quad j = \overline{1, M-1}, i \text{ is fixed}$$

$$A_{ij} = -\frac{\alpha^2}{2\Delta y^2}, \quad B_{ij} = \frac{1}{\Delta t} + \frac{\alpha^2}{\Delta y^2}, \quad C_{ij} = -\frac{\alpha^2}{2\Delta y^2}$$

$$D_{ij} = \frac{U_{ij}^{n+1/2}}{\Delta t} - \frac{1}{2} \left( \alpha^2 \frac{U_{ij+1}^n - 2U_{ij}^n + U_{ij-1}^n}{\Delta y^2} \right)$$

a) Implicit method ADM (Alternating direction method)

First step:

$$\frac{U_{ij}^{n+1/2} - U_{ij}^n}{\Delta t} = \Lambda_1 U^{n+1/2} + \Lambda_2 U^n$$

Second step:

$$\frac{U_{ij}^{n+1} - U_{ij}^{n+1/2}}{\Delta t} = \Lambda_1 U^{n+1/2} + \Lambda_2 U^{n+1}$$

Here is the  $\Lambda_1$  and  $\Lambda_2$  operators have the following form:

$$\Lambda_1 = \alpha^2 \frac{\partial^2}{\partial x^2}, \quad \Lambda_2 = \alpha^2 \frac{\partial^2}{\partial y^2}$$

First step:

$$\frac{U_{ij}^{n+1/2} - U_{ij}^n}{\Delta t} = \alpha^2 \frac{U_{i+1j}^{n+1/2} - 2U_{ij}^{n+1/2} + U_{i-1j}^{n+1/2}}{\Delta x^2} + \alpha^2 \frac{U_{ij+1}^n - 2U_{ij}^n + U_{ij-1}^n}{\Delta y^2}$$

The equation is converted in 1D, and we can solve it by the Thomas algorithm.

$$A_{ij} U_{i+1j}^{n+1/2} + B_{ij} U_{ij}^{n+1/2} + C_{ij} U_{i-1j}^{n+1/2} = D_{ij}, \quad i = \overline{1, N-1}, j \text{ is fixed}$$

$$A_{ij} = -\frac{\alpha^2}{\Delta x^2}, \quad B_{ij} = \frac{1}{\Delta t} + \frac{2\alpha^2}{\Delta x^2}, \quad C_{ij} = -\frac{\alpha^2}{\Delta x^2}$$

$$D_{ij} = \frac{U_{ij}^n}{\Delta t} + \alpha^2 \frac{U_{ij+1}^n - 2U_{ij}^n + U_{ij-1}^n}{\Delta y^2}$$

Second step:

$$\frac{U_{ij}^{n+1} - U_{ij}^{n+1/2}}{\Delta t} = \alpha^2 \frac{U_{i+1j}^{n+1/2} - 2U_{ij}^{n+1/2} + U_{i-1j}^{n+1/2}}{\Delta x^2} + \alpha^2 \frac{U_{ij+1}^{n+1} - 2U_{ij}^{n+1} + U_{ij-1}^{n+1}}{\Delta y^2}$$

Thomas algorithm for y:

$$A_{ij} U_{ij+1}^{n+1} + B_{ij} U_{ij}^{n+1} + C_{ij} U_{ij-1}^{n+1} = D_{ij}, \quad j = \overline{1, M-1}, i \text{ is fixed}$$

$$A_{ij} = -\frac{\alpha^2}{\Delta y^2}, \quad B_{ij} = \frac{1}{\Delta t} + \frac{2\alpha^2}{\Delta y^2}, \quad C_{ij} = -\frac{\alpha^2}{\Delta y^2}$$

$$D_{ij} = \frac{U_{ij}^{n+1/2}}{\Delta t} + \alpha^2 \frac{U_{i+1j}^{n+1/2} - 2U_{ij}^{n+1/2} + U_{i-1j}^{n+1/2}}{\Delta x^2}$$