

Pathway of Greenland Meltwater Revealed by Passive Tracers in a NEMO Simulation

Xianmin Hu and Paul G. Myers

Department of Earth and Atmosphere Sciences
University of Alberta

3rd Annual VITALS Science Meeting, Halifax, NS, Oct 19-21, 2015


Outline

- **Model configuration and experiment setup**
- **Results**
 - evolution of passive tracer content
 - travel time to the Labrador Sea
 - vertical distribution of passive tracers
- **Summary and future work**

Outline


- **Model configuration and experiment setup**
- **Results**
 - evolution of passive tracer content
 - travel time to the Labrador Sea
 - vertical distribution of passive tracers
- **Summary and future work**

Model Configuration



ANHA: Arctic and Northern Hemisphere Atlantic

Experiment Setup



Initialization:

3D T, S, U and V (GLORYS2v3, Jan02)
SSH and Sea Ice

Atmospheric forcing (CGRF, hourly):

T2, Q2, U10, V10
Precipitation ~ 33km
Radiation (SW & LW)

Runoff:

Inter-annual Dai and Trenberth's runoff
+ Jonathan Bamber's Greenland melt

OBC:

U, V, T and S (GLORYS2v3)

NO temperature & salinity restoring

Jan 2002 – Dec 2013


CGRF: CMC GDPS reforecasts

GDPS: Global Deterministic Prediction System


CMC: Canadian Meteorological Centre

GLORYS: GLObal Ocean ReanalYses and Simulations

How to Create the Runoff Data




How to Add the Passive Tracers



- five passive tracers
- proportional to the amount of runoff
- start adding tracers from Jan 2004

$$\Delta c = \frac{\Delta t}{\rho_o \cdot e3t_1} \cdot rnf$$


$$\frac{s}{\frac{kg}{m^3} \cdot m} \cdot \frac{kg}{m^2 \cdot s} = \textit{unitless}$$



Outline


- Model configuration and experiment setup
- **Results**
 - evolution of passive tracer content
 - travel time to the Labrador Sea
 - vertical distribution of passive tracers
- Summary and future work

Evolution of Total Passive Tracer Content





$$TC(x, y, t) = \int c(x, y, z, t) dz$$

Travel Time to Reach the Labrador Sea




$$TS(t) = \iiint c(x, y, z, t) dz dy dx$$


Travel Time to Reach the Labrador Sea




Vertical Distribution of Passive Tracer




Vertical Distribution of Passive Tracer



Vertical Distribution of Passive Tracer



Vertical Distribution of Passive Tracer



Outline

- Model configuration and experiment setup
- Results
 - evolution of passive tracer content
 - travel time to the Labrador Sea
 - vertical distribution of passive tracers
- **Summary and future work**

Summary and Future Work

- passive tracers nicely show the pathway of Greenland meltwater
- Greenland meltwater, particularly from the southeast, can reach Labrador Sea continent shore and slope region within one year, then spread in the subpolar gyre.
- Labrador Sea interior is accumulating over the whole simulation period
- large amount of passive tracers accumulated within Baffin Bay, even the deep basin, why?
- how does the Greenland meltwater impact coastal currents