

The Metric side of Optimal Transport

Jonas Müller

Outline

1 A metric for probabilities?

- The Wasserstein distance
- Dudley Lemma
- Comparing L^2 norm and Wasserstein distance
- Wasserstein for normal distributions

2 Lifting completeness from X to $\mathcal{P}_p(X)$

- Iterated Dudley Lemma

3 Some things in case time permits

- Wasserstein ∞ metric
- Some other properties of the Wasserstein metric

The Wasserstein distance for finite p

As a reminder, we have for any $p \in [1, \infty[$

$$\mathcal{P}_p(X) := \left\{ \mu \in \mathcal{P}(X) \mid \int_X d^p(x, x_0) d\mu(x) < \infty \right\}$$

and the following will be a metric on this space.

Definition

For any $p \in [1, \infty[$ and $\nu, \mu \in \mathcal{P}_2(X)$

$$W_p^p(\mu, \nu) := \min \left\{ \int_{X \times X} d^p(x, y) d\pi(x, y) \mid \pi \in \Gamma(\mu, \nu) \right\}.$$

Theorem

$(\mathcal{P}_p(X), W_p)$ is a metric space!

Proof of Theorem

We assume $p = 2$, which makes it less messy, the other cases are similar.

1. $W_2(\mu, \nu) < \infty$ always
2. $\mu = \nu \Leftrightarrow W_2(\mu, \nu) = 0$
3. $W_2(\mu, \nu) = W_2(\nu, \mu)$
4. Dudley Lemma for triangle inequality
5. $W_2(\mu_1, \mu_3) \leq W_2(\mu_1, \mu_2) + W_2(\mu_2, \mu_3)$.

Reminder

Let $T : X \rightarrow Y$ be a measurable function and let μ, ν be measures on X, Y , then we have the Change of Variables Formula:

$$\int_Y f(y) d(T_{\#}\mu)(y) = \int_X f(T(x)) d\mu(x).$$

Additionally, for any metric d on X it holds for any $x_0 \in X$ that

$$d^2(x, y) \leq (d(x, x_0) + d(x_0, y))^2 \leq 2(d^2(x, x_0) + d^2(x_0, y))$$

because $2ab \leq a^2 + b^2$ always.

The Dudley Lemma

In the Literature this is often called Gluing Lemma!

Lemma

Let $(X_1, \mu_1), (X_1, \mu_2), (X_1, \mu_3)$ be Polish and $\pi^{1,2} \in \Gamma(\mu_1, \mu_2)$ and $\pi^{2,3} \in \Gamma(\mu_2, \mu_3)$. Then there exists some $\pi \in \mathcal{P}(X_1 \times X_2 \times X_3)$ such that

$$p_{\#}^{1,2}(\pi) = \pi^{1,2} \quad \text{and} \quad p_{\#}^{2,3}(\pi) = \pi^{2,3}$$

where $p^{1,2}(x_1, x_2, x_3) = (x_1, x_2)$ and $p^{2,3}(x_1, x_2, x_3) = (x_2, x_3)$.

Proof is essentially just taking the product measures of the Disintegration's of $\pi^{2,3}$ and $\pi^{1,2}$ and "Gluing" them together with μ_2 .

Triangle inequality

Let $\mu_1, \mu_2, \mu_3 \in \mathcal{P}(X)$, $\pi^{1,2}$ and $\pi^{2,3}$ be optimal. Let $\pi \in \mathcal{P}(X \times X \times X)$ be the measure from Dudley and $p_{\#}^{1,3}(\pi) = \pi^{1,3} \in \Gamma(\mu_1, \mu_3)$.

$$W_2(\mu_1, \mu_3) \leq \left(\int_{X \times X} d^2(x_1, x_3) d\pi^{1,3}(x_1, x_3) \right)^{1/2}$$

Triangle inequality

Let $\mu_1, \mu_2, \mu_3 \in \mathcal{P}(X)$, $\pi^{1,2}$ and $\pi^{2,3}$ be optimal. Let $\pi \in \mathcal{P}(X \times X \times X)$ be the measure from Dudley and $p_{\#}^{1,3}(\pi) = \pi^{1,3} \in \Gamma(\mu_1, \mu_3)$.

$$\begin{aligned} W_2(\mu_1, \mu_3) &\leq \left(\int_{X \times X} d^2(x_1, x_3) d\pi^{1,3}(x_1, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X \times X} d^2(x_1, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \end{aligned}$$

Triangle inequality

Let $\mu_1, \mu_2, \mu_3 \in \mathcal{P}(X)$, $\pi^{1,2}$ and $\pi^{2,3}$ be optimal. Let $\pi \in \mathcal{P}(X \times X \times X)$ be the measure from Dudley and $p_{\#}^{1,3}(\pi) = \pi^{1,3} \in \Gamma(\mu_1, \mu_3)$.

$$\begin{aligned} W_2(\mu_1, \mu_3) &\leq \left(\int_{X \times X} d^2(x_1, x_3) d\pi^{1,3}(x_1, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X \times X} d^2(x_1, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\leq \left(\int_{X \times X \times X} [d(x_1, x_2) + d(x_2, x_3)]^2 d\pi(x_1, x_2, x_3) \right)^{1/2} \end{aligned}$$

Triangle inequality

Let $\mu_1, \mu_2, \mu_3 \in \mathcal{P}(X)$, $\pi^{1,2}$ and $\pi^{2,3}$ be optimal. Let $\pi \in \mathcal{P}(X \times X \times X)$ be the measure from Dudley and $p_{\#}^{1,3}(\pi) = \pi^{1,3} \in \Gamma(\mu_1, \mu_3)$.

$$\begin{aligned} W_2(\mu_1, \mu_3) &\leq \left(\int_{X \times X} d^2(x_1, x_3) d\pi^{1,3}(x_1, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X \times X} d^2(x_1, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\leq \left(\int_{X \times X \times X} [d(x_1, x_2) + d(x_2, x_3)]^2 d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\stackrel{*}{\leq} \left(\int_{X \times X \times X} d^2(x_1, x_2) d\pi(x_1, x_2, x_3) \right)^{1/2} + \left(\int_{X \times X \times X} d^2(x_2, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \end{aligned}$$

Triangle inequality

Let $\mu_1, \mu_2, \mu_3 \in \mathcal{P}(X)$, $\pi^{1,2}$ and $\pi^{2,3}$ be optimal. Let $\pi \in \mathcal{P}(X \times X \times X)$ be the measure from Dudley and $p_{\#}^{1,3}(\pi) = \pi^{1,3} \in \Gamma(\mu_1, \mu_3)$.

$$\begin{aligned} W_2(\mu_1, \mu_3) &\leq \left(\int_{X \times X} d^2(x_1, x_3) d\pi^{1,3}(x_1, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X \times X} d^2(x_1, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\leq \left(\int_{X \times X \times X} [d(x_1, x_2) + d(x_2, x_3)]^2 d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\stackrel{*}{\leq} \left(\int_{X \times X \times X} d^2(x_1, x_2) d\pi(x_1, x_2, x_3) \right)^{1/2} + \left(\int_{X \times X \times X} d^2(x_2, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X} d^2(x_1, x_2) d\pi^{1,2}(x_1, x_2) \right)^{1/2} + \left(\int_{X \times X} d^2(x_2, x_3) d\pi^{2,3}(x_2, x_3) \right)^{1/2} \end{aligned}$$

Triangle inequality

Let $\mu_1, \mu_2, \mu_3 \in \mathcal{P}(X)$, $\pi^{1,2}$ and $\pi^{2,3}$ be optimal. Let $\pi \in \mathcal{P}(X \times X \times X)$ be the measure from Dudley and $p_{\#}^{1,3}(\pi) = \pi^{1,3} \in \Gamma(\mu_1, \mu_3)$.

$$\begin{aligned} W_2(\mu_1, \mu_3) &\leq \left(\int_{X \times X} d^2(x_1, x_3) d\pi^{1,3}(x_1, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X \times X} d^2(x_1, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\leq \left(\int_{X \times X \times X} [d(x_1, x_2) + d(x_2, x_3)]^2 d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\stackrel{*}{\leq} \left(\int_{X \times X \times X} d^2(x_1, x_2) d\pi(x_1, x_2, x_3) \right)^{1/2} + \left(\int_{X \times X \times X} d^2(x_2, x_3) d\pi(x_1, x_2, x_3) \right)^{1/2} \\ &\stackrel{!}{=} \left(\int_{X \times X} d^2(x_1, x_2) d\pi^{1,2}(x_1, x_2) \right)^{1/2} + \left(\int_{X \times X} d^2(x_2, x_3) d\pi^{2,3}(x_2, x_3) \right)^{1/2} \\ &= W_2(\mu_1, \mu_2) + W_2(\mu_2, \mu_3). \end{aligned}$$

Comparing L^2 norm and Wasserstein distance

Let $\nu \ll \lambda^n$ with Radon Nikodym derivative f such that $\text{supp}(f) \subseteq \overline{B_1}$.

Additionally, let $\mu_h \ll \lambda^n$ such that $f_h(x) = f(x + h)$. Then we have

$$\|f_h - f\|_{L^2(\mathbb{R}^n)}^2 = 2\|f\|_{L^2(\mathbb{R}^n)}^2$$

but on the other hand

$$W_2(\mu_h, \mu) = \|h\|_2$$

which for large $\|h\|$ implies

$$W_2(\mu_h, \mu) \gg \|f_h - f\|_{L^2(\mathbb{R}^n)}.$$

In case of small $\|h\|$ we can also find $W_2(\mu_h, \mu) \ll \|f_h - f\|_{L^2(\mathbb{R}^n)}$.

Wasserstein distance for normal distributions

For any $\mu_X, \mu_Y \in \mathcal{P}_2(\mathbb{R}^d)$ with $\mathcal{L}(X) = \mu_X$ and $\mathcal{L}(Y) = \mu_Y$ where X and Y are normally distributed, the Wasserstein distance can be calculated as follows

$$W_2(\mu_X, \mu_Y) = \|m_X - m_Y\|^2 + \text{tr} \left(\Sigma_X - 2 \cdot \left(\Sigma_Y^{1/2} \Sigma_X \Sigma_Y^{1/2} \right)^{1/2} + \Sigma_Y \right)$$


If Σ_X and Σ_Y commute, this simplifies to

$$W_2(\mu_X, \mu_Y) = \|m_X - m_Y\|^2 + \sum_{i=1}^d \left(\sqrt{\lambda_i^X} - \sqrt{\lambda_i^Y} \right)^2$$

where λ_i^X and λ_i^Y are the Eigenvalues of Σ_X and Σ_Y respectively.

Normally distributed examples

Assume $\Sigma_X = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $\Sigma_Y = \begin{pmatrix} \frac{1}{2} & \frac{1}{3} \\ \frac{1}{3} & \frac{1}{2} \end{pmatrix}$ with $m_X = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ and $m_Y = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$.



Iterated Dudley Lemma

We want to represent a sequence of measures (μ_n) as the marginals of some measure in an infinite product space.

For that we need the iterated Dudley Lemma. Note that we also allow $N = \infty$ in this Lemma, in that case, the \leq become $<$.

Lemma

Let $N \geq 3$ and for any $n \leq N$ (X_n, d_n) Polish, $\mu_n \in \mathcal{P}(X_n)$ and $\theta_n \in \Gamma(\mu_{n-1}, \mu_n)$. Then there exists $\pi_n \in \mathcal{P}(X_1 \times \dots \times X_n)$ for any $n \leq N$ such that.

1. $p_{\#}^{1, \dots, n-1} \pi_n = \pi_{n-1}$ for $2 \leq n \leq N$
2. $p_{\#}^i \pi_n = \mu_i$ for $1 \leq i \leq n \leq N$
3. $p_{\#}^{i-1, i} \pi_n = \theta_i$ for $2 \leq i \leq n \leq N$

Proof is just applying the Dudley Lemma iteratively (hence the name :).

Lifting completeness from X to $\mathcal{P}_p(X)$

For $N = \infty$ we get a sequence (π_n) and can find a unique measure π_∞ on $\mathbb{X} = \prod_{i=1}^{\infty} X_i$ such that

$$p_{\#}^{(1, \dots, n)} \pi_\infty = \pi_n.$$

Additionally, we remind ourselves of the metric version of the L^p spaces.

$$L^p(\Omega, \mathcal{F}, P, X) := \left\{ f : \Omega \rightarrow X \mid f \text{ measurable, } \int_{\Omega} d^p(f, z_0) dP < \infty \right\}$$

with

$$d_{L^p}^p(f, g) := \int_{\Omega} d^p(f, g) dP$$

These are not necessarily vector spaces, but one can show that this more general notion of L^p space is still complete. (Which we will need in the following)

Theorem

Let (X, d) be a complete metric space, then $(\mathcal{P}_p(X), W_p)$ is complete.

Duality for the Wasserstein Distance

By the Kantorovich-Rubinstein Duality we have

$$\begin{aligned} W_1(\mu, \nu) &= \sup_{(\phi, \psi) \in I_c} \left\{ \int_X \phi d\mu + \int_X \psi d\nu \right\} \\ &= \sup_{\|\phi\|_{Lip} \leq 1} \left\{ \int_X \phi d\mu + \int_X \phi^c d\nu \right\} \\ &= \sup_{\|\phi\|_{Lip} \leq 1} \left\{ \int_X \phi d\mu - \int_X \phi d\nu \right\} \end{aligned}$$

where the first equality holds for any W_p distance, the latter need that $c = d$.

Wasserstein distance for $p = \infty$

If we restrict ourselves to $\mathcal{P}_\infty(X)$, the space of probability measures with bounded support. We can also define W_∞ as the limit of W_p .

$$\begin{aligned}\lim_{p \rightarrow \infty} W_p(\mu, \nu) &= W_\infty(\mu, \nu) = \inf\{\|d(x, y)\|_{L^\infty}(\pi) \mid \pi \in \Gamma(\mu, \nu)\} \\ &= \inf_{\pi \in \Gamma(\mu, \nu)} \inf\{C \geq 0 \mid |d(x, y)| \leq C \text{ for } \pi \text{ almost all } (x, y)\}\end{aligned}$$

and we have

$(\mathcal{P}_\infty(X), W_\infty)$ is a metric space